


# ARCHIVES

OF

# FAMILY MEDICINE

FEBRUARY 1993

PROPERTY OF  
BRANDYWINE HOSPITAL  
& TRAUMA CENTER  
HEALTH SCIENCES LIBRARY



Electron photomicrograph of didanosine (DDI), second drug approved for combating the AIDS virus. See page 158.

---

## CLINICAL FOLLOW-UP OF CERVICAL SAMPLING WITH THE AYRE SPATULA AND ZELSMYR CYTOBRUSH

B. J. Crouse, B. A. Elliott, N. Nesin

---

## PREDICTING EXERCISE AND SMOKING BEHAVIORS IN DIABETIC AND HYPERTENSIVE PATIENTS: AGE, RACE, SEX, AND PSYCHOLOGICAL FACTORS

J. G. Spangler, J. C. Konen

---

## IMMOBILITY IN THE AGED

P. Rousseau

---

## UNPROVEN DIET THERAPIES IN THE TREATMENT OF THE CHRONIC FATIGUE SYNDROME

D. H. Morris, F. J. Stare

---

## HEPATIC ABSCESS PRESENTING AS SEVERE FATIGUE AND ANEMIA

V. Nguyen, J. P. Richardson

---

## GUIDELINES FOR THE MEDICAL MANAGEMENT OF THE HOME-CARE PATIENT

American Medical Association  
Home Care Advisory Panel

American Medical Association

Physicians dedicated to the health of America



#BXNBBP\*\*\*\*\*SCF 189  
#00002894416# 97 32 21 DEC93  
BRANDYWINE HOSPITAL  
3 TRAUMA CENTER  
HEALTH SCIENCE LIBRARY  
ROUTE 30 BYPASS  
COATESVILLE PA  
19320

**A**LTHOUGH IMMOBILITY is not a disease in the traditional sense, with a specific identifiable cause and a specific cure, it has a powerful influence on the outcome of many diseases in elderly patients. Understanding the effects of immobility gives us an opportunity to make a real difference for many of our patients.

We have all had the experience of "curing" a case of pneumonia in an elderly patient only to find that he or she never returns to his or her previous level of overall function and independence. Fortunately, as this article points out, much of that functional decline is potentially preventable.

How can we prevent the complications of immobility? First, we must remember to assess mobility in elderly patients. After some disappointing outcomes, I have learned to spend extra time both asking elderly patients about their functional abilities at home and getting them off the emergency department stretcher to stand and walk as part of the initial examination. Finally, we must mobilize the patient as early as possible. This includes passive mobility for the very ill, early ambulation for those who are able, and resistive exercise when appropriate.

Dan Brewer, MD  
Maryville, Tenn