

PHOTODEHYDROXYLATION OF SOME 3-HYDROXYOXINDOLES¹


Noboru Shoji, Yoshikazu Kondo*, and Tsunematsu Takemoto

Pharmaceutical Institute, Tohoku University, Aobayama, Sendai, Japan

Photoreduction of 3-substituted 3-hydroxyoxindoles I-IV in the presence of sodium hydrosulfite led to loss of the 3-hydroxyl group. Irradiation at longer wavelengths ($\lambda > 3000 \text{ \AA}$) did not cause this photodehydroxylation.

While reduction of simple 3-hydroxyoxindoles usually gave the corresponding oxindoles, 3-alkyl-3-hydroxyoxindoles sometimes produced the 3-alkylindoles together with the oxindoles.² Generalization on the reduction of the 3-hydroxyoxindoles was so far unknown. We now wish to report that irradiation of some 3-substituted 3-hydroxyoxindoles I-IV in aqueous solution in the presence of sodium hydrosulfite resulted in dehydroxylation of the 3-hydroxyl group (Scheme 1).

Scheme 1.



I: R=COCH₃, X=H

II: R=C(NO₂)CH₃, X=H

III: R=Py, X=H

IV: R=Py, X=Cl


V: R=COCH₃, X=H

VI: R=C(NO₂)CH₃, X=H

VII: R=Py, X=H

VIII: R=Py, X=Cl


In a typical run, a 10^{-2} M aqueous solution of 3-substituted 3-hydroxyoxindole in the presence of 10^{-1} M sodium hydrosulfite was irradiated at ambient temperature with a 450 W medium-pressure lamp (Ushio UM-452, 1.9 W at 2894 Å) equipped with a Vycor filter. After irradiation for 10 h, most of the starting material disappeared by monitoring silica gel thin-layer chromatography. Isolation by column chromatography gave a major photoproduct³ (25-43% yield) which was confirmed to be 3-substituted oxindole (V-VIII) by direct comparison with an independently prepared sample in the following manner (Scheme 2).



Irradiation with light above 3000 Å did not cause dehydroxylation. Nmr data and melting points of these products were shown in Table I. The oxindole (V-VIII) was converted to the corresponding 3-hydroxyoxindole by aerial oxidation.⁴ When sodium borohydride⁵ was used in the place of sodium hydrosulfite, irradiation of the 3-hydroxyoxindole (I-IV) produced a mixture of several unidentified substances.

Addition of the low energy triplet-quencher such as biacetyl⁶ gave no apparent effect on this photodehydration. A reasonable mechanism involve a reduced intermediate X *via* zwitterion IX as shown in Scheme 3.


Table I*



	mp	δ_{CH}	δ_{CHH}	δ_{CHH}
V	99-101°	3.86 (q, $J_1=4.2$ Hz, $J_2=8.2$ Hz)	2.83 (q, $J_1=8.2$ Hz, $J_2=17.5$ Hz)	3.27 (q, $J_1=4.2$ Hz, $J_2=17.5$ Hz)
VI	175° (decomp.)	3.69 (q, $J_1=4.7$ Hz, $J_2=8$ Hz)	2.43 (q, $J_1=8$ Hz, $J_2=15.3$ Hz)	2.87 (q, $J_1=4.7$ Hz, $J_2=15.3$ Hz)
VII	130°	4.16 (q, $J_1=5.2$ Hz, $J_2=9$ Hz)	3.12 (q, $J_1=9$ Hz, $J_2=15$ Hz)	3.64 (q, $J_1=5.2$ Hz, $J_2=15$ Hz)
VIII	170-172°	4.15 (q, $J_1=5$ Hz, $J_2=8.7$ Hz)	3.08 (q, $J_1=8.7$ Hz, $J_2=15$ Hz)	3.60 (q, $J_1=5$ Hz, $J_2=15$ Hz)

* Nmr spectra were recorded on a Hitachi R-20 spectrometer (60 MHz) and the chemical shifts are in ppm from TMS.

Scheme 3



REFERENCES

- 1 On Photoreduction. Part V. Part IV: *Tetrahedron*, 1972, 28, 797.
- 2 B. Robinson, *Chem. Rev.*, 1969, 69, 786.
- 3 All compounds gave satisfactory elemental analyses.
- 4 L. P. Julian and J. Píkl, *J. Amer. Chem. Soc.*, 1935, 57, 539; S. Toyoshima, N. Hirose, K. Yamatsu, and S. Sohda, *Yakugaku Zasshi*, 1970, 90, 1524; N. Hirose, S. Sohda, and S. Toyoshima, *Yakugaku Zasshi*, 1971, 91, 1323.
- 5 Cf. Y. Kondo, *J. Synth. Org. Chem. Japan*, 1971, 29, 1109.
- 6 W. A. Noyes, Jun., G. B. Porter, and J. E. Jolley, *Chem. Rev.*, 1958, 58, 49.

Received, 22nd November, 1973