

HETEROCYCLES, Vol. 74, 2007, pp. 961 - 968. © The Japan Institute of Heterocyclic Chemistry
Received, 7th August, 2007, Accepted, 23rd October, 2007, Published online, 26th October, 2007. COM-07-S(W)26

ONE-POT SYNTHESIS OF DIBENZOFURAN-1,4-DIONES

Tetsuya Takeya,* Hiromu Kondo, Kazuho Tomita, Iwao Okamoto,
Nobuyoshi Morita, and Osamu Tamura*

Showa Pharmaceutical University, 3-3165 Higashi-tamagawagakuen, Machida,
Tokyo 194-8543, Japan. E-mail: tamura@ac.shoyaku.ac.jp

Dedicated to Professor Dr. Ekkehard Winterfeldt on the occasion of his 75th birthday

Abstract – One-pot synthesis of dibenzofuran-1,4-diones **8** from 4-methoxyphenols (or 4-methoxy-1-naphthols) **4** was achieved by oxidative dimerization over a semiconductor in heated, O₂-saturated toluene, followed by selective monodemethylation, and oxidative cyclization.

INTRODUCTION

The dibenzofuran-1,4-dione core **1** is frequently found in natural products, such as popolohuanone E (**2**) and violet-quinone (**3**) (Figure 1),¹⁻⁴ and several methods have been reported for efficient construction of this structure.⁵⁻⁸ In this context, we recently disclosed an efficient synthetic method of dibenzofuran-1,4-diones **8** having an oxygen functionality at C8, featuring a three-step sequence of oxidative dimerization of hydroquinone monomethyl ethers **4** leading to 2,2'-biarenyl-1,1'-quinones **6** via the 2,2'-biarenols **5**, followed by mono-demethylation of **6**, and oxidative cyclization of the quinone-arenols **7** (Scheme 1).⁹





Figure 1

If the three-step sequence could be conducted in one pot, it would be very convenient for synthesizing a range of compounds **8** for studies on structure-activity relationships. We therefore sought appropriate oxidative conditions, and found that one-pot synthesis of several dibenzofuran-1,4-diones **8** from **4** could be achieved by employing a semiconductor and molecular oxygen.



RESULTS AND DISCUSSION

We recently reported that semiconductors, such as activated carbon, ZrO_2 , Nb_2O_5 , TiO_2 , SnO_2 , and Ag_2O , could be used for dimerization of 1-naphthols **4**, leading to the dimers **5** or **6**, and that SnO_2 effected mono-demethylation of **6**.¹⁰ We initiated our investigation by examining the reaction of the naphthol **4a** with SnO_2 in the presence of various oxidants (Table 1). Since benzoquinone was found to be an excellent oxidant for dimerization of **4a** to **6a**,¹¹ SnO_2 and benzoquinone were used in combination. Heating a mixture of **4a**, SnO_2 , and benzoquinone in toluene in a sealed tube at 100 °C gave a mixture of





Table 1. One-pot synthesis of dibenzofuranquinone **8a** from **4a**.

Entry	Conditions	Product (% yield)		
		5a	6a	8a
1	SnO_2 / benzoquinone (1.2 equiv.) toluene, 100 °C, 7 h	30	26	11
2	SnO_2 / air, toluene, 100 °C, 21 h			36
3	SnO_2 / O_2 , toluene, 100 °C, 7 h			62


three products **5a**, **6a**, and the desired quinone **8a** in 30, 26, and 11% yields, respectively (entry 1). When the oxidation reaction was conducted in air-saturated toluene, the yield of **8a** was slightly improved (entry 2). The best result was obtained by the use of oxygen-saturated toluene as the solvent. Thus, on heating with SnO_2 in oxygen-saturated toluene, the naphthol **4a** underwent oxidative dimerization followed by cyclization to afford **8a** in 62% yield (entry 3).

Next, reaction of the naphthol **4b** was examined. Since the use of conditions similar to those employed for the reaction of **4a** (SnO_2/O_2 , toluene, 100 °C) gave a complex mixture, other semiconductors such as TiO_2 , Nb_2O_5 , and ZrO_2 were examined. Among them, ZrO_2 afforded **8b** in 19% yield, along with **5b** (8%) and unidentified products (Scheme 2).



Scheme 2

One-pot oxidative dimerization-cyclization sequences of 4-methoxyphenols were also examined. Unfortunately, treatment of the phenol **4c** with SnO_2 in O_2 -saturated toluene at 100 °C again resulted in the formation of a complex mixture. After extensive experimentation, the use of Nb_2O_5 in place of SnO_2 was found to afford the quinone **8c** in 21% yield although unidentified by-products were still produced (Scheme 3).



Scheme 3

In the case of **4d**, treatment with SnO_2 in O_2 -saturated toluene at 100 °C gave the dibenzofuran-1,4-dione **8d** in 15% yield, along with unidentified products (Table 2, entry 1). Combined use of ZrO_2 and molecular oxygen appeared to be a milder oxidation system, and the starting **4d** was recovered despite the prolonged reaction time (entry 2). Addition of K_2CO_3 shortened the reaction time and improved the yield of **8d**, probably due to activation of substrate by conversion of the phenol **4d** to phenoxide (entry 3).

Finally, we attempted a one-pot synthesis of the dibenzofuran-1,4-dione **8e**, a model compound of popolohuanone E,^{6b,9} from the phenol **4e**. However, the use of ZrO₂, Nb₂O₅, TiO₂, SnO₂, and Ag₂O gave **8e** only in very low yield (<10%) although the reason remains unclear. For example, heating **4e** with ZnO₂ in O₂-saturated toluene at 100 °C gave only 6% yield of **8e** accompanied by many unidentified products.





Table 2. One-pot synthesis of dibenzofuranquinone **8d** from **4d**.

Entry	Conditions	Product (% yield)	
		5d	8d
1	SnO ₂ / O ₂ , toluene, 100 °C, 7 h		15
2	ZrO ₂ / O ₂ , toluene, 100 °C, 3 days	18	15
3	ZrO ₂ / O ₂ , K ₂ CO ₃ , toluene, 100 °C, 19 h	4	32



In summary, we examined the feasibility of one-pot synthesis of dibenzofuran-1,4-diones **8a-e** from 4-methoxyphenols (or 4-methoxy-1-naphthols) **4a-e** and found that the use of a combination of a semiconductor and molecular oxygen afforded the desired product **8** in some cases.

EXPERIMENTAL

General

Melting points are uncorrected. IR spectra were recorded with a Shimadzu FTIR-3200A spectrophotometer. ¹H and ¹³C NMR spectra were measured on JEOL JNM-AL300 and JEOL JNM-α-500 spectrometers. High-resolution mass spectra (HRMS) were obtained with a JEOL JMS-D300 or a JEOL JMS-HX110 instrument. Column chromatography was performed on Silica gel 60 (0.040-0.063 mm) under pressure.

4,4'-Dimethoxy[2,2']binaphthalenyl-1,1'-diol (5a),

4,4'-Dimethoxy[2,2']binaphthalenyldiene-1,1'-dione (6a),

5-Methoxydinaphtho[1,2-b;2',3'-d]furan-7,12-dione (8a) (Table 1).

Entry 1: Argon gas was bubbled into a solution of 4-methoxy-1-naphthol (**4a**) (50 mg, 0.29 mmol)

in toluene (15 mL) until saturation. Stannic oxide (5 g) and 1,4-benzoquinone (38 mg, 0.35 mmol) were added to the solution, and the mixture was heated in a sealed tube at 100 °C for 7 h, then filtered, and the filtrate was concentrated under reduced pressure. The residue was chromatographed on silica gel with CH₂Cl₂-hexane (1:1) to give **5a** (15 mg, 30%), **6a** (13 mg, 26%), **8a** (5.2 mg, 11%). **5a**: Colorless needles, mp 223.0-224.0 °C (benzene). IR (KBr): 3435 cm⁻¹; ¹H-NMR (CDCl₃) δ 3.99 (6H, s), 5.44 (2H, s), 6.73 (2H, s), 7.58-7.61 (4H, m), 8.28-8.30 (4H, m); MS *m/z* 346 (M⁺); HRMS calcd for C₂₂H₁₈O₄ 346.1200, found 346.1222. **6a**: Deep blue needles, mp 257-258 °C (benzene); IR (KBr) 1605, 1585 cm⁻¹; ¹H-NMR (CDCl₃) δ 4.08 (6H, s), 7.48 (2H, t, *J* = 7.7 Hz), 7.61 (2H, t, *J* = 7.7 Hz), 7.79 (2H, d, *J* = 7.7 Hz), 8.17 (2H, d, *J* = 7.7 Hz), 8.42 (2H, s); MS *m/z* 344 (M⁺); HRMS calcd for C₂₂H₁₈O₄ 344.1044, found 344.1029. **8a**: ¹H NMR (CDCl₃) δ 4.12 (3H, s), 7.48 (1H, s), 7.61-7.80 (4H, m), 8.22-8.45 (4H, m); ¹³C NMR (CDCl₃) δ 56.2, 95.9, 119.3, 121.0, 121.5, 123.6, 125.3, 126.7, 126.97, 127.03, 127.6, 128.1, 132.9, 133.4, 133.86, 133.90, 149.1, 152.6, 155.0, 174.4, 182.1. These ¹H and ¹³C NMR spectra were consistent with those reported.⁹

Entry 3: Oxygen gas was bubbled into a solution of 4-methoxy-1-naphthol (**4a**) (50 mg, 0.29 mmol) in toluene (15 mL) until saturation. Stannic oxide (5 g) was added to the solution, and the mixture was heated in a sealed tube at 100 °C for 7 h, then filtered, and the filtrate was concentrated under reduced pressure. The residue was chromatographed on silica gel with CH₂Cl₂-hexane (1:1) to give **8a** (29.5 mg, 62%).

4,4',8,8'-Tetramethoxy[2,2']binaphthalenyl-1,1'-diol (5b) and 1,5,8-trimethoxydinaphtho-[1,2-*b*;2',3'-*d*]furan-7,12-dione (8b).

Using a procedure similar to that for **8a** (Table 1, entry 3), 4,8-dimethoxy-1-naphthol (**4b**) (50 mg, 0.25 mmol) was treated with zirconium oxide (5 g) in oxygen-saturated toluene (15 mL). After work-up, the crude material was chromatographed on silica gel with CH₂Cl₂ to give **5b** (4.0 mg, 8%) and **8b** (9.2 mg, 19%). **5b**: Colorless needles, mp 207.0-209.0 °C (CHCl₃-hexane); IR (KBr) 3350 cm⁻¹; ¹H NMR (CDCl₃) δ 3.95 (6H, s), 4.03 (6H, s), 6.87 (2H, d, *J* = 8.6 Hz), 6.92 (2H, s), 7.35 (2H, t, *J* = 8.6 Hz), 7.89 (2H, d, *J* = 8.6 Hz), 9.87 (2H, s); ¹³C NMR (CDCl₃) δ 56.0, 56.1, 105.2, 109.4, 115.7, 116.0, 119.4, 125.0, 127.6, 144.7, 147.6, 156.2. These ¹H and ¹³C NMR spectra were consistent with those reported.^{10b} **8b**: IR (KBr) 1665, 1580 cm⁻¹; ¹H NMR (CDCl₃) δ 4.081 (3H, s), 4.085 (3H, s), 4.16 (3H, s), 7.10 (1H, d, *J* = 7.9 Hz), 7.35 (1H, d, *J* = 8.2 Hz), 7.55 (1H, dd, *J* = 7.9, 8.2 Hz), 7.67 (1H, s), 7.70 (1H, dd, *J* = 7.3, 8.2 Hz), 7.96 (2H, m); ¹³C NMR (CDCl₃) δ 56.1, 56.2, 56.5, 97.1, 107.6, 113.7, 115.4, 118.4, 119.9, 120.5, 120.7, 125.6, 127.8, 128.7, 134.9, 135.7, 148.7, 150.9, 154.4, 155.4, 160.4, 173.6, 182.3. These ¹H and ¹³C NMR spectra were consistent with those reported.⁹

3,7,8-Trimethoxydibenzofuran-1,4-dione (8c).

Using a procedure similar to that for **8a** (Table 1, entry 3), 3,4-dimethoxyphenol (**4c**) (50 mg, 0.32 mmol) was treated with niobium(V) oxide (5 g) in oxygen-saturated toluene (15 mL). After work-up, the crude material was chromatographed on silica gel with CH₂Cl₂-AcOEt (10:1) to give **8c** (10.4 mg, 21%). IR (KBr) 1680, 1645, 1605, 1560 cm⁻¹; ¹H NMR (CDCl₃) δ 3.90 (3H, s), 3.98 (3H, s), 3.99 (3H, s), 5.86 (1H, s), 7.07 (1H, s), 7.47 (1H, s); ¹³C NMR (CDCl₃) δ 56.47, 56.53, 57.0, 95.2, 102.5, 106.8, 114.9, 123.7, 149.5, 149.7, 152.8, 153.2, 159.8, 170.1, 183.5. These ¹H and ¹³C NMR spectra were consistent with those reported.⁹

3,3'-di-tert-Butyl-5,5'-dimethoxybiphenyl-2,2'-diol (5d) and 2,6-Di-tert-butyl-8-methoxydibenzofuran-1,4-dione (8d) (Table 2, entry 3).

Using a procedure similar to that for **8a** (Table 1, entry 3), 3-tert-butyl-4-hydroxyanisole (**4d**) (50 mg, 0.28 mmol) was treated with zirconium oxide (5 g) and potassium carbonate (383 mg, 2.8 mmol) in oxygen-saturated toluene (15 mL). After work-up, the crude material was chromatographed on silica gel with CH₂Cl₂-AcOEt (8:1) to give **5d** (2 mg, 4%) and **8d** (15 mg, 32%). **5d**: Colorless powder, mp 228.5-229.5 °C (CH₂Cl₂); IR (KBr) 3420 cm⁻¹; ¹H NMR (CDCl₃) δ 1.43 (18H, s), 3.78 (6H, s), 5.01 (2H, s), 6.63 (2H, d, *J* = 3.1 Hz), 6.96 (2H, d, *J* = 3.1 Hz); ¹³C NMR (CDCl₃) δ 27.5, 33.2, 53.8, 109.7, 113.3, 121.2, 136.9, 143.9, 151.2; MS (*m/z*) 358 (M⁺); HRMS calcd for C₂₄H₂₂O₆ 358.2136, found 358.2133. **8d**: ¹H NMR (CDCl₃) δ 1.38 (9H, s), 1.49 (9H, s), 3.90 (3H, s), 6.60 (1H, s), 7.04 (1H, d, *J* = 2.6 Hz), 7.45 (1H, d, *J* = 2.6 Hz); ¹³C NMR (CDCl₃) δ 29.7, 34.6, 35.6, 55.9, 100.5, 116.9, 123.4, 124.1, 131.0, 137.9, 150.1, 151.0, 156.9, 158.3, 177.6, 184.5. These ¹H and ¹³C NMR spectra were consistent with those reported.⁹

2,6-Bis-cyclohexylmethyl-3,7,8-trimethoxydibenzofuran-1,4-dione (8e).

Using a procedure similar to that for **8a** (Table 1, entry 3), 2-cyclohexylmethyl-3,4-dimethoxyphenol (**4e**) (50 mg, 0.20 mmol) was treated with zirconium oxide (5 g) oxygen-saturated toluene (15 mL). After work-up, the crude material was chromatographed on silica gel with hexane-AcOEt (20:1) to give **8e** (2.7 mg, 6%). ¹H NMR (CDCl₃) δ 0.95-1.75 (22H, m), 2.43 (2H, d, *J* = 7.2 Hz), 2.81 (2H, d, *J* = 7.3 Hz), 3.88 (3H, s), 3.95 (3H, s), 4.08 (3H, s), 7.41 (1H, s). The signals were consistent with reported data.⁹

ACKNOWLEDGEMENTS

This research was supported by a Grant-in-aid from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

REFERENCES (AND NOTES)

1. For popolohuanone E, see: J. R. Carney and P. J. Scheuer, *Tetrahedron Lett.*, 1993, **34**, 3727.
2. For violet-quinone, see: A. V. B. Sankaram, V. V. N. Reddy, and G. S. Sidhu, *Phytochemistry*, 1981, **20**, 1093.
3. For other natural products containing a dibenzofuran-1,4-dione core, see: (a) K. Ishiguro, Y. Ohira, and H. Oku, *J. Nat. Prod.*, 1998, **61**, 1126. (b) M. Xu, Z. Deng, M. Li, J. Li, H. Fu, P. Proksch, and W. Lin, *J. Nat. Prod.*, 2004, **67**, 762. (c) C. Puder, K. Wagner, R. Vettermann, R. Hauptmann, and O. Potterat, *J. Nat. Prod.*, 2005, **68**, 323. (d) O. Kayser, A. F. Kiderlen, H. Laatsch, and S. L. Croft, *Acta Tropica*, 2000, **77**, 307.
4. For biological activities of synthetic compounds containing a dibenzofuran-1,4-dione core, see: (a) C. C. Cheng, Q. Dong, D. F. Liu, Y. L. Luo, L. F. Liu, A. Y. Chen, C. Yu, N. Savaraj, and T. C. Chou, *J. Med. Chem.*, 1993, **36**, 4108. (b) H. A. M. Hejaz, L. W. L. Woo, A. Purohit, M. J. Reed, and B. V. L. Potter, *Bioorg. Med. Chem.*, 2004, **12**, 2759. (c) K. Lee, Y. Park, S. Park, J. Hwang, S. Lee, G. Kim, W. Park, S. Lee, D. Jeong, J. Kong, H. Kang, and H. Cho, *Bioorg. Med. Chem. Lett.*, 2006, **16**, 737.
5. For construction of **1** by cyclization of 2,2'-biquinones, see: (a) R. Buchan, and O. C. Musgrave, *J. Chem. Soc., Perkin Trans. 1*, 1980, 90. (b) T. Ogata, I. Okamoto, E. Kotani, and T. Takeya, *Tetrahedron*, 2004, **60**, 3941. (c) J. C. Anderson, R. M. Denton, and C. Wilson, *Org. Lett.*, 2005, **7**, 123.
6. For construction of **1** by coupling reaction of dichloro-quinones with phenols, see: (a) A. Shand and R. H. Thomson, *Tetrahedron*, 1963, **19**, 1919. (b) Y. Ueki, M. Itoh, T. Katoh, and S. Terashima, *Tetrahedron Lett.*, 1996, **37**, 5719. (c) T. Katoh, M. Nakatani, S. Shikita, R. Sampe, A. Ishiwata, O. Ohmori, M. Nakamura, and S. Terashima, *Org. Lett.*, 2001, **3**, 2701. See also ref. 4.
7. For construction of **1** by transition metal-mediated cyclization, see: (a) E. Martinez, L. Martinez, J. C. Estevez, R. J. Estevez, and L. Castedo, *Tetrahedron Lett.*, 1998, **39**, 2175. (b) H. Chang, T. Chou, N. Savaraj, L. F. Liu, C. Yu, and C. C. Cheng, *J. Med. Chem.*, 1999, **42**, 405. (c) E. Martinez, L. Martinez, M. Treus, J. C. Estevez, R. J. Estevez, and L. Castedo, *Tetrahedron*, 2000, **56**, 6023. (d) A. Martinez, M. Fernandez, J. C. Estevez, R. J. Estevez, and L. Castedo, *Tetrahedron*, 2005, **61**, 1353. (e) J. M. Miguel del Corral, M. A. Castro, M. Gordaliza, M. L. Martin, A. M. Gamito, C. Cuevas, and A. San Feliciano, *Bioorg. Med. Chem.*, 2006, **14**, 2816.
8. For construction of **1** by lithiation-mediated cyclization, see: M. S. Azevedo, G. B. C. Alves, J. N. Cardoso, R. S. C. Lopes, and C. C. Lopes, *Synthesis*, 2004, 1262.
9. T. Takeya, H. Kondo, T. Otsuka, K. Tomita, I. Okamoto, and O. Tamura, *Org. Lett.*, 2007, **9**, 2807.
10. (a) T. Otsuka, I. Okamoto, E. Kotani, and T. Takeya, *Tetrahedron Lett.*, 2004, **45**, 2643. (b) T.

- Takeya, H. Doi, T. Ogata, T. Otsuka, I. Okamoto, and E. Kotani, *Tetrahedron*, 2004, **60**, 6295. (c) T. Takeya, T. Otsuka, I. Okamoto, and E. Kotani, *Tetrahedron*, 2004, **60**, 10681.
11. T. Takeya, H. Kondo, T. Otsuka, H. Doi, I. Okamoto, and E. Kotani, *Chem. Pharm. Bull.*, 2005, **53**, 199.