

CHLOROSULFONYL ISOCYANATE: A NOVEL REAGENT FOR THE SYNTHESIS OF HETEROCYCLES

Ahmed Kamal and Pralhad B. Sattur

Medicinal Chemistry & Natural Products Division
Regional Research Laboratory
Hyderabad 500 007, India

Abstract - Chlorosulfonyl isocyanate, a novel reagent employed towards the synthesis of heterocycles and related compounds, finds extensive use. An effort has been made in this review to catalogue and update its applications for the synthesis of nitrogen, oxygen and sulphur containing ring systems.

Contents

1. Introduction
2. β -Lactams
- 2.1 Azetidinones
- 2.2 4-Acyloxy-2-azetidinones
3. Oxetes
4. Diazetidinone-imine
5. Lactone
6. Oxazolones
7. 3-Oxo-2-isothiazolidine-1,1-dioxides
8. 5-Oxo- Δ^2 -triazoline
9. 2,3-Dihydro-3,4-oxadiazoles
10. Benzimidazole-2-one
11. Benzoisothiazole-1,1-dioxide
12. Tetrazolones
13. 2-Pyridone
14. 1,4-Cyclo-adducts
15. 1,3-Oxazin-2-ones
16. 1,3- Oxazine-2,4-diones
17. Uracils

18. 2(1H)-Quinazolinones
19. 1,3-Benzoxazin-2-ones
20. 1,3,5-Triazine-diones
21. 1,2,4-Benzothiadiazinones
22. 1,2,3-Oxathiazine-2,2-dioxides
23. 1,2,3-Benzoxathiazine-2,2-dioxides
24. 1,2,4,6-Thiatriazine-1,1-dioxide-3-ones
25. 2-Azepinone
26. 1,2,4,7-Thiatriazepine-3(2H)-one-1,1-dioxides and 2,1,3,5-Benzothia-triazepin-4(5H)-one-2,2-dioxides
27. 1,2,3,5(3H)-Benzoxathiadiazepine-2,2-dioxide-4(5H)-ones
28. 1,5,2,3(3H)-Benzodioxathiazepine-2,2-dioxide-4-ones
29. Azocine
30. Miscellaneous fused heterocycles
- 30.1 Triazolotriazolone
- 30.2 Triazolopyrimidine
- 30.3 Fused 1,2,4,6-Thiatriazine-1,1-dioxides
- 30.4 Thiatriazinobenzimidazole
- 30.5 Azabullvalene
31. Novel unusual heterocycles
32. Conclusion

1. Introduction

Chlorosulfonyl isocyanate (CSI), the most reactive isocyanate discovered by Graf 30 years ago, finds extensive use in the synthesis of various heterocyclic ring systems. Its chemistry and synthetic reactions have been the subject of many reviews¹⁻³. In view of its versatile nature and high degree of reactivity it was considered of interest to broaden the scope of its usefulness in the synthesis of heterocycles and also to update its applications in this review.


Reactions of CSI cited in literature have been broadly classified based mainly on the site of initial attack, i.e., type I generally involving the initial attack at the carbon of the isocyanate moiety, while type II deals with cycloadditions (2+2) across the isocyanate C=N bond and whereas type III involves nucleophilic additions at the sulfonyl group.

Though CSI has enabled numerous types of synthetic transformations which is evident from the rich chemistry of this reagent covering the literature through 1968, its usefulness in the synthesis

of a variety of heterocyclic compounds and often novel systems is of immense importance. Thus, the present review describes the applications of CSI for simple, small heterocycles to large, fused and unusual heterocycles.


2. β -Lactams

2.1 2-Azetidinones: 2-Azetidinones 3 are produced by the (2+2) cycloaddition of chlorosulfonyl isocyanate to a variety of unsaturated linkages⁴⁻⁷. Cycloaddition reactions of CSI with different types of olefins have been studied thoroughly⁸⁻¹¹. Although for the formation of β -lactam, a two-step mechanism, has been proposed, a concerted cycloaddition can also take place. There are citations^{6,9} of evidence for both the mechanisms.






Examples illustrating the degree of usefulness of chlorosulfonyl isocyanate for the synthesis of a variety of 2-azetidinones 4-16¹²⁻²² are as follows.




SCHEME I





SCHEME I (Contd.)





5




6






7



8



















9




10


SCHEME I (Contd.)


11

12

13

14

15

16


Reaction of CSI with conjugated or non-conjugated dienes also leads to *N*-chlorosulfonyl- β -lactams which on hydrolysis give 2-azetidinones^{10,23}.



2.2 4-Acyloxy-2-azetidinones: The reaction of CSI with heterosubstituted olefins such as vinyl acetates leads to 4-acyloxy-2-azetidinones²⁴ upon hydrolysis.




The 2-azetidinones obtained by the reaction of CSI with a variety of olefins like vinyl esters present ideal building blocks for the total synthesis of penicillin, cephalosporin and oxacephem class of antibiotics²⁵⁻³⁰. The general reaction scheme II is given below where the cyclization of the 2-azetidinones usually involves a Wittig-type of reaction via the intermediate **20**.



SCHEME II


3. Oxetes

The reaction of CSI with acetylenes such as 1-diethylamino-1-propyne at low temperatures leads to oxete-type system 22³¹.




4. Diazetidinone-imine

Reaction of CSI with dicyclohexylcarbodiimide gives diazetidinone-imine 23³². In this reaction the mode of addition plays an important role, i.e., only the addition of CSI to dicyclohexylcarbodiimide will lead to diazetidinone.




5. Lactone

The reaction of CSI with diphenylmethylenecyclopropane afforded ring-opened iminolactone 25³³. The proposed mechanism involves the initial formation of β -lactam, which could then open to dipolar intermediate 24, followed by ring opening of the cyclopropylcarbanyl cation and closure through oxygen.




6. Oxazolones

The facile reaction of CSI with the hydroxy group of 2-hydroxy ketones leads to O-carbamoyl ketones 26. These carbamates on heating to 175-250°C undergo ring closure which results in the formation of oxazolones 27³⁴.




7. 3-Oxo-2-isothiazolidine-1,1-dioxides

CSI reacts with olefins under free radical conditions to yield 2-chloroalkanesulfonyl isocyanates. Interestingly, an excess of olefin changes the course of reaction and 3-oxo-2-(2-chloroalkyl)-isothiazolidine-1,1-dioxides 28³⁵ are obtained in good yields.




8. 5-Oxo- Δ^2 -triazoline

Reaction of CSI with diazoacetic esters leads to an ester of 1-chlorosulfonyl-5-oxo- Δ^2 -triazoline-4-carboxylic acid 29¹.




9. 2,3-Dihydro-3,4-oxadiazoles

The reaction of CSI with 1,3-diarylnitrilimine gives exclusively 2-chlorosulfonylimino-3,5-diphenyl-2,3-dihydro-1,3,4-oxadiazole 30³⁶ via addition of the 1,3-dipole nitrilimine to the C=O function of CSI.




10. Benzimidazole-2-one

Reaction of CSI with 2-nitroaniline followed by hydrolysis leads to a urea derivative which on reductive cyclization yields benzimidazol-2-one 31³⁷.




11. Benzoisothiazole-1,1-dioxide

CSI reacts with non-enolizable ketones such as benzophenone to give a cyclized product, i.e., benzoisothiazole-1,1-dioxide 32³⁸ via a N-chlorosulfonylazomethine intermediate.




12. Tetrazolones

Reaction of CSI with alkylazides at ambient temperature gives alkylchlorosulfonyl tetrazolones which on hydrolysis afford stable alkyltetrazolones 33³⁹.




13. 2-Pyridone

Addition of CSI with dimethylphenylethylnamine has been reported to yield 2-pyridone 34⁴⁰.




14. 1,4-Cyclo-adducts

Though CSI reacts with conjugated 1,3-dienes to form N-chlorosulfonyl- β -lactams, these can be thermally rearranged to form 1,4-cyclo-adducts 35⁴¹.





The reaction of CSI with cyclohexa-1,3-diene yields N-chlorosulfonyl- β -lactam which opens to a dipolar intermediate and provides access to 1,4-addition products, 36 and 37⁴².




15. 1,3-Oxazin-2-ones

CSI reacts with α,β -unsaturated ketones to yield 3,4-dihydro-1,3-oxazin-2-ones. Hydrolysis produces the unsubstituted compounds 38⁴³.




16. 1,3-Oxazine-2,4-diones

Reaction of CSI with enolizable ketones by an electrophilic attack on the 2-methylenic carbon atom provides unstable N-chlorosulfonyl- β -ketocarboxamides. These, on reaction with excess of CSI followed by hydrolysis, yield 3,4-dihydro-2H-1,3-oxazine-2,4-diones 39^{44,45}. Further, 1,3-oxazine-2,4-diones can be easily converted to 5,6-disubstituted uracils 40⁴⁴.




17. Uracils

Recently it has been reported that the reaction of CSI with various 2-dialkylaminostyrenes gives 6-substituted dihydrouracils 41⁴⁶, i.e., by involving two equivalents of CSI.




The formation of uracils has been explained on the basis of a distinct 1,4-dipolar intermediate with opposing charges being stabilized by the dialkylamino group at the positive end of the sulfonyl chloride and by the carbonyl moiety at the negative end.




18. 2(1H)-Quinazolinones



The reaction of CSI with 2-aminoacetophenones/benzophenones leads to 2(1H)-quinazolinones 42⁴⁷ in good yields. This is a convenient alternate route of synthesis for 2(1H)-quinazolinones.




CSI reacts with methyl anthranilates to give urethanes upon hydrolysis which are cyclized in presence of a base to afford 2,4-quinazolinediones 43⁴⁸.

19. 1,3-Benzoxazin-2-ones


Reaction of CSI with 2-hydroxybenzaldehydes and 2-hydroxyacetophenones gives 1,3-benzoxazin-2-ones 44⁴⁹ whereas CSI reacts with 2-hydroxybenzophenones to yield O-carbamoyl compounds, and these are also cyclized in presence of a base to 44a.



Reaction of CSI with 2-hydroxybenzoates gives O-carbamoyloxybenzoates 45⁵⁰ which are cyclized in presence of a base to 1,3-benzoxazine-2,4-diones 46⁴⁸.




In a recent report CSI has been further employed for the synthesis of 1,3-benzoxazine-2,4-diones by another route⁵¹. Salicylic acid chloride is reacted with CSI, cyclized by the addition of base and hydrolyzed to afford 1,3-benzoxazine-2,4-dione as shown below.




20. 1,3,5-Triazine-diones


The reactions of CSI (2 moles) with compounds having carbon-nitrogen double bonds give triazine-dione systems. Thus, CSI reacts with azomethines to produce 1,3,5-triazine-diones 47³⁹ in the following way.



CSI also reacts with amidines to give amino-substituted triazine-diones 48⁵².




While addition of dialkylcarbodiimide to two moles of CSI affords triazinedioneimine 49³², in this case the mode of addition, i.e., the addition of carbodiimide to two moles of CSI, only leads to the formation of triazinedione system.




It is evident from the above reactions that CSI provides a convenient route to hexahydrotriazine-2,4-diones.

21. 1,2,4-Benzothiadiazinones

The reaction of CSI with aniline derivatives under Friedel-Crafts conditions leads to 2H-1,2,4-benzothiadiazin-3-one-1,1-dioxides 50⁵³.




Antihypertensive drug, diazoxide 51, can be prepared starting from 4-chloroaniline employing CSI under the Friedel-Crafts conditions as illustrated above and later heating with acetic acid. Thus CSI offers an alternative method of synthesis for this pharmaceutically important heterocyclic compound.




22. 1,2,3-Oxathiazine-2,2-dioxides


The reaction of CSI with many acetylenic compounds gives 1,2,3-oxathiazine-2,2-dioxides³¹. The ratio between the two isomers 52 and 53 depends on the nature of R^1 and R^2 substituents.



CSI reacts with enolizable ketones via α -methylenic carbon atom to produce N-chlorosulfonyl- β - ketocarboxamides 54^{44,54}. Excess of CSI in ether solvent acts as a Lewis acid in promoting the cyclization of 54 to 1,2,3-oxathiazin-4(3H)-one-2,2-dioxide compounds 56, probably through an N-sulfonylamine intermediate 55⁴⁵.




CSI also adds to the methylene group of t-butyl acetoacetate to give N-chlorosulfonyl adduct which upon thermolysis yields N-chlorosulfonylacetoacetamide 57 along with carbon dioxide and isobutylene. Then successive cyclodehalogenation gives an access to oxathiazinonedioxide 58⁵⁵.




23. 1,2,3-Benzoxathiazine-2,2-dioxides

Though CSI at room temperature reacts with phenolic compounds at carbon atom of its isocyanate function, at elevated temperature CSI reacts at the sulfur atom of the chlorosulfonyl group. Thus, the reactions of CSI with 2-hydroxybenzaldehyde, 2-hydroxyacetophenones or 2-hydroxybenzophenones at higher temperatures (100-105°C) undergo cycloaddition to yield 1,2,3-benzoxathiazine-2,2-dioxides 59⁵⁶.




Similarly, reaction of CSI with 2-hydroxybenzoates at higher temperatures (100-108°C) gives 2-sulfamoyloxybenzoates 60 which are cyclized in presence of a base to 1,2,3-benzoxathiazine-2,2-dioxides 61⁴⁸.




24. 1,2,4,6-Thiatriazine-1,1-dioxide-3-ones

Isothiureas react with CSI to afford substituted 1,2,4,6-thiatriazine-1,1-dioxide-3-ones ⁶²⁵⁷.




The reaction of CSI with 5-aminotetrazole and subsequent treatment with ethyldiisopropylamine gives the thiatriazine azide ⁵⁸. Treatment of this with water converted it to the urea derivative in which the tetrazole system was reconstituted. Thus, CSI provides an alternative synthesis for 5-tetrazolyl-ureas too.




25. 2-Azepinone

Addition of CSI to trans-2-phenylisopropenylcyclopropane yields N-chlorosulfonyl-2-azepinone ⁶³⁵⁹, the formation of which has been rationalized on the basis of the different stabilizations available to the dipolar intermediates.




26. 1,2,4,7-Thiatriazepin-3(2H)-one-1,1-dioxides and 2,1,3,5-benzothiatriazepin-4(5H)-one-2,2-dioxides

The reaction of 1,2-diamine with CSI gives 4,5,6,7-tetrahydro-1,2,4,7-thiatriazepin-3(2H)-one-1,1-dioxide 64 while CSI reacts with o-phenylenediamine to afford 1,3-dihydro-2,1,3,5-benzothiatriazepin-4(5H)-one-2,2-dioxide 65⁶⁰.




27. 1,2,3,5(3H)-Benzoxathiadiazepine-2,2-dioxide-4(5H)-ones

Addition of CSI to 2-aminophenols at higher temperature affords (3H)-1,2,3,5-benzoxathiadiazepine-2,2-dioxide-4(5H)-ones 66⁴⁸, a new class of seven-membered heterocycles.




28. 1,5,2,3(3H)-Benzodioxathiazepine-2,2-dioxide-4-ones

The reactions of CSI with catechols produce a novel family of seven-membered heterocycles 1,5,2,3(3H)-benzodioxathiazepine-2,2-dioxide-4-ones 67⁶¹.




29. Azocine

Treatment of 1,4-cyclohexadiene with CSI results in the formation of β -lactam which on O-methylation followed by monobromination and finally dehydrohalogenation yields 2-methoxy-1-azocine 68⁶². Thus this route employing CSI offers a direct and novel synthesis of azocine system.




30. Miscellaneous fused heterocycles



30.1 Triazolotriazolone: Azines may be considered as bis-anils which are versatile partners in cycloaddition reactions. The two azomethine groups may react with dipolarophiles in 'criss-cross' additions. At ambient temperature with CSI 'criss-cross' adducts are obtained in high yields. The reaction of CSI, for instance, with benzaldehyde azine gives diphenyltriazolotriazolone 69³².




30.2 Thiazolopyrimidine CSI reacts with penams such as methyl 6- β -phthalimidopenicillinate to give a thiazolo[3,2-c]pyrimidine 70⁶³ derivative by ring expansion.




30.3 Fused 1,2,4,6-Thiatriazine-1,1-dioxides: Reaction of 2-aminopyridine with CSI produces the N-chlorosulfonyl intermediate which cyclizes on the addition of a base to afford a fused 1,2,4,6-thiatriazine-1,1-dioxide system 71⁶⁴. Such systems have also been obtained by the reaction of CSI with 2-aminopyrazine and 2-aminothiazoline.




In case of 2-aminothiazoline it has been established that CSI attacked the ring nitrogen preferentially to give product 72⁵⁷.




30.4 **Thiatriazinobenzimidazoles** Addition of CSI to 2-aminobenzimidazole leads to 2H[1,2,4,6]-thiatriazino[2,3-a]benzimidazol-3(4H)-one-1,1-dioxide 73⁶⁵.



30.5 **Thiatriazinoquinazoline**: Reaction of CSI with 2-amino-4-arylquinazoline yields 6-aryl[1,2,4,6]-thiatriazino[2,3-a]quinazoline-3(2H)-one-1,1-dioxide 74⁶⁵.






30.6 **Azabullvalene**: It has been demonstrated by Paquette that the reaction of CSI with bullvalene emerges as a practical route of synthesis for methoxyazabullvalene 75⁶⁶.



31. Novel (unusual) heterocycles

Some bicyclic hydrocarbons undergo formal cycloaddition reactions with CSI at the strained, carbon-carbon single bonds on account of their high degree of π character⁶⁷. The products formed are often novel heterocycles and may be difficult to prepare by any other method. Some examples^{68,69} are illustrated below.



32. Conclusion

It is seen from the foregoing pages that CSI has truly been a reactive, broad based reagent for the synthesis of a variety of novel heterocycles. In addition, it affords an alternate and convenient method of synthesis for a number of known heterocyclic systems. It is hoped that CSI will further open up new vistas in heterocyclic synthesis.

REFERENCES

1. R. Graf, Angew. Chem., Int. Ed. Engl., 1968, 7, 172.
2. J.K. Rasmussen and A. Hassner, Chem. Rev., 1976, 76, 389.
3. W.A. Szabo, Aldrichimica Acta, 1977, 10, 23.
4. H. Ulrich, "Cycloaddition Reactions of Heterocumulenes", Academic Press, Inc., New York, N.Y., 1967, pp. 135-141.
5. K. Clauss, Liebigs Ann. Chem., 1969, 722, 110.
6. H. Bestian, Pure Appl. Chem., 1971, 27, 611.
7. N.S. Isaacs, Chem. Soc. Rev., 1976, 5, 181.
8. R. Graf, Liebigs Ann. Chem., 1963, 661, 111.
9. E.J. Moriconi, "Mechanisms of Reactions of Sulfur Compounds", Vol. 3, Intra-Science Research Foundation, Sanata Monica, Calif., 1968, p. 131.
10. E.J. Moriconi and W.C. Meyer, J. Org. Chem., 1971, 36, 2841.
11. R.B. Woodward and R. Hoffmann, Angew. Chem., Int. Ed. Engl., 1969, 8, 781.
12. K. Clauss, Tetrahedron Lett., 1974, 1271.
13. T.J. Barton and R.J. Rogido, Tetrahedron Lett., 1972, 3901.
14. T.J. Barton and R.J. Rogido, Chem. Commun., 1972, 878.
15. D.H. Ane, H. Iwahashi and D.F. Shellhamer, Tetrahedron Lett., 1973, 3719.
16. L.A. Paquette, T. Kakihana, J.F. Hansen and J.C. Philips, J. Am. Chem. Soc., 1971, 93, 152.
17. A. de S. Gomes and A.M. Figueiredo, Org. Prep. Proced. Int., 1973, 5, 13.
18. E.J. Moriconi and P.H. Mazzochi, J. Org. Chem., 1966, 31, 1372.
19. J.R. Malpass and N.J. Tweddle, Chem. Commun., 1972, 1255.
20. G.T. Furst, M.A. Wachsman, J. Pieroni, J.G. White and E.J. Moriconi, Tetrahedron, 1973, 29, 1675.
21. T. Sasaki, S. Eguchi and H. Yamada, J. Org. Chem., 1973, 38, 679.
22. T. Sasaki, S. Eguchi and Y. Hirako, Tetrahedron, 1976, 32, 437.
23. A.J.G. Baxter and K.H. Dickinson, Chem. Commun., 1979, 236.
24. K. Clauss, D. Grimm and G. Prossel, Liebigs Ann. Chem., 1974, 539.
25. P.H. Bentley and G. Brooks, Tetrahedron Lett., 1979, 21, 1889.
26. G. Gosteli, M. Lang and K. Prasat, J. Am. Chem. Soc., 1979, 101, 6296.
27. G. Gosteli, H. Pfaendler and R.B. Woodward, J. Am. Chem. Soc., 1980, 102, 2039.
28. P. Lombardi, G. Franceschi and R. Arcamone, Tetrahedron Lett., 1979, 39, 3777.
29. S. Oida and A. Yoshida, J. Antibiot., 1980, 33, 107.
30. K. Hirai, J. Synth. Org. Chem. Japan, 1980, 38, 2.

31. E.J. Moriconi and Y. Shimakawa, J. Org. Chem., 1972, 37, 196.
32. H. Suschitzky, E.R. Walrond and R. Hill, J. Chem. Soc., Perkin Trans. 1, 1977, 47.
33. T.W. Doyle and T.T. Conway, Tetrahedron Lett., 1969, 1889.
34. H. Hofmann, R. Wagner and J. Uhl, Chem. Ber., 1971, 104, 2134.
35. Hoechst, A.G., German Patent, 1,230,016 (8 Dec. 1966); CA, 67, 65095s.
36. D.N. Dhar and R. Ragunathan, Synthesis, 1982, 1095.
37. Hoechst, A.G., German Patent, 2,855,884 (10 Jul. 1980); CA, 94, 15413b.
38. K. Clauss, Liebigs Ann. Chem., 1974, 561.
39. H. Suschitzky and E.R. Walrond, J. Chem. Soc., Chem. Commun., 1973, 16, 570.
40. K. Hirai and H. Matsuda, Sankyo Kentyushi Nempo, 1972, 24, 108.
41. K. Clauss and N.J. Tweddle, Liebigs Ann. Chem., 1969, 722, 122.
42. J.R. Malpass and N.J. Tweddle, J. Chem. Soc., Chem. Commun., 1972, 1246.
43. D.N. Dhar, Indian J. Chem., 1976, 14B, 477.
44. J.K. Rasmussen and A. Hassner, J. Am. Chem. Soc., 1975, 97, 1451.
46. F. Hollywood, H. Suschitzky and R. Hull, Synthesis, 1982, 662.
47. A. Kamal, K.R. Rao and P.B. Sattur, Synth. Commun., 1980, 10, 799.
48. A. Kamal, Ph.D. Thesis, Aligarh Muslim University, Aligarh, 1982.
49. A. Kamal and P.B. Sattur, Synth. Commun., 1982, 12, 157.
50. A. Kamal, P.B. Sattur and G. Thyagarajan, Indian Patent Appl. 178/Del/82 (1982); European Patent Appl. 833011745 (1983).
51. D.N. Dhar and A.K. Bag, Indian J. Chem., 1982, 21B, 366.
52. T. Wagner-Tauregg, Synthesis, 1980, 10, 785.
53. Y. Grand, J.G. Atkinson and J. Rokach, J. Chem. Soc., Perkin Trans. 1, 1979, 1043.
54. K. Clauss, H.J. Friedrich and H. Jensen, Liebigs Ann. Chem., 1974, 561.
55. Hoechst, A.G., German Patent 2,327,804 (2 Jan. 1975) CA, 82, 171097.
56. A. Kamal and P.B. Sattur, Synthesis, 1981, 272.
57. S. Karady, Heterocycles, 1979, 12, 1199.
58. G.K. Denny and E.J. Cragoe, J. Org. Chem., 1980, 45, 1662.
59. D.J. Pasto and A.F.T. Chen, Tetrahedron Lett., 1973, 713.
60. E. Otto, W. Durkheimer and R. Muschweek, German Patent, 2,409,355.
61. M. Hedayatullah and J.F. Brault, Phosphorus and Sulfur, 1981, 11, 255.
62. L.A. Paquett and T. Kakihana, J. Am. Chem. Soc., 1968, 90, 3897.
63. M.M. Campbell, R.G. Harcus and K.H. Nelson, J. Chem. Soc., Perkin Trans. 1, 1978, 817.

64. S. Karady, Heterocycles, 1979, 12, 815.
65. A. Kamal and P.B. Sattur, Unpublished results from this laboratory, 1982.
66. L.A. Paquette, S. Kirschener and J.R. Malpass, J. Am. Chem. Soc., 1970, 92, 4330.
67. L.A. Paquette, G.R. Krow and T.J. Barton, in "Organic Photochemical Synthesis", Vol. 1, John Wiley & Sons, Inc., New York, N.Y., 1971, pp. 67-69.
68. L.A. Paquette, G.R. Allen, Jr., and M.J. Broadhurst, J. Am. Chem. Soc., 1971, 93, 4503.
69. W.E. Volz, L.A. Paquette, R.J. Rogido and T.J. Barton, Chem. Ind. (London), 1974, 771.

Received, 23rd May, 1986