

INDEX TO ADVERTISERS

Lederle Laboratories 799-800
 Lederle Praxis Biologicals 820A-D, 821

Mead Johnson Nutritionals 833
 Mead Johnson Pharmaceuticals 807-808

Pratt Pharmaceutical, a Division of Pfizer, Inc Cover 2- 798

Rhone-Poulenc Rorer 804A-B, 812, 834-836, 889

SmithKline-Beecham Laboratories 803
 Syntex Laboratories Cover 3- Cover 4

Wyeth-Ayerst Laboratories 815-816


Zeneca Pharmaceuticals 825-826

While every precaution is taken to ensure accuracy, we cannot guarantee against the possibility of an occasional change or omission in the preparation of this index.

Downloaded from www.archfammed.com at STANFORD Univ Med Center, on Nov 10, 2015

NAPROSYN®
 (NAPROXEN) 500 mg tablets

Brief Summary:
Contraindications: Patients who have had allergic reactions to NAPROSYN, ANAPROX or ANAPROX DS or in whom aspirin or other NSAIDs induce the syndrome of asthma, rhinitis, and nasal polyps. Because anaphylactic reactions usually occur in patients with a history of such reactions, question patients for asthma, nasal polyps, urticaria, and hypotension associated with NSAIDs before starting therapy. If such symptoms occur, discontinue the drug. **Warnings:** Serious GI toxicity such as bleeding, ulceration, and perforation can occur at any time, with or without warning symptoms, in patients treated chronically with NSAIDs. Remain alert for ulceration and bleeding in such patients even in the absence of previous GI tract symptoms. In clinical trials, symptomatic upper GI ulcers, gross bleeding or perforation appear to occur in approximately 1% of patients treated for 3-6 months, and in about 2-4% of patients treated for one year. Inform patients about the signs and/or symptoms of serious GI toxicity and what steps to take if they occur. Studies have not identified any subset of patients not at risk of developing peptic ulceration and bleeding. Except for a prior history of serious GI events and other risk factors known to be associated with peptic ulcer disease, such as alcoholism, smoking, etc., no risk factors (e.g., age, sex) have been associated with increased risk. Elderly or debilitated patients seem to tolerate ulceration or bleeding less well than others and most spontaneous reports of fatal GI events are in this population. In considering the use of relatively large doses (within the recommended dosage range), sufficient benefit should be anticipated to offset the potential increased risk of GI toxicity. **Precautions:** DO NOT GIVE NAPROSYN® (NAPROXEN) CONCOMITANTLY WITH ANAPROX® (NAPROXEN SODIUM) OR ANAPROX® DS (NAPROXEN SODIUM) SINCE THEY BOTH CIRCULATE IN PLASMA AS THE NAPROXEN ANION. Acute interstitial nephritis with hematuria, proteinuria, and nephrotic syndrome has been reported. Patients with impaired renal function, heart failure, liver dysfunction, patients taking diuretics, and the elderly are at greater risk of overt renal decompensation. If this occurs, discontinue the drug. Use with caution and monitor serum creatinine and/or creatinine clearance in patients with significantly impaired renal function. Use caution in patients with baseline creatinine clearance less than 20 mL/minute. Use the lowest effective dose in the elderly or in patients with chronic alcoholic liver disease or cirrhosis. With NSAIDs, borderline elevations of liver tests may occur in up to 15% of patients. They may progress, remain unchanged, or be transient with continued therapy. Elevations of SGPT or SGOT occurred in controlled clinical trials in less than 1% of patients. Severe hepatic reactions, including jaundice and fatal hepatitis, have been reported rarely. If liver disease develops or if systemic manifestations occur (e.g., eosinophilia or rash), discontinue therapy. If steroid dosage is reduced or eliminated during therapy, do so slowly and observe patients closely for adverse effects, including adrenal insufficiency and exacerbation of arthritis symptoms. Determine hemoglobin values periodically for patients with initial values of 10 grams or less who receive long-term therapy. Peripheral edema has been reported. Therefore, use with caution in patients with fluid retention, hypertension or heart failure. The drug's antipyretic and anti-inflammatory activities may reduce fever and inflammation, diminishing their diagnostic value. Conduct ophthalmic studies if any change or disturbance in vision occurs. For patients with restricted sodium intake, note that the suspension contains 8 mg/mL of sodium. **Information for Patients:** Side effects of NSAIDs can cause discomfort and, rarely, there are more serious side effects, such as GI bleeding, which may result in hospitalization and rarely, death. Physicians may wish to discuss with patients the potential risks and likely benefits of NSAID treatment, particularly when they are used for less serious conditions where treatment without NSAIDs may be an acceptable alternative. Patients should use caution for activities requiring alertness if they experience drowsiness, dizziness, vertigo or depression during therapy. **Laboratory Tests:** Because serious GI tract ulceration and bleeding can occur without warning symptoms, follow chronically treated patients for signs and symptoms of these and inform them of the importance of this follow-up. **Drug Interactions:** Use caution when giving concomitantly with coumarin-type anticoagulants; a hydantoin, sulfonamide or sulfonurea; furosemide; lithium; beta-blockers; probenecid; or methotrexate. **Drug/Laboratory Test Interactions:** The drug may decrease platelet aggregation and prolong bleeding time or increase urinary values for 17-ketogenic steroids. Temporarily stop therapy for 72 hours before doing adrenal function tests. The drug may interfere with urinary assays of SHIAA. **Carcinogenesis:** A 2-year rat study showed no evidence of carcinogenicity. **Pregnancy:** Category B. Do not use during pregnancy unless clearly needed. Avoid use during late pregnancy. **Nursing Mothers:** Avoid use in nursing mothers. **Pediatric Use:** Single doses of 2.5-5 mg/kg, with total daily dose not exceeding 15 mg/kg/day, are safe in children over 2 years of age. **Adverse Reactions:** In a study, GI reactions were more frequent and severe in rheumatoid arthritis patients on 1,500 mg/day than in those on 750 mg/day. In studies in children with juvenile arthritis, rash and prolonged bleeding times were more frequent. GI and CNS reactions about the same, and other reactions less frequent than in adults. **Incidence Greater Than 1%:** Probable Causal Relationship: GI: The most frequent complaints related to the GI tract: constipation, heartburn, abdominal pain, nausea, dyspepsia, diarrhea, stomatitis. CNS: headache, dizziness, drowsiness, light-headedness, vertigo. Dermatologic: itching (pruritus), skin eruptions, ecchymoses, sweating, purpura. Special Senses: tinnitus, hearing disturbances, visual disturbances. Cardiovascular: edema, dyspnea, palpitations. General: thirst. **Incidence Less Than 1%:** Probable Causal Relationship: GI: abnormal liver function tests, colitis, GI bleeding and/or perforation, hematemesis, jaundice, melena, peptic ulceration with bleeding and/or perforation, vomiting. Renal: glomerular nephritis, hematuria, hyperkalemia, interstitial nephritis, nephrotic syndrome, renal disease, renal failure, renal papillary necrosis. Hematologic: agranulocytosis, eosinophilia, granulocytopenia, leukopenia, thrombocytopenia. CNS: depression, dream abnormalities, inability to concentrate, insomnia, malaise, myalgia and muscle weakness. Dermatologic: alopecia, photosensitive dermatitis, skin rashes. Special Senses: hearing impairment. Cardiovascular: congestive heart failure. Respiratory: eosinophilic pneumonitis. General: anaphylactoid reactions, menstrual disorders, pyrexia (chills and fever). Causal Relationship Unknown: Hematologic: aplastic anemia, hemolytic anemia. CNS: aseptic meningitis, cognitive dysfunction. Dermatologic: epidermal necrolysis, erythema multiforme, photosensitivity reactions resembling porphyria cutanea tarda and epidermolysis bullosa, Stevens-Johnson syndrome, urticaria. GI: non-peptic GI ulceration, ulcerative stomatitis. Cardiovascular: vasculitis. General: angioneurotic edema, hyperglycemia, hypoglycemia. **Overdosage:** May have drowsiness, heartburn, indigestion, nausea, vomiting. A few patients have had seizures. Empty stomach and use usual supportive measures. In animals 0.5 g/kg of activated charcoal reduced plasma levels of naproxen. **Caution:** Federal law prohibits dispensing without prescription. See package insert for full Prescribing information.

Incidence of reported reaction 3%-9%.
 Where unmarked, incidence less than 3%.

 U.S. patent nos. 3,904,682, 3,998,966 and others.
 © 1991 Syntex Puerto Rico, Inc. Rev. 39 September 1990