


CONJUGATE ADDITION OF ARYLMETAL REAGENTS TO AN α,β -UNSATURATED KETONE. SHORT SYNTHESSES OF (-)-AGATHARESINOL DIMETHYL ETHER AND (-)-SUGIRESINOL DIMETHYL ETHER

Keizo Matsuo*, Yuji Ono, Atsutaka Seki, Hiroshi Kuwajima, and Keiji Nishiwaki

Faculty of Pharmaceutical sciences, Kinki University
3-4-1 Kowakae, Higashiosaka, Osaka 577-8502, Japan

Abstract – (-)-Agatharesinol dimethyl ether and (-)-sugiresinol dimethyl ether, derivatives of (-)-agatharesinol and (-)-sugiresinol isolated from *Agathis australis* and *Cryptomeria Japonica*, respectively were synthesized starting from a protected D-glyceraldehyde in five steps through 1,4-conjugate addition of arylmetal reagents to an α,β -unsaturated ketone.

(-)-Agatharesinol (**1a**) and (-)-sugiresinol (**2a**) are norlignans isolated from heartwood of *Agathis australis* by Enzell and Thomas¹ in 1965 and *Cryptomeria Japonica* by Funaoka *et al.*² in 1963, respectively, and their chemical structures including absolute stereochemistry were determined by Enzell *et al.*³ in 1966. The norlignans (**1a** and **2a**) have been reported to show biological activities such as antifungal activity,⁴ inhibitory effect on cyclic AMP phosphodiesterase,⁵ inhibitory activity against *C. shiitake* hyphae growth and fruiting body formation,⁶ and vinyl polymerization-inhibitory activity.⁷ Syntheses of agatharesinol dimethyl ether (**1b**) and sugiresinol dimethyl ether (**2b**) have been reported by Horii *et al.*⁸ and Beracierta and Whiting⁸ as racemic form and by Muraoka *et al.*⁹ and Dujardin *et al.*¹⁰ as optically active form, respectively. Recently, we also reported the synthesis of (-)-**2b** utilizing (-)-quinic acid as the chiral source.¹¹




Scheme 1


As a part of our synthetic studies on biologically active natural products, we would like to report here the

short syntheses (five steps) of (-)-**1b** and (-)-**2b** starting from a protected D-glyceraldehyde¹² through 1,4-conjugate addition of arylmetal reagents to the α,β -unsaturated chiral ketone (**3**).¹³

Leonard and Ryan¹⁴ have reported that isopropenylcopper reacted with α,β -unsaturated ketone (**4**) (*E:Z*=6:1) which has an oxygen function at γ -position to give a mixture of *anti*- and *syn*-adducts (**5:6**=7:1) in 79% yield, on the other hand, in the reaction of isopropenyllithium with **4** (*E:Z*=7:2), a mixture of *anti*- and *syn*-adducts (**5:6**=1:36) in 60% yield (Scheme 2). These results indicated that the copper reagent added to **4** from the less hindered side to afford the *anti*-isomer (**5**) predominantly (kinetic control), but the lithium reagent chelated first to the oxygen atom at γ -position and then the alkyl group attacked the β -position of the α,β -unsaturated ketone from the same face to give the *syn*-isomer (**6**) predominantly (chelation control) (Scheme 3).




Scheme 2



Scheme 3

According to the results of above Leonard and Ryan's experiment, if the 4-methoxyphenyl-metal reagents react with **3** under chelation control, the *syn*-adduct (**8**) would be obtained predominantly (Scheme 4). Two stereogenic centers of **8** suit for the stereochemistry of 2 and 3 positions of (-)-**1b** and 4 and 5 positions of (-)-**2b**.



Scheme 4

In our synthetic work on (-)-**1b** and (-)-**2b**, the α,β -unsaturated chiral ketone (**3**) was synthesized first to investigate the 1,4-conjugate addition of arylmetal reagents. Thus, the protected D-glyceraldehyde,¹²

which was derived from D-mannitol, was treated with 4-methoxybenzoylmethylenetriphenylphosphorane¹⁵ to give a mixture (3:1) of (*E*)- and (*Z*)- α,β -unsaturated ketones (**3**) in quantitative yield, which possessed chiral oxygen functions at γ -position. Then the conjugate addition reaction of the various 4-methoxyphenyl-metal reagents to **3** was investigated. The results are summarized in Table 1.

Table 1. Conjugate Addition of Arylmetal Reagents to **3**

Entry	Reagents ^{a)}	Temp. (°C)	Yield (%)	7 : 8
1	ArMgBr (3.3 eq.) CuBr-Me ₂ S (1.6 eq.), THF	-15	46	3 : 1 ^{b)}
2	Ar-Li (1.2 eq.), THF	-78	45	4 : 1 ^{c)}
3	Ar-Li (1.8 eq.), Et ₂ O	-78	38	nd ^{d)}
4	Ar-I (1.2 eq.), (Ph ₃ P) ₄ Pd, Et ₃ N	reflux	28	3 : 1 ^{c)}


a) Ar=4-Methoxyphenyl.

b) Each compound was isolated by preparative SiO₂ TLC.

c) The ratio was determined with ¹H-NMR.

d) Not determined.

When a mixture of 4-methoxyphenylmagnesium bromide (3.3 eq.) and CuBr-Me₂S complex (1.6 eq.)¹⁶ was treated with **3** (a mixture of *E*:*Z*=3:1) in THF at -15 °C, a mixture (3:1) of **7** and **8** was obtained in 46% yield (Entry 1). The reaction of 4-methoxyphenyllithium (prepared from 4-iodoanisole and *n*-butyllithium)¹⁷ with **3** in THF at -78 °C resulted in formation of a mixture (4:1) of **7** and **8** in 45 % yield (Entry 2). In this reaction, change of the solvent from THF to Et₂O caused to lower the yield (Entry 3). When a mixture of 4-iodoanisole and (Ph₃P)₄Pd¹⁸ was treated with **3** in the presence of Et₃N at refluxing temperature, a mixture (3:1) of **7** and **8** was also produced in 28% yield (Entry 4). These results suggested that the reaction of the arylmetal reagents used above took place to give the kinetically controlled product (**7**) predominantly, even when the lithium reagent was employed (Entry 2). These evidences are probably attributable to the steric hindrance of both the arylmetal reagents and the



Scheme 5

cyclohexylidene protective group (Scheme 5). Although the starting compound (**3**) was almost

internal standard. High-resolution MS spectra (HRMS) were measured with a JEOL JMS-HX100 instrument at 70 eV.

(4S)-4,5-Cyclohexylidenedioxy-1-(4-methoxyphenyl)pent-2-en-1-one (3)

4-Methoxybenzoylmethylenetriphenylphosphorane¹⁵ (6.20 g, 15.1 mmol) was added to a solution of (2R)-2,3-cyclohexylidenedioxypropanal¹² (2.57 g, 15.1 mmol) in MeCN (100 mL), and the whole was stirred at room rt for 1 h and refluxed for 30 min. The reaction mixture was concentrated under reduced pressure and ether (100 mL) was added to the residue. The mixture was stirred vigorously to form precipitates, which were removed by filtration. The filtrate was concentrated under reduced pressure and the residue was purified by silica gel column chromatography (*n*-hexane:AcOEt=4:1) to give a mixture of (*E*)- and (*Z*)-**3** (4.56 g, quantitative yield). ¹H-NMR spectrum of the mixture showed *E*:*Z*=3:1. Repeated silica gel column chromatography (*n*-hexane:AcOEt=6:1) of the mixture afforded (*E*)- and (*Z*)-**3** as oils. (*E*)-**3**: IR (neat): 1665 cm⁻¹. ¹H-NMR (CDCl₃) δ: 1.38-1.74 (10H, m, cyclohexylidene), 3.73 (1H, dd, *J*= 8.5, 7.0 Hz, OCH₂), 3.88 (3H, s, OCH₃), 4.23 (1H, dd, *J*= 8.0, 7.0 Hz, OCH₂), 4.79 (1H, m, OCH), 6.94 (1H, dd, *J*= 15.5, 5.5 Hz, OCHCH=CH), 6.96 (2H, dt, *J*=9.0, 2.5 Hz, ArH), 7.20 (1H, dd, *J*=15.5, 1.5 Hz, CH=CHCO), 7.97 (2H, dt, *J*=9.0, 2.5 Hz, ArH). HRMS (*m/z*): Calcd for C₁₈H₂₂O₄ (M⁺): 302.1518. Found: 302.1523. (*Z*)-**3**: IR (neat): 1665 cm⁻¹. ¹H-NMR (CDCl₃) δ: 1.38-1.74 (10H, m, cyclohexylidene), 3.69 (1H, dd, *J*=8.5, 7.0 Hz, OCH₂), 3.88 (3H, s, OCH₃), 4.50 (1H, dd, *J*=8.0, 7.0 Hz, OCH₂), 5.35 (1H, m, OCH), 6.43 (1H, dd, *J*=12.0, 7.0 Hz, OCHCH=CH), 6.92 (1H, dd, *J*=12.0, 2.0 Hz, CH=CHCO), 6.96 (1H, dt, *J*=9.0, 2.5 Hz, ArH), 7.97 (1H, dt, *J*=9.0, 2.5 Hz, ArH).

(4S)-4,5-Cyclohexylidenedioxy-1,3-bis(4-methoxyphenyl)pentan-1-ones (7 and 8)

Entry 1. A THF solution of 4-methoxyphenylmagnesium bromide was prepared from magnesium turnings (80 mg, 3.3 mmol), 4-bromoanisole (0.42 mL, 3.4 mmol) and THF (1.5 mL) by usual method. A cold solution of CuBr-Me₂S complex (0.322 g, 1.6 mmol) and Me₂S (1.8 mL) in THF (3.5 mL) was added through cannula to the above Grignard reagent at -40 °C. A solution of **3** (*E*:*Z*=3:1) (0.294 g, 1.0 mmol) in THF (2.0 mL) was added dropwise to the above reaction mixture at -15 °C and the whole was stirred at that temperature for 15 min. After the reaction mixture was warmed to rt, saturated aqueous solution of NH₄Cl was added and then the solution was concentrated to remove THF under reduced pressure. Water (10 mL) was added to the concentrated solution and the mixture was extracted with AcOEt. The extract was washed with 10% ammonia solution, H₂O and brine, successively. After drying over anhydrous Na₂SO₄, the organic layer was concentrated under reduced pressure and the residue was purified by silica gel column chromatography (benzene:AcOEt=30:1) to give a mixture of **7** and **8** (0.399 g, 46%) as a pale yellow oil. The compounds (**7**) and (**8**) were isolated from the mixture by repeated silica gel preparative TLC (*n*-hexane: AcOEt=10:1) and the ratio of **7** and **8** was 3:1. **7**: oil. IR (neat): 1675 cm⁻¹. ¹H-NMR (CDCl₃) δ: 1.32–1.68 (10H, m, cyclohexylidene), 3.28 (1H, dd, *J*= 15.5, 9.5 Hz, ArCHCH₂CO), 3.39 (1H, td, *J*=9.5, 3.0 Hz, OCHCHAr), 3.58 (1H, dd, *J*=9.0, 6.0 Hz, OCH₂CHO), 3.62 (1H, dd, *J*=15.5, 3.0 Hz, ArCHCH₂CO), 3.71-3.78 (1H, m, OCH₂CHO), 3.75 (3H, s, OCH₃), 3.85 (3H, s, OCH₃), 4.26 (1H, td, *J*=9.5, 6.0 Hz, OCH₂CHO), 6.79 (2H, dt, *J*=9.0, 2.5 Hz, ArH), 6.89 (2H, dt, *J*=9.0, 2.5 Hz, ArH), 7.16 (2H, dt, *J*=9.0, 2.5 Hz, ArH), 7.90 (2H, dt, *J*=9.0, 2.5 Hz, ArH). **8**: oil. IR (neat): 1675 cm⁻¹. ¹H-NMR (CDCl₃) δ: 1.32-1.68 (10H, m, cyclohexylidene), 3.34 (1H, dd, *J*=16.5, 8.0 Hz, ArCHCH₂CO), 3.47 (1H,

dd, $J=16.5, 8.0$ Hz, ArCHCH₂CO), 3.52 (1H, t, $J=8.0$ Hz, ArCHCH₂), 3.56-3.63 (1H, m, OCH₂CH), 3.75 (3H, s, OCH₃), 3.85 (3H, s, OCH₃), 3.92 (1H, dd, $J=8.0, 6.0$ Hz, OCH₂CH), 4.36 (1H, ddd, $J=8.0, 6.0, 5.0$ Hz, OCH₂CHO), 6.79 (2H, dt, $J=9.0, 2.5$ Hz, ArH), 6.89 (2H, dt, $J=9.0, 2.5$ Hz, ArH), 7.16 (2H, dt, $J=9.0, 2.5$ Hz, ArH), 7.90 (2H, dt, $J=9.0, 2.5$ Hz, ArH). HRMS (m/z): Calcd for C₂₅H₃₀O₅ (M⁺): 410.2093. Found: 410.2101.

Entry 2. *n*-BuLi (1.58 M *n*-hexane solution, 1.27 mL, 2.01 mmol) was added to a stirred solution of 4-iodoanisole (0.46 g, 1.98 mmol) in THF (2.5 mL) at -78 °C. A solution of **3** (*E*:*Z*=3:1)(0.50 g, 1.65 mmol) in THF (1.0 mL) was added to the above reaction mixture at -78 °C and the whole was stirred at that temperature for 20 min. After addition of H₂O, the mixture was extracted with AcOEt (x 3) and the combined extract was washed with H₂O and brine, successively. The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to give a residue, which was purified by silica gel column chromatography (benzene:AcOEt=30:1) as above. A mixture of **7** and **8** (0.39 g, 45%) was obtained as a pale yellow oil and the ratio of each determined as 4:1 by ¹H-NMR spectral data.

Entry 3. *n*-BuLi (1.58 M *n*-hexane solution, 0.78 mL, 1.23 mmol) was added to a stirred solution of 4-iodoanisole (0.28 g, 0.79 mmol) in Et₂O (5.0 mL) at -78 °C. A solution of **3** (*E*:*Z*=3:1)(0.20 g, 0.66 mmol) in Et₂O (1.5 mL) was added to the above reaction mixture at -78 °C and the whole was stirred at that temperature for 25 min. After addition of H₂O, the mixture was extracted with AcOEt and the combined extract was washed with H₂O and brine, successively. The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to give a residue, which was purified by silica gel column chromatography (benzene:AcOEt=30:1) as above. A mixture of **7** and **8** (0.10 g, 38%) was obtained as a pale yellow oil.

Entry 4. A mixture of **3** (*E*:*Z*=3:1)(0.20 g, 0.66 mmol), 4-iodoanisole (0.19 g, 0.80 mmol), (Ph₃P)₄Pd (7 mg, 6×10^{-3} mmol) and Et₃N (0.30 mL, 2.15 mmol) was refluxed for 7 h. After addition of 10% HCl and H₂O, the mixture was extracted with AcOEt (x 2) and the combined extract was washed with H₂O and brine, respectively. The organic layer was dried over anhydrous Na₂SO₄ and then concentrated under reduced pressure to give a residue, which was purified by silica gel column chromatography (*n*-hexane:AcOEt=19:1) as above. A mixture of **7** and **8** (76 mg, 28%) was obtained as pale yellow oil and the ratio of each determined as 3:1 by ¹H-NMR spectral data.

(2*S*)-1,2-Cyclohexylidenedioxy-3,5-bis(4-methoxyphenyl)pentan-5-ol (**9**)

NaBH₄ (93.5 mg, 2.5 mmol) was added to a solution of **7** and **8** (3:1 mixture, 0.496 g, 1.2 mmol) in EtOH (6.0 mL) under ice-cooling and the whole was stirred at rt for 7 h. After addition of H₂O and sodium chloride, the mixture was extracted with CHCl₃ (x 3) and the combined organic layer was washed with brine. The organic layer was dried over anhydrous Na₂SO₄ and then concentrated under reduced pressure to give **9** (0.491 g, 99%) as a pale yellow oil. Since this compound was relatively unstable, it was submitted for the next reaction immediately after taking the IR spectrum. IR (neat): 3450 cm⁻¹.

(2*S*,3*R*)-(4*E*)- and (2*S*,3*S*)-(4*E*)-1,2-Cyclohexylidenedioxy-3,5-bis(4-methoxyphenyl)pent-4-enes (**10** and **11**)

Methanesulfonyl chloride (0.375 mL, 4.8 mmol) was added to a solution of **9** (0.483 g, 1.2 mmol) in dry pyridine (12.0 mL) and the reaction mixture was stirred at rt for 14 h and then refluxed for 5 h. The reaction was quenched by addition of cold water and the mixture was extracted with CHCl₃ (x 5). The

combined extract was washed with 5% HCl, saturated NaHCO₃ aqueous solution and brine, successively and dried over anhydrous Na₂SO₄. The organic layer was concentrated under reduced pressure to give a brown oil which was purified by silica gel column chromatography (*n*-hexane:AcOEt=10:1) to furnish a mixture of **10** and **11** (0.322 g, 70%) as a pale yellow oil. The oil was separated into each **10** and **11** (3:1) by careful silica gel column chromatography (benzene). **10**: oil. IR (neat): 965 cm⁻¹. ¹H-NMR (CDCl₃) δ: 1.30-1.75 (10H, m, cyclohexylidene), 3.45 (1H, t, *J*=7.0 Hz, ArCH), 3.60 (1H, dd, *J*=8.0, 7.0 Hz, OCH₂), 3.79 (3H, s, OCH₃), 3.80 (3H, s, OCH₃), 3.82 (1H, dd, *J*=8.0, 6.0 Hz, OCH₂), 4.40 (1H, ddd, *J*=8.0, 7.0, 6.0 Hz, OCH), 6.28 (1H, d, *J*=16.0 Hz, CH=CHAr), 6.40 (1H, dd, *J*=16.0, 6.5 Hz, CHCH=CH), 6.81 (2H, dt, *J*=9.0, 2.5 Hz, ArH), 6.86 (2H, dt, *J*=9.0, 2.5 Hz, ArH), 7.18 (2H, dt, *J*=9.0, 2.5 Hz, ArH), 7.28 (2H, dt, *J*=9.0, 2.5 Hz, ArH). [α]_D²²: -10.8° (*c*=1.00, CHCl₃). **11**: oil. IR (neat): 960 cm⁻¹. ¹H-NMR (CDCl₃) δ: 1.29-1.68 (10H, m, cyclohexylidene), 3.55 (1H, t, *J*=8.0 Hz, ArCH), 3.79 (6H, s, 2 x OCH₃), 3.80 (1H, dd, *J*=8.0, 7.0 Hz, OCH₂), 4.03 (1H, dd, *J*=8.0, 6.0 Hz, OCH₂), 4.41 (1H, ddd, *J*=8.0, 7.0, 6.0 Hz, OCH), 6.16 (1H, dd, *J*=15.5, 8.0 Hz, CHCH=CH), 6.41 (1H, d, *J*=15.5 Hz, CH=CHAr), 6.82 (2H, dt, *J*=9.0, 2.5 Hz, ArH), 6.86 (2H, dt, *J*=9.0, 2.5 Hz, ArH), 7.23 (2H, dt, *J*=9.0, 2.5 Hz, ArH), 7.27 (2H, dt, *J*=9.0, 2.5 Hz, ArH). HRMS (*m/z*): Calcd for C₂₅H₃₀O₄ (M⁺): 394.2144. Found: 394.2150. [α]_D²²: -12.0° (*c*=1.06, CHCl₃).

(-)-Agatharesinol dimethyl ether (**1b**)

Concentrated HCl (0.52 mL) was added to a solution of **11** (20 mg, 0.051 mmol) in MeOH (3.5 mL) under ice-cooling and the whole was stirred at rt for 2 h. After addition of saturated NaHCO₃ aqueous solution, the mixture was extracted with CHCl₃ (x 5) and the combined extract was washed with brine. The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to give a residue, which was purified by silica gel PTLC (*n*-hexane:AcOEt=1:1) to give (-)-**1b** (16 mg, 98%) as a colorless oil. IR (neat): 3400, 1610, 1510, 1250 cm⁻¹. ¹H-NMR (CDCl₃) δ: 1.95 (2H, br s, 2 x OH), 3.52 (1H, t, *J*=9.0 Hz, OCHCHAr), 3.62 (1H, dd, *J*=11.5, 6.5 Hz, CH₂OH), 3.79 (3H, s, OCH₃), 3.80 (3H, s, OCH₃), 3.82 (1H, dd, *J*=11.5, 3.0 Hz, CH₂OH), 3.99 (1H, ddd, *J*=9.0, 6.5, 3.0 Hz, CHOH), 6.16 (1H, dd, *J*=16.0, 9.0 Hz, CH=CHAr), 6.42 (1H, d, *J*=16.0 Hz, CH=CHAr), 6.82 (2H, dt, *J*=9.0, 2.5 Hz, ArH), 6.90 (2H, dt, *J*=9.0, 2.5 Hz, ArH), 7.24 (2H, dt, *J*=9.0, 2.5 Hz, ArH), 7.26 (2H, dt, *J*=9.0, 2.5 Hz, ArH). [α]_D²⁰: -24.8° (*c*=1.24, CHCl₃). HRMS (*m/z*): Calcd for C₁₉H₂₂O₄ (M⁺): 314.1518. Found: 314.1513.

(-)-Sugiresinol dimethyl ether (**2b**)

Concentrated HCl (0.52 mL) was added to a solution of **11** (20 mg, 0.05 mmol) in MeOH (3.5 mL) and the whole was refluxed for 20 h. After cooling and neutralization by addition of saturated NaHCO₃ aqueous solution, the mixture was extracted with CHCl₃ (x 5). The combined extract was washed with brine and then dried over anhydrous Na₂SO₄. Removal of the solvent under reduced pressure gave a solid, which was purified by silica gel PTLC (*n*-hexane:AcOEt=1:1) to give (-)-**1b** (14 mg, 89%) as colorless crystalline powder. Sample for spectra was obtained by recrystallization from *n*-hexane. mp 103.5-104.5 °C (lit.³ 104-105 °C). IR (Nujol): 3400, 1610, 1585, 1510 cm⁻¹. ¹H-NMR (CDCl₃) δ: 1.62 (1H, br s, OH), 1.91 (1H, ddd, *J*=14.0, 12.5, 11.0 Hz, ArCHCH₂CHAr), 2.07 (1H, ddd, *J*=14.0, 4.0, 2.0 Hz, ArCHCH₂CHAr), 2.78 (1H, ddd, *J*=12.5, 10.0, 4.0 Hz, CH₂CHArCH), 3.48 (1H, dd, *J*=11.0, 10.0 Hz, OCH₂CH), 3.79 (3H, s, OCH₃), 3.80 (3H, s, OCH₃), 3.86 (1H, dt, *J*=10.0, 5.0 Hz, CHOH), 4.29 (1H, dd, *J*=11.0, 5.0 Hz, OCH₂CH), 4.47 (1H, dd, *J*=11.0, 2.0 Hz, OCHArCH₂), 6.86 (2H, dt, *J*=9.0, 2.5 Hz, ArH),

6.89 (2H, dt, $J=9.0$, 2.5 Hz, ArH), 7.22 (2H, dt, $J=9.0$, 2.5 Hz, ArH), 7.30 (2H, dt, $J=9.0$, 2.5 Hz, ArH). HRMS (m/z): Calcd for $C_{19}H_{22}O_4$: 314.1518. Found: 314.1499. $[\alpha]_D^{21}$: -4.0° ($c=1.01$, $CHCl_3$)(lit., $^{3,19}[\alpha]_D$: -4.0° ($CHCl_3$)).

REFERENCES

1. C. R. Enzell and B. R. Thomas, *Acta Chem. Scand.*, 1965, **19**, 913; R. C. Cambie, J. M. Coddington, M. J. Stone, N. Tanaka, Y.-H. Li, and S. Arigayo, *Phytochemistry*, 1989, **28**, 1675.
2. K. Funaoka, Y. Kuroda, Y. Kai, and T. Kondo, *Nippon Mokuzai Gakkaishi*, 1963, **9**, 139 [*Chem. Abstr.*, 1964, **60**, 1626].
3. C. R. Enzell and B. R. Thomas, *Tetrahedron Lett.*, **1966**, 2395; C. R. Enzell, Y. Hirose, and B. R. Thomas, *Tetrahedron Lett.*, **1967**, 793.
4. K. Ootsuka, K. Sakurada, and H. Nakamura, *Jpn. Kokai Tokkyo Koho*, JP 07 25719, **1995** [*Chem. Abstr.*, 1995, **122**, 233344t].
5. T. Nikaido, T. Ohmoto, H. Noguchi, T. Kinoshita, H. Saitoh, and U. Sankawa, *Planta Med.*, 1981, **43**, 18; U. Sankawa, *Saengyak Hakhoe Chi*, 1980, **11**, 125 [*Chem. Abstr.*, 1981, **95**, 125747s].
6. T. Sato and T. Sato, *Gyomu Hokoku-Akita-ken Ringyo Senta*, **1979**, 185 [*Chem. Abstr.*, 1981, **95**, 93999g].
7. H. Funakoshi, K. Nobashi, and T. Yokota, *Mokuzai Gakkaishi*, 1978, **24**, 141 [*Chem. Abstr.*, 1978, **88**, 170579e].
8. Z. Horii, C. Iwata, T. Tanaka, and T. Momose, *Heterocycles*, 1977, **6**, 697; A. P. Beracierta and D. A. Whiting, *J. Chem. Soc., Perkin Trans. 1*, **1978**, 1257.
9. O. Muraoka, N. Fujiwara, G. Tanabe, and T. Momose, *Tetrahedron: Asymmetry*, 1991, **2**, 357; O. Muraoka, B. Zheng, N. Fujiwara, and G. Tanabe, *J. Chem. Soc., Perkin Trans. 1*, **1996**, 405.
10. G. Dujardin, M. Maudet, and E. Brown, *Tetrahedron Lett.*, 1997, **38**, 1555; E. Brown, G. Dujardin, and M. Maudet, *Tetrahedron*, 1997, **53**, 9679.
11. K. Matsuo, W. Sugimura, Y. Shimizu, K. Nishiwaki, and H. Kuwajima, *Heterocycles*, 2000, **53**, 1505.
12. H. Ronneberg, G. Borch, R. Buchecker, N. Arpin, and S. L. Jensen, *Phytochemistry*, 1982, **21**, 2087; T. Sugiyama, H. Sugawara, M. Watanabe, and K. Yamashita, *Agric. Biol. Chem.*, 1984, **48**, 1841; S. Takano, A. Kurotaki, and K. Ogasawara, *Synthesis*, **1986**, 403.
13. K. Matsuo, Y. Ono, A. Seki, H. Kuwajima, and K. Nishiwaki, *Bull. Pharm. Res. Tech. Inst.*, **1999**, 105.
14. J. Leonard and G. Ryan, *Tetrahedron Lett.*, 1987, **28**, 2525.
15. F. Ramirez and S. Dershowitz, *J. Org. Chem.*, 1957, **22**, 41.
16. E. Nicolas, K. C. Russell, and V. J. Hruby, *J. Org. Chem.*, 1993, **58**, 766.
17. T. Iwasaki, K. Kondo, T. Nishitani, T. Kuroda, K. Hirakoso, A. Ohtani, and K. Takashima, *Chem. Pharm. Bull.*, 1995, **43**, 1701.
18. G. E. Stokker, *Tetrahedron Lett.*, 1987, **28**, 3179.
19. P. H. Smith and D. A. Whiting, *Phytochemistry*, 1976, **15**, 1285.