

ENANTIOCONTROL IN TANDEM ALLYLIC SULFONIUM YLIDE GENERATION AND [2,3] SIGMATROPIC REARRANGEMENT CATALYZED BY CHIRAL DIRHODIUM(II) COMPLEXES[†]

Shinji Kitagaki, Yoshimasa Yanamoto, Hisae Okubo, Makoto Nakajima, and Shunichi Hashimoto*


Graduate School of Pharmaceutical Sciences, Hokkaido University, Sapporo 060-0812, Japan

Abstract – Tandem sulfonium ylide formation and [2,3] sigmatropic rearrangement sequence from allylic sulfides and diazoacetates has been effected by using dirhodium(II) tetrakis[3(*S*)-phthalimido-2-piperidinonate], Rh₂(*S*-PTPI)₄, as a chiral catalyst, displaying enantioselectivities of up to 58%. It has been demonstrated that the [2,3] sigmatropic rearrangement proceeds through chiral, nonracemic free sulfonium ylides dissociated from the dirhodium(II) complex.

Allylic sulfonium ylides readily undergo a [2,3] sigmatropic rearrangement which features a carbon-carbon bond formation with a high degree of stereocontrol.¹ Considering that catalytic generation and [2,3] sigmatropic rearrangement of sulfonium ylides from allylic sulfides and a variety of metal carbenes have recently enjoyed widespread attention as an alternative to the process *via* deprotonation or desilylation of the corresponding sulfonium salts,² the development of an enantioselective version of this sequence mediated by chiral catalysts should be a significant addition to the field of asymmetric synthesis.³

Uemura and co-workers were the first to demonstrate asymmetric induction (up to 20% ee) in the reaction of (*E*)-cinnamyl phenyl sulfide (**1a**) and ethyl diazoacetate (**2a**) employing bis(oxazoline)-Cu(I) complex (**3a**) or Rh₂(*5S*-MEPY)₄ as a catalyst.⁴ Recently, the bis(oxazoline)-Cu(I)-catalyzed process has been systematically investigated by McMillen and co-workers, in which a significant enhancement of up to 52% ee can be attained by the combinational use of the sterically demanding allyl 2,6-dimethylphenyl sulfide and bis(oxazoline)-Cu(I) complex (**3b**).⁵ Katsuki and co-workers have reported that chiral salen-Co(III)-complex (**4**) is an efficient catalyst for the reaction of **1a** with *tert*-butyl diazoacetate (**2b**) to display the highest degree of enantioselectivity (64% ee) known for this type of reactions.⁶ They have also demonstrated that the [2,3] sigmatropic rearrangement proceeds through chiral, nonracemic free sulfonium ylides rather than through the Co-salen-associated ylides. In this

[†] Dedicated to Professor Shô Itô on the occasion of his 77th birthday.




respect, it has recently been suggested that metal-associated ylides are involved in the rearrangement step with their iodonium and oxonium counterparts.^{7,8} We have recently documented that tandem formation and 1,3-dipolar cycloaddition of keto- or ester-carbonyl ylides under the influence of chiral dirhodium(II) carboxylates incorporating *N*-phthaloyl-(*S*)-amino acids as bridging ligands gives cycloadducts in good yields and with up to 93% ee,⁹ which provides conclusive evidence for the intermediacy of the chiral rhodium(II)-associated carbonyl ylides in the cycloaddition step. Thus, our interest has now been centered on the mechanistic pathway for asymmetric induction in tandem allylic sulfonium ylide formation and [2,3] sigmatropic rearrangement sequence catalyzed by our dirhodium(II) complexes. At the outset, we explored the reaction of (*E*)-cinnamyl phenyl sulfide (**1a**) and ethyl diazoacetate (**2a**) in the presence of 1 mol % of chiral dirhodium(II) carboxylates. The use of dirhodium(II) tetrakis[*N*-phthaloyl-(*S*)-phenylalaninate], Rh₂(S-PTPA)₄, afforded rearrangement products (**5a** and **6a**) as a 62:38

Table 1. Enantioselective Intermolecular Sulfonium Ylide Formation / [2,3] Sigmatropic Rearrangement Catalyzed by Rh(II) Complexes

entry	diazo ester		Rh(II) catalyst	% yield of 5+6 ^a	ratio of 5 : 6 ^b	% ee	
	2a-c	R				5	6
1	2a	Et	Rh ₂ (S-PTPA) ₄	61	62 : 38	0 ^c	0 ^c
2	2a	Et	Rh ₂ (S-PTTL) ₄	60	62 : 38	0 ^c	0 ^c
3	2a	Et	Rh ₂ (S-PTPI) ₄	81	66 : 34	14 ^c	14 ^c
4	2b	<i>t</i> -Bu	Rh ₂ (S-PTPA) ₄	46	80 : 20	<2 ^d	<2 ^d
5	2b	<i>t</i> -Bu	Rh ₂ (S-PTTL) ₄	42	83 : 17	<2 ^d	<2 ^d
6	2b	<i>t</i> -Bu	Rh ₂ (S-PTPI) ₄	61	83 : 17	23 ^d	24 ^d
7	2c	CH(<i>i</i> -Pr) ₂	Rh ₂ (S-PTPI) ₄	61	94 : 6	53 ^d	- ^e

^a Isolated yield. Carbene dimers, maleate and fumarate were obtained as the only detectable by products in all cases. ^b Determined by ¹H NMR after LiAlH₄ reduction and acetylation. ^c Determined by HPLC (Daicel Chiralcel OD). ^d Determined by HPLC (Daicel Chiralcel OJ) after LiAlH₄ reduction. ^e Could not be accurately determined due to an exceptionally high order of diastereoselectivity.

mixture of *erythro* and *threo* isomers in 61% yield, but no asymmetric induction was observed in either case (Table 1, entry 1). A virtually similar result was obtained with the use of dirhodium(II) tetrakis[*N*-phthaloyl-(*S*)-*tert*-leucinate], Rh₂(*S*-PTTL)₄, characterized by a bulky *tert*-butyl group (entry 2). While a significant increase in diastereoselectivity was observed with the sterically demanding *tert*-butyl diazoacetate (**2b**), we were disappointed here again to find that there was little asymmetric induction (entries 4 and 5). After several unsuccessful attempts, it was found that this goal could be achieved by the use of dirhodium(II) carboxamidate complex, dirhodium(II) tetrakis[3(*S*)-phthalimido-2-piperidinonate], Rh₂(*S*-PTPI)₄, which has proven to be the catalyst of choice for intermolecular enantioselective cyclopropanations of styrenes or 1,1-disubstituted alkenes with diazoacetates.¹⁰ A steady increase in diastereo- and enantioselectivity was observed on increasing the steric bulk of the ester alkyl group (entries 3, 6 and 7), in which 2,4-dimethyl-3-pentyl diazoacetate (**2c**) provided the highest values (**5c**:**6c**=94:6, ee of **5c**; 53%) (entry 7).^{11,12} From the fact that *erythro*/*threo* diastereoselectivities are independent of the nature of dirhodium(II) catalysts used but markedly dictated by the size of the ester group (entries 1-3 vs. 4-6), it is evident that the [2,3] sigmatropic rearrangement proceeds through free sulfonium ylides formed by metal dissociation from the dirhodium(II)-bound intermediate (**7**) as mentioned by Katsuki with chiral salen-Co(III) complex.⁶ Based on the further findings that virtually similar enantioselectivities were obtained with *erythro* and *threo* products (**5** and **6**) (entries 1-6), the observed *erythro*/*threo* diastereoselectivity can be rationalized by assessing two envelope transition states (**A**) and (**B**).^{6b} Clearly, the transition state (**A**) leading to *erythro* isomer (**5**) is preferred over the transition state (**B**) to give *threo* isomer (**6**) because of severe steric repulsion between the ester group and the phenyl group in **B**, which also explains a function of the size of the ester group in diazoacetates.



Provided that the [2,3] sigmatropic rearrangement occurs at a faster rate than the racemization at the sulfonium center,^{13,14} the magnitude of enantioselection observed here indicates the level of differentiation between enantiotopic lone pairs on the sulfur atom by the chiral rhodium(II) carbene intermediate (**8**).¹⁵ In this regard, it is worthy of note that the chiral rhodium(II) carbene intermediate (**8**) derived from Rh₂(*S*-PTPI)₄ captures selectively one of the enantiotopic sulfur lone pairs to form the configurationally stable sulfonium ylide, while that from chiral dirhodium(II) carboxylates such as Rh₂(*S*-PTPA)₄ or Rh₂(*S*-PTTL)₄ exhibits virtually no selectivity at this step.

In order to establish the efficacy of using Rh₂(*S*-PTPI)₄ in conjunction with 2,4-dimethyl-3-pentyl diazoacetate (**2c**), we then extended the present protocol to other allylic phenyl sulfides (**1b-e**) than

Table 2. Enantioselective Intermolecular Sulfonium Ylide Formation / [2,3] Sigmatropic Rearrangement Catalyzed by $\text{Rh}_2(\text{S-PTPI})_4$

entry	sulfide 1			products			
	R^1	R^2	R^3	% yield ^a	% ee		
1	1a	Ph	H	H	5,6	61 ^b	53 ^{c,d}
2	1b	H	H	H	9	75	58 ^e
3	1c^f	Me	H	H	10	75 ^g	45 ^{c,h}
4	1d	Me	Me	H	11	72	27 ^e
5	1e	H	H	Me	12	63	19 ^e

^a Isolated yield. ^b Yield of a 94:6 mixture of *erythro* and *threo* products. ^c Ee of *erythro* isomer. ^d Determined by HPLC (Daicel Chiralcel OJ) after LiAlH_4 reduction. ^e Determined by HPLC (Daicel Chiralcel OD). ^f *E:Z* = 93:7. ^g Yield of a 85:15 mixture of *erythro* and *threo* products. ^h Determined by HPLC (Daicel Chiralcel OD) after LiAlH_4 reduction.

cinnamyl phenyl sulfide (**1a**). The results are summarized in Table 2. Allyl phenyl sulfide (**1b**) used as the standard for purposes of comparison provided the highest value (58% ee) reported to date for dirhodium(II)- and copper(I)-catalyzed reactions of achiral allylic sulfides with diazoacetates (entry 2). Lower enantioselectivities were generally observed with phenyl or methyl substitution on the carbon-carbon double bond relative to **1b**, in which a dramatic drop in enantioselectivities was observed with methallyl and prenyl phenyl sulfides (**1d** and **1e**) (entries 4 and 5).

In summary, we have demonstrated the effective use of $\text{Rh}_2(\text{S-PTPI})_4$ as a catalyst for enantioselective tandem allylic sulfonium ylide formation and [2,3] sigmatropic rearrangement sequence, albeit with a limited range of allylic sulfides. It has also been demonstrated that the [2,3] sigmatropic rearrangement proceeds through chiral, nonracemic free sulfonium ylides dissociated from the dirhodium(II) complex. Further extension of the present method to rearrangements *via* oxonium or ammonium ylide formation is currently in progress.

ACKNOWLEDGEMENT

This research was supported in part by a Grant-in-Aid for Scientific Research from the Ministry of Education, Science, Sports and Culture of Japan. We are grateful to Ms. S. Oka and N. Hazama of Center for Instrumental Analysis, Hokkaido University, for mass measurements.

REFERENCES AND NOTES

- R. Brückner, in 'Comprehensive Organic Synthesis,' Vol. 6, ed. by B. M. Trost and I. Fleming, Pergamon Press, Oxford, 1991, pp. 873-908.
- (a) M. P. Doyle, *Chem. Rev.*, 1986, **86**, 919; (b) A. Padwa and M. D. Weingarten, *Chem. Rev.*, 1996,

- 96, 223; (c) M. P. Doyle, M. A. McKervey, and T. Ye, 'Modern Catalytic Methods for Organic Synthesis with Diazo Compounds,' Wiley-Interscience, New York, 1998.
- (a) A.-H. Li, L.-X. Dai, and V. K. Aggarwal, *Chem. Rev.*, 1997, **97**, 2341; (b) M. P. Doyle and D. C. Forbes, *Chem. Rev.*, 1998, **98**, 911.
 - Y. Nishibayashi, K. Ohe, and S. Uemura, *J. Chem. Soc., Chem. Commun.*, 1995, 1245.
 - D. W. McMillen, N. Varga, B. A. Reed, and C. King, *J. Org. Chem.*, 2000, **65**, 2532.
 - (a) T. Fukuda and T. Katsuki, *Tetrahedron Lett.*, 1997, **38**, 3435; (b) T. Fukuda, R. Irie, and T. Katsuki, *Tetrahedron*, 1999, **55**, 649.
 - (a) N. McCarthy, M. A. McKervey, T. Ye, M. McCann, E. Murphy, and M. P. Doyle, *Tetrahedron Lett.*, 1992, **33**, 5983; (b) K. Ito, M. Yoshitake, and T. Katsuki, *Heterocycles*, 1996, **42**, 305; (c) M. P. Doyle, D. G. Ene, D. C. Forbes, and J. S. Tedrow, *Tetrahedron Lett.*, 1997, **38**, 4367; (d) N. Pierson, C. Fernández-García, and M. A. McKervey, *Tetrahedron Lett.*, 1997, **38**, 4705; (e) J. S. Clark, M. Fretwell, G. A. Whitlock, C. J. Burns, and D. N. A. Fox, *Tetrahedron Lett.*, 1998, **39**, 97; (f) M. P. Doyle, D. C. Forbes, M. M. Vasbinder, and C. S. Peterson, *J. Am. Chem. Soc.*, 1998, **120**, 7653.
 - Aggarwal and co-workers have recently shown that metal-associated ylides are involved in allylic sulfonium ylide rearrangements using trimethylsilyldiazomethane as a diazo substrate, in which a marked influence on the diastereoselectivity was observed with the metal catalyst used: V. K. Aggarwal, M. Ferrara, R. Hainz, and S. E. Spey, *Tetrahedron Lett.*, 1999, **40**, 8923.
 - (a) S. Kitagaki, M. Anada, O. Kataoka, K. Matsuno, C. Umeda, N. Watanabe, and S. Hashimoto, *J. Am. Chem. Soc.*, 1999, **121**, 1417; (b) S. Kitagaki, M. Yasugahira, M. Anada, M. Nakajima, and S. Hashimoto, *Tetrahedron Lett.*, 2000, **41**, 5931.
 - (a) N. Watanabe, H. Matsuda, H. Kuribayashi, and S. Hashimoto, *Heterocycles*, 1996, **42**, 537; (b) S. Kitagaki, H. Matsuda, N. Watanabe, and S. Hashimoto, *Synlett*, 1997, 1171.
 - It is noteworthy that a similar trend has been observed in Rh₂(S-PTPI)₄-catalyzed intermolecular cyclopropanations.
 - Representative procedure.* A solution of 2,4-dimethyl-3-pentyl diazoacetate (**2c**) (40.0 mg, 0.22 mmol) in CH₂Cl₂ (1 mL) was added through a syringe pump over 5 h to a refluxing CH₂Cl₂ (1 mL) solution of (*E*)-cinnamyl phenyl sulfide (**1a**) (73.7 mg, 0.33 mmol) and THF adduct of Rh₂(S-PTPI)₄ (2.6 mg, 1 mol %). After stirring at this temperature for an additional 1 h, the solvent was removed *in vacuo*. The residue was purified by column chromatography (silica gel 15 g, 1:2 benzene/hexane) to give **5c** and **6c** as an inseparable mixture of *erythro* and *threo* isomers in a ratio of 94:6 (50.4 mg, 61%); white solid; mp 81-82 °C; [α]_D²³ -68.6 (c=1.53, CHCl₃); IR ν_{max} (film) 2969, 1723 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ: 0.49 (d, *J* = 6.6 Hz, 3H), 0.51 (d, *J* = 6.6 Hz, 3H), 0.65 (d, *J* = 6.6 Hz, 3H), 0.70 (d, *J* = 6.6 Hz, 3H), 1.61-1.78 (m, 2H), 3.78 (dd, *J* = 8.3, 11.2 Hz, 1H), 3.98 (d, *J* = 11.2 Hz, 1H), 4.38 (t, *J* = 5.9 Hz, 1H), 5.08 (d, *J* = 17.2 Hz, 1H), 5.18 (d, *J* = 10.2 Hz, 1H), 6.15 (ddd, *J* = 8.3, 10.2, 17.2 Hz, 1H), 7.15-7.55 (m, 10H); ¹³C NMR (100 MHz, CDCl₃) δ: 16.7, 17.2, 19.1, 19.3, 29.1, 29.6, 51.1, 55.6, 83.8, 117.5, 127.0, 127.7, 128.2, 128.6, 128.8, 132.5,

134.1, 138.4, 141.2, 171.0; HR-EI-MS m/z : Calcd for $C_{24}H_{30}O_2S$ (M^+) 382.1966, found 382.1975. According to the procedure of Katsuki,^{6b} a mixture of *erythro* and *threo* products (**5c** and **6c**) was submitted to reduction with $LiAlH_4$ and acetylation of the resulting alcohols. The *erythro*/*threo* ratio was determined by 1H NMR spectroscopy [methyl signals in acetyl group: δ 1.95 (*erythro*) and 1.99 (*threo*)]. The enantiomeric excess of the *erythro* isomer (**5c**) was determined to be 53% by HPLC analysis of the corresponding alcohols, whereas that of the *threo* isomer (**6c**) could not be determined accurately owing to an exceptionally high order of diastereoselectivity: Daicel Chiralcel OJ, 9:1 hexane/2-propanol, 0.5 mL/min, t_R = 40 min (*erythro* major), t_R = 56 min (*erythro* minor), t_R = 65 min (*threo* major), t_R = 79 min (*threo* minor).

13. B. M. Trost and R. F. Hammen, *J. Am. Chem. Soc.*, 1973, **95**, 962.
14. For racemization of chiral sulfonium ylides, see: (a) D. Darwish and R. L. Tomilson, *J. Am. Chem. Soc.*, 1968, **90**, 5938; (b) S. J. Campbell and D. Darwish, *Can. J. Chem.*, 1974, **52**, 2953.
15. The role of the ester group in enantioselection has not yet been defined; however, it seems likely that the steric effects would give priority to the approach of one of the two enantiotopic sulfur lone pairs to the rhodium(II) carbene center.