

[2□ + 2□] TYPE CYCLOADDITION REACTIONS OF NAPHTHO[*b*]-
CYCLOPROPENE WITH TRITHIOCARBONYL COMPOUNDS TO FORM
1,3-DIHYDRONAPHTHO[2,3-*c*]THIOPHENE DERIVATIVES

Katsuhiro Saito,* Yasuhiro Ohzone, Yoji Kondo, Katsuhiko Ono, and Masakazu Ohkita

Department of Applied Chemistry, Nagoya Institute of Technology, Gokiso-cho,
Showa-ku, Nagoya 466-8555, Japan

Abstract – Reactions of naphtho[*b*]cyclopropene with trithiocarbonyl derivatives in chloroform afforded 1,3-dihydrobenzo[2,3-*c*]thiophene derivatives in moderate yields through a [2□ + 2□] type cycloaddition process. The similar reaction with a dithiocarbonate compound gave a recovery of the starting material. A reaction with thiotropone resulted in a [8□ + 2□] type adduct in 5% yield. Solvent effects on the product yields revealed that the reactions proceeded *via* ionic intermediates.

Naphtho[*b*]cyclopropene (**1**) is a stable colorless crystalline material in spite of its large strain energy resulted from the distorted bond angles of the bridge-head sp² hybridized carbon atoms. However, it is known that compound **1** is highly reactive as a 2□ component in cycloaddition reactions owing to the strained structure and a deficient aromaticity.¹ Recently, the carbenic behavior of **1** was also proposed.² The reactivities of cyclopropene derivatives attached to aromatic rings such as benzocyclopropene,³ 5*H*-cycloprop[*f*]isobenzofuran,⁴ or **1** have been researched. Recently, [4□ + 2□] and [8□ + 2□] types cycloadduct were obtained in reactions of **1** with nitrones and tropones, respectively.⁵

As a series of our research, reactions of **1** with trithiocarbonyl compounds were performed to give 1,3-dihydronaphtho[2,3-*c*]thiophene derivatives. We report here the results of the reactions.

Compound (**1**) and an equimolar amount of dimethyltrithiocarbonate (**2**) in chloroform were refluxed for 3 days. The reaction mixture was separated with column chromatography on silica gel to give a

cycloaddition product (**3**) in 21% yield. The analogous reactions of compound (**1**) with 1,3-dithiol-2-thione (**4**) afforded **5** in moderate yields (**5a**: 43%, **5b**: 55%, **5c**: 66%, and **5d**: 63%). The structures of the products were deduced on the basis of their spectral properties. The molecular structures were ultimately determined by an X-ray structural analysis of **5b** as shown in Fig. 1.

As a π -electron system similar to **4**, thiotropone (**6**)⁸ was also reacted with **1** to afford a $[8\pi + 2\pi]$ type cycloaddition product (**7**) in low yield (5%).






Figure 1. Molecular structure of **5b** (ORTEP).



The reactions of **1** with **4b** in several solvents were examined. The yields of **5b** in the reactions and the solvent parameters⁷ are summarized in the following table. The cycloaddition reaction did not proceed in benzene.

Table. Reactions of **1** with **4b** in solvents and the solvent parameters

Solvent	Time (days)	Yield (%)	Relative Permittivity	Swain's Parameter
chloroform	2	43	4.81	1.15
tetrahydrofuran	3	32	7.58	0.84
ethyl acetate	3	21	6.02	0.79
benzene	3	0	2.28	0.73

The table shows that the polarity of the solvents increases the product yields. The electron densities on the carbon atoms of **1** are known to be high at the bridgehead carbon atoms as shown in Figure 2.⁵






Figure 2. Electron densities of naphtho[*b*]cyclopropene (**1**).

These facts indicate that the reaction of **1** with **2** or **4** proceed through an ionic process as follows. An anionic carbon atom of **1** nucleophilically attacks to the trithiocarbonyl carbon atom (**2**, **4**) to generate a zwitter ionic intermediate (**8**), which then cyclized to the final product (**3**, **5**).



The reaction of **1** with thiotropone (**6**) may proceed *via* a reaction path similar to those of **2** and **4**. Thus the initial product is supposed to be spiro-type intermediate (**9**), which subsequently isomerizes to the final product (**7**).⁵



EXPERIMENTAL

NMR, MS, and IR spectra were measured with Varian GEMINI 200/300, Hitachi 220A, and JASCO FT/IR-5300 spectrometers, respectively. On the separation of reaction mixture, Wakogel[®] C-200 (100–200 mesh; 75–150 μm) was used.

Typical reactions are mentioned below.

Reaction of 1 with 2. A mixture of **1** (140 mg, 1 mmol) and **2** (138 mg, 1 mmol) in chloroform (15 mL) was refluxed for 3 days. The solvent was removed in vacuo. The resulted residue was separated with column chromatography on silica gel (eluent: 30 vol% benzene in *n*-hexane) to give **3** (88 mg, 21%).

3: ^1H NMR (CDCl_3) δ : 2.18 (s, 6H, Me), 4.37 (s, 2H, $-\text{CH}_2-$), 7.45–7.58 (m, 2H), 7.71 (s, 1H), 7.79–7.90 (m, 2H), 8.43 (s, 1H). ^{13}C NMR (CDCl_3) δ : 12.3, 30.6, 74.3, 126.3, 127.4, 128.5, 132.4, 133.4, 137.3, 140.1. IR (KBr): 3056, 2913, 1659 cm^{-1} . MS m/z (rel. intensity): 278 (M^+ , 100), 248, 184, 140.

Reaction of 1 with 4. A mixture of **1** (140 mg, 1 mmol) and **4a–d** (1 mmol) in chloroform (15 mL) was refluxed for 3 days. The solvent was removed in vacuo. The resulted residue was separated with column chromatography on silica gel (eluent: 30 vol% benzene in *n*-hexane) to gave **5**.

5a: ^1H NMR (CDCl_3) δ : 4.47 (s, 2H, $-\text{CH}_2-$), 6.13 (s, 2H, CH), 7.45–7.55 (m, 2H), 7.72 (s, 1H), 7.78–7.95 (m, 2H), 8.42 (s, 1H). ^{13}C NMR (CDCl_3) δ : 37.2, 86.0, 123.1, 123.2, 126.2, 126.3, 127.3, 128.7, 128.9, 132.7, 133.5, 135.6, 140.1. IR (KBr): 3057, 1522 cm^{-1} . MS m/z (rel. intensity): 274 (M^+ , 100), 184, 139.

5b: ^1H NMR (CDCl_3) δ : 2.46 (s, 6H, Me), 4.46 (s, 2H, $-\text{CH}_2-$), 7.49–7.51 (m, 2H), 7.71 (s, 1H), 7.79–7.90 (m, 2H), 8.40 (s, 1H). ^{13}C NMR (CDCl_3) δ : 19.4, 36.6, 83.2, 123.4, 124.2, 125.3, 126.4, 126.9, 127.1, 128.5, 132.4, 133.4, 137.3, 140.0. IR (KBr): 3052, 2912 cm^{-1} . MS m/z (rel. intensity): 366 (M^+ , 96), 231, 184.0, 150. A crystal data: colorless plate, $\text{C}_{16}\text{H}_{14}\text{S}_5$, $M = 366.59$, triclinic, space group $P\bar{1}(\#2)$, $a = 7.2598(8)\text{\AA}$, $b = 7.791(1)\text{\AA}$, $c = 16.714(4)\text{\AA}$, $\beta = 80.07(4)^\circ$, $\gamma = 79.43(3)^\circ$, $\delta = 63.44(3)^\circ$, $V = 826.8(3)\text{\AA}^3$, $Z = 2$, $D_{\text{calcd}} = 1.472\text{ g cm}^{-3}$, $\rho(\text{MoK}\alpha) = 6.90\text{ cm}^{-1}$, $F_{000} = 380$. A total of

3610 reflections for $2\theta_{\max} = 55.0^\circ$ was collected with $I > 2\sigma(I)$ using a Rigaku / MSC Mercury CCD diffractometer (MoK α radiation, $\lambda = 0.71070$ Å) at 296 K. The structure was solved using direct method (SIR97) and refined by full-matrix least-squares analysis giving values of $R = 0.109$, $R_w = 0.173$, $R_1 = 0.075$, $S = 1.48$, $\rho_{\max} / \rho_{\min} = 0.51 / -0.48$ e Å $^{-3}$.

5c: ^1H NMR (CDCl_3) δ : 4.54 (s, 2H, $-\text{CH}_2-$), 7.46 (t, $J = 7.4$ Hz, 4H), 7.53 (dd, $J = 6.2, 3.2$ Hz, 2H), 7.60 (tt, $J = 1.4, 7.4$ Hz, 2H), 7.73 (s, 1H), 7.82 (dd, $J = 5.9, 3.2$ Hz, 1H), 7.97 (dd, $J = 7.1, 1.4$ Hz, 4H), 8.03 (dd, $J = 5.8, 3.6$ Hz, 1H), 8.86 (s, 1H). ^{13}C NMR (CDCl_3 , 300 MHz) δ : 37.2, 86.3, 122.8, 123.1, 123.2, 126.2, 126.3, 126.3, 127.2, 127.4, 127.8, 128.8, 132.7, 133.5, 134.2, 135.6, 136.6, 140.9, 187.2. IR (KBr): 3400, 3055, 1684 cm^{-1} . MS m/z (rel. intensity): 546 (M^+ , 1), 216, 184, 139.

5d: ^1H NMR (CDCl_3) δ : 3.30–3.43 (m, 4H, $-\text{CH}_2-$), 4.46 (s, 2H, $-\text{CH}_2-$), 7.46–7.55 (m, 2H), 7.71 (s, 1H), 7.78–7.83 (m, 1H), 7.86–7.91 (m, 1H), and 8.40 (s, 1H). ^{13}C NMR (CDCl_3) δ : 26.0, 30.7, 36.5, 111.7, 123.3, 123.5, 124.8, 126.3, 127.1, 128.4, 132.2, 133.4, 137.7, 113.8. IR (KBr): 3056, 1530 cm^{-1} . MS m/z (rel. intensity): 365 (M^+ , 100), 224, 139.

Reaction of 1 with 6. A mixture of **1** (281 mg, 2 mmol), P_2S_5 (3.80 g, 17 mmol), and tropone (1.00 g, 9.5 mmol) in dichloromethane (7 mL) was treated with NEt_3 (1.07 g, 10 mmol) at 0°C under a nitrogen stream. The mixture was stirred for 1 h, then filtrated in the reduced pressure and then the solvent was removed on a rotary evaporator. The resulted residue was separated with column chromatography on silica gel (eluent: 40 vol% benzene in *n*-hexane) to give **7** (28 mg, 5.4%).

7: ^1H NMR (CDCl_3) δ : 3.15 (d, $J = 6.0$ Hz, 1H), 3.94 (d, $J = 14.3$ Hz, 1H, $-\text{CH}_2-$), 4.10 (d, $J = 14.3$ Hz, 1H, $-\text{CH}_2-$), 5.28 (dd, $J = 9.1, 6.3$ Hz, 1H), 6.34 (d, $J = 9.1$ Hz, 1H), 6.53 (t, $J = 3.2$ Hz, 1H), 6.72 (t, $J = 3.2$ Hz, 2H), 7.48–7.52 (m, 2H), 7.72 (s, 1H), 7.76–7.86 (m, 4H). ^{13}C NMR (CDCl_3) δ : 34.1, 44.5, 122.3, 124.5, 125.0, 125.4, 125.8, 126.0, 127.0, 127.1, 127.2, 127.3, 128.5, 129.7, 130.3, 131.8, 132.9, 136.7. IR (KBr): 3002, 1578, 1501, 914, 874, 758, 721 cm^{-1} . MS m/z (rel. intensity): 262 (M^+ , 100), 261, 229, 228, 185.

REFERENCES

1. B. Halton and M. C. Banwell, "Cyclopropenes" in "The Chemistry of Functional Groups" ed. by Z. Rappoport, Wiley, Chichester, 1978, pp. 1223–1340; B. Halton, *Chem. Rev.*, 1989, **89**, 1161; P. C. Hiberty, G. Ohanessian, and F. Delbecq, *J. Am. Chem. Soc.*, 1985, **107**, 3095; C. Wentrup, C. Mayor, J. Becker, and H. J. Linder, *Tetrahedron*, 1985, **41**, 1601.
2. W. Pan, M. Jones, Jr., B. East, and P. Lahti, *Tetrahedron Lett.*, 1998, **39**, 1505.

3. K. Saito, H. Ishihara, and S. Kagabu, *Bull. Chem. Soc. Jpn.*, 1987, **60**, 4141; S. Kagabu and K. Saito, *Tetrahedron Lett.*, 1988, **29**, 675; S. Kagabu and T. Inoue, *Chem. Lett.*, 1989, 2181; B. Halton, and S. G. Russell, *Aust. J. Chem.*, 1990, **43**, 2099; S. Kagabu, K. Saito, H. Watanabe, K. Takahashi, and K. Wada, *Bull. Chem. Soc. Jpn.*, 1991, **64**, 106; K. Saito, K. Ito, K. Takahashi, and S. Kagabu, *Org. Prep. Proc. Internat.*, 1991, **23**, 196; K. Ito and K. Saito, *Recent Res. Devel. Pure & Applied Chem.*, 2000, **4**, 81.
4. I. J. Anthony and D. Wege, *Tetrahedron Lett.*, 1987, **28**, 4217.
5. S. Ando, J. Imamura, A. Hattori, M. Tajitsu, and K. Saito, *Heterocycles*, 1998, **48**, 1769; K. Saito, S. Ando, and Y. Kondo, *ibid.*, 2000, **53**, 2601; K. Saito, K. Ono, N. Ito, N. Tada, and S. Ando, *ibid.*, 2002, **57**, 235.
6. K. Saito, K. Ito, C. Kabuto, and K. Takahashi, *Bull. Chem. Soc. Jpn.*, 1991, **64**, 2387.
7. C. G. Swain, R. B. Moseley, and D. E. Bown, *J. Am. Chem. Soc.*, 1955, **77**, 3721.
8. T. Machiguchi, T. Hasegawa, H. Otani, and S. Yamabe, *J. Am. Chem. Soc.*, 1994, **116**, 407; K. Ito, K. Saito, S. Takeuchi, and K. Takahashi, *Heterocycles*, 1992, **34**, 1415.