

NUCLEOPHILIC SUBSTITUTION REACTION OF 5-*t*-BUTYL-2-METHOXY-3*H*-AZEPINE WITH ALKOXIDES AND ALKYL LITHIUM REAGENTS: A FORMATION OF BIS(5-*t*-BUTYL-3*H*-AZEPIN-2-YL)-METHANE HAVING A VINAMIDINE CONJUGATION

Kyosuke Satake,* Yasuhiro Kubota, Hideki Okamoto, and Masaru Kimura

Department of Chemistry, Faculty of Science, Okayama University

Tsushima-Naka 3-1-1, Okayama, 700-8530, Japan


satake@cc.okayama-u.ac.jp

Abstract – Reaction of 5-*t*-butyl-2-methoxy-3*H*-azepine (**2**) with nucleophiles (alkoxides or *t*-butyllithium) gave respective substitution product at 2-position of the ring. Unexpectedly, when methyllithium was used as a nucleophile, bis(5-*t*-butyl-3*H*-azepin-2-yl)methane (**7**), the structure of which found to be tautomeric vinamidine (**7a,b**), was formed. Tautomerization between **7** and **7a,b** was characterized spectroscopically and theoretically based on the levels of B3LYP/6-31G(d).

Among the four possible azepine isomers (1*H*-, 2*H*-, 3*H*-, and 4*H*-azepine), 2-alkoxy- or 2-alkylamino-3*H*-azepines have been studied extensively,¹ because the synthesis of such 3*H*-azepines has been well explored.² In addition, these 3*H*-azepines were relatively stable making exploration of their reactivities


possible, although unsubstituted 3*H*-azepine has been reported to be a labile substance.³ Nucleophilic substitution reactions at the 2-position of the ring with amines or active methylene compounds have been reported.⁴ We report here alternative nucleophilic substitution reactions with alkoxide and alkyllithium reagents.

Ring expansion reaction of 4-*t*-butylnitrobenzene (**1**) was carried out by heating a methanol solution of **1** and tributylphosphine in a sealed tube at 150°C for 24 h. Subsequent distillation under reduced pressure of the reaction mixture gave **2** in 74% yield.⁵ In order to examine the nucleophilic substitution reaction by an alkoxide ion, an ethanol solution of **2** and excess of sodium ethoxide was refluxed for 24 h. An effective ether exchange reaction occurred to give 2-ethoxy-5-*t*-butyl-3*H*-azepine (**3a**) in 78% yield. Similar procedures for **2** using propanoxide, isopropanoxide, and triethylene monoglycoxide gave **3b** (74%), **3c** (62%), and **3d** (24%), respectively.⁶



We next examined the possibility for the synthesis of 2,5-dialkyl-3*H*-azepines by nucleophilic alkyl substitution reaction. Previously, we have reported the synthesis of 2,5-dialkyl-3*H*-azepines by means of a demethoxycarbonylation reaction of methyl 2,5-dialkyl 1*H*-azepine-1-carboxylate.⁷ To inspect an alternative and effective synthetic method of 2,5-di-*t*-butyl-3*H*-azepine (**4**), *t*-butyllithium was used as a nucleophile. To a dry ether solution of **2**, excess of *t*-butyllithium ether solution was added at room temperature with stirring. After quenching the reaction mixture with water, di-*t*-butyl derivative (**4**) was obtained from the organic layer in 44% yield. On the other hand, when methyllithium was used as a nucleophile under similar conditions, the reaction gave not only the expected 2-methyl-5-*t*-butyl-3*H*-azepine (**5**) in 12% yield, but also 2,3-dihydro-2,2-dimethyl-5-*t*-butyl-1*H*-azepine (**6**) and bis(5-*t*-butyl-

3*H*-azepin-2-yl)methane (**7**) in 37 and 25% yields. Structures for **4** and **5** were confirmed by comparing the spectral data with those of previously reported.⁷ The formation of **6** is presumed to doubly occurred nucleophilic attacks of methyllide on **2**, because the reaction of **5** with methyllithium also gave **6**. The enamine structure of **6** was supported by $\nu_{\text{N-H}}$ band at 3295 cm^{-1} and three olefinic proton signals at $\delta 4.63$ (dd, $J = 10.0, 1.7\text{ Hz}$, 1H), 5.37 (dt, $J = 6.8, 1.7\text{ Hz}$, 1H), and 6.12 (d, $J = 10.0\text{ Hz}$, 1H) ppm in the IR and ^1H NMR spectra, respectively.⁸



Another sequential nucleophilic reaction between **2** and the carbanion from **5** gave **7** which showed m/z 311.2483 $[(\text{M}+\text{H})^+ \text{ calcd for } \text{C}_{21}\text{H}_{31}\text{N}_2: 311.2487]$ in HRMS (FAB). The NMR spectral signals could not be assigned to **7** due to an ^1H NMR signal of olefinic methyne at $\delta 4.42$ (s, 1H) ppm and an acidic proton at $\delta 12.0$ (br, 1H) ppm and the ^{13}C NMR signal at $\delta 93.9$ (d) ppm.⁹ Reasonable assignment for the NMR spectra was made possible by considering the tautomers (**7a,b**), which are considered to be promoted by intramolecular hydrogen bond. As well, the stretching band for N-H at 3046 cm^{-1} indicates the existence of an effective hydrogen bond. D_2O exchange experiment showed instant disappearance of the signal at $\delta 12.0$ ppm (N-H) and then gradually decreasing of the signal intensity at $\delta 4.42$ ppm (central $=\text{CH}-$) with a half-life period by 81 min at 22°C . The observed independent rates for D_2O exchange suggest the interconversion between **7a** and **7b** is not only attributed to 1,3-prototropy but mainly attributed to a degenerate 1,5-hydrogen shift between both rings with lower activation energy (E_a).

In order to investigate an energy profile for a diimine (**7**), tautomers (**7a,b**) and the transition structure of 1,5-hydrogen shift (**7_{TS}**), an *ab initio* energy analysis was carried out using GAUSSIAN98 program.¹⁰ Theoretical levels of B3LYP/6-31G(d)¹¹ were applied and zero-point-energy correction was made on all calculations.¹² Calculated energy for **7**, **7a,b**, and **7_{TS}** were -927.2880, -927.2971, and -927.2909 Hartrees, respectively. The vibrational analysis for the optimized **7_{TS}** showed only one imaginary frequency by -1431.1 cm⁻¹. The six central hexagonally arranged atoms of **7a,b** and **7_{TS}** are on a plane; each of them corresponds to the classical and nonclassical expression for a vinamidine conjugation,¹³ respectively. Calculated internuclear distances between central hydrogen and both sides of nitrogen of **7a,b** are 1.03 and 1.88 Å. In the case of **7_{TS}**, the location of the hydrogen is on a C₂ symmetry axis of the molecule with an internuclear distance of 1.28 Å. Both the transition structure (**7_{TS}**) and the diimine (**7**) are less stable than **7a,b** by 3.9 and 5.7 kcal/mol, respectively. This explains that the D₂O exchange rates for N–H proton and central C–H proton are attributed to the degenerate 1,5-hydrogen shift (calcd Ea = 3.8 kcal/mol) and 1,3-prototropy (calcd Ea > 5.7 kcal/mol), respectively (Figure 1).




Figure 1 Calculated energy profile for **7**, **7_{TS}**, and **7a,b**.

We are interested in the character of the unique vinamidine conjugation proposed here, which is incorporated in a seven-membered conjugated olefinic system. Further investigation for the chemistry of bisazepinylmethane is underway.

We thank the SC-NMR Laboratory of Okayama University for ^1H and ^{13}C NMR spectral measurements.

REFERENCES AND NOTES

1. L. Wolf, *Liebigs Ann. Chem.*, 1912, **394**, 59; R. Huisgen, *Angew. Chem.*, 1955, **67**, 756; R. Huisgen and D. Vossius, *M. Appl. Chem. Ber.*, 1958, **91**, 1; W. von E. Doering and R. A. Odum, *Tetrahedron*, **1966**, 81; R. Purvis, R. K. Smalley, W. A. Strachan, and H. Suschitzky, *J. Chem. Soc., Perkin Trans. 1*, **1978**, 191.
2. J. I. G. Cadogan and R. K. Mackie, *J. Chem. Soc., C*, **1969**, 2819; T. de Bore, J. I. G. Cadogan, H. M. McWilliam, and A. G. Rowley, *J. Chem. Soc., Perkin Trans. 2*, **1975**, 554; M. Masaki, K. Fukui, and J. Kita, *Bull. Chem. Soc. Jpn.*, 1977, **50**, 2013.
3. E. Vogel, H.-J. Altenbach, J.-M. Drossard, H. Schmickler, and H. Stegelmeier, *Angew. Chem.*, 1980, **92**, 1053; *Angew. Chem., Int. Ed. Engl.*, 1980, **19**, 1015.
4. S. Batori, R. Gompper, J. Meier, and H.-U. Wagner, *Tetrahedron*, 1988, **44**, 3309; H. A. Daboun and M. A. A. Aziz, *Heterocycles*, 1982, **19**, 1375.
5. Selected data for **2**; colorless oil (bp 105–109°C); ^1H NMR (200 MHz, CDCl_3) δ 1.10 (s, 9H), 2.59 (d, J = 6.8 Hz, 2H), 3.71 (s, 3H), 5.15 (dt, J = 6.8, 1.0 Hz, 1H), 6.15 (dd, J = 8.8, 1.0 Hz, 1H), 6.94 (d, J = 8.8 Hz, 1H); ^{13}C NMR (50 MHz, CDCl_3) δ 30.0 (q), 32.5 (t), 34.5 (s), 54.4 (q), 110.1 (d), 115.5 (d), 137.4 (d), 148.6 (s), 154.8 (s); IR (neat) ν_{max} 1628 cm^{-1} (C=N); UV-Vis λ_{max} (EtOH) 254 nm ($\log \epsilon$ = 3.61).
6. Selected data for **3a**; ^1H NMR (200 MHz, CDCl_3) δ 1.10 (s, 9H), 1.26 (t, J = 7.2 Hz, 3H), 2.54 (d, J = 6.8 Hz, 2H), 4.12 (q, J = 7.2 Hz, 2H), 5.15 (dt, J = 6.8, 1.4 Hz, 1H), 6.13 (dd, J = 8.6 Hz, 1.4 Hz, 1H), 6.92 (d, J = 8.6 Hz, 1H); IR ν_{max} (neat) 1626 cm^{-1} (C=N); UV-Vis λ_{max} (EtOH) 254 nm ($\log \epsilon$ = 3.77); **3b**; ^1H NMR (200 MHz, CDCl_3) δ 0.89 (t, J = 7.2 Hz, 3H), 1.06 (s, 9H), 1.61 (dq, J = 7.2, 6.8 Hz, 2H), 2.50 (d, J = 6.7 Hz, 2H), 3.98 (t, J = 6.8 Hz, 2H), 5.11 (dt, J = 6.7, 1.4 Hz, 1H), 6.09 (dd, J = 8.8, 1.4 Hz, 1H), 6.88 (d, J = 8.8 Hz, 1H); IR ν_{max} (neat) 1624 cm^{-1} (C=N); UV-Vis λ_{max} (EtOH) 254 nm ($\log \epsilon$ = 3.70); **3c**; ^1H NMR (200 MHz, CDCl_3) δ 1.10 (s, 9H), 1.19 (d, J = 6.3 Hz, 6H), 2.51 (d, J = 7.4 Hz, 2H), 4.99 (hept, J = 6.3 Hz, 1H), 5.12 (dt, J = 7.4, 1.2 Hz, 1H), 6.12 (dd, J = 8.8, 1.2 Hz, 1H), 6.92 (d, J = 8.8, 1H); IR ν_{max} (neat) 1620 cm^{-1} (C=N); UV-Vis λ_{max} (EtOH) 254 nm ($\log \epsilon$ = 3.44); **3d**; ^1H NMR (200 MHz, CDCl_3) δ 1.10 (s, 9H), 2.58 (d, J = 6.6 Hz, 2H), 3.66–3.75 (m, 10H), 4.25 (t, J =

- 4.7 Hz, 2H), 5.15 (t, $J = 6.6$ Hz, 1H), 6.15 (d, $J = 8.6$ Hz, 1H), 6.90 (d, $J = 8.6$ Hz, 1H); IR λ_{max} (neat) 1624 cm^{-1} (C=N); UV-Vis λ_{max} (EtOH) 254 nm ($\log \epsilon = 3.52$).
7. K. Satake, R. Okuda, M. Hashimoto, Y. Fujiwara, H. Okamoto, M. Kimura, and S. Morosawa, *J. Chem. Soc., Perkin Trans. 1*, **1994**, 1753.
8. Selected data for **6**; ^1H NMR (200 MHz, CDCl_3) δ 1.07 (s, 6H), 1.09 (s, 9H), 2.08 (d, $J = 6.8$ Hz, 2H), 3.80 (br s, 1H), 4.63 (dd, $J = 10.0, 1.7$ Hz, 1H), 5.37 (dt, $J = 6.8, 1.7$ Hz, 1H), 6.12 (d, $J = 10.0$ Hz, 1H); ^{13}C NMR (50 MHz, CDCl_3) δ 29.9 (q), 30.0 (q), 35.2 (s), 42.6 (t), 55.7 (s), 94.5 (d), 113.0 (d), 132.0 (d), 148.3 (s); IR ν_{max} (neat) 3295 cm^{-1} (NH).
9. Selected data for **7**; yellow needles (mp $145\text{--}146^\circ\text{C}$); ^1H NMR (200 MHz, CDCl_3) δ 1.09 (s, 18H), 2.63 (d, $J = 6.8$ Hz, 4H), 4.42 (s, 1H), 5.24 (dt, $J = 6.8, 1.2$ Hz, 2H), 5.85 (dd, $J = 9.0, 1.2$ Hz, 2H), 6.83 (d, $J = 9.0$ Hz, 2H), 12.0 (br s, 1H); ^{13}C NMR (50 MHz, CDCl_3) δ 29.9 (q), 34.7 (s), 36.2 (t), 93.9 (d), 113.4 (d), 114.1 (d), 134.8 (d), 148.6 (s), 155.4 (s); IR ν_{max} (KBr) 3046 cm^{-1} (NH); UV-Vis λ_{max} (EtOH) 380 nm ($\log \epsilon = 4.40$).
10. *Gaussian 98, Revision A.9*, M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, V. G. Zakrzewski, J. A. Montgomery, Jr., R. E. Stratmann, J. C. Burant, S. Dapprich, J. M. Millam, A. D. Daniels, K. N. Kudin, M. C. Strain, O. Farkas, J. Tomasi, V. Barone, M. Cossi, R. Cammi, B. Mennucci, C. Pomelli, C. Adamo, S. Clifford, J. Ochterski, G. A. Petersson, P. Y. Ayala, Q. Cui, K. Morokuma, D. K. Malick, A. D. Rabuck, K. Raghavachari, J. B. Foresman, J. Cioslowski, J. V. Ortiz, A. G. Baboul, B. B. Stefanov, G. Liu, A. Liashenko, P. Piskorz, I. Komaromi, R. Gomperts, R. L. Martin, D. J. Fox, T. Keith, M. A. Al-Laham, C. Y. Peng, A. Nanayakkara, M. Challacombe, P. M. W. Gill, B. Johnson, W. Chen, M. W. Wong, J. L. Andres, C. Gonzalez, M. Head-Gordon, E. S. Replogle, and J. A. Pople, Gaussian, Inc., Pittsburgh PA, 1998.
11. P. J. Stevens, F. J. Devlin, C. F. Chabowski, and M. J. Frisch, *J. Phys. Chem.*, 1994, **98**, 11623.
12. C. W. Bauschlicher, Jr. and H. Partridge, *J. Chem. Phys.*, 1995, **103**, 1788; M. W. Wong, *Chem. Phys. Lett.*, 1996, **256**, 391; A. P. Scott and L. Radom, *J. Phys. Chem.*, 1996, **100**, 16502.
13. Review for vinamidine conjugations; D. Lloyd and H. McNab, *Angew. Chem., Int. Ed. Engl.*, 1976, **15**, 459.