

STUDIES ON THE CONSTITUENTS OF INDONESIAN *BORRERIA*
*LATIFOLIA*¹

Kohei Kamiya,^{a,b} Yasuhiro Fujita,^a Yasuhisa Saiki,^a Endang Hanani,^c Umar
Mansur,^c and Toshiko Satake^{*a,b}

Faculty of Pharmaceutical Sciences,^a and High Technology Research Center,^b
Kobe Gakuin University, Nishi-Ku, Kobe, 651-2180, Japan and Jursan Farmasi,
FMIPA, University of Indonesia,^c Depok Baru, Jawa Barat, Indonesia

Abstract - Seven known iridoid glycosides (**1~7**), together with one known
diterpenoid (**8**) were isolated from the aerial parts of *Borreria latifolia* K. Schum.
(Rubiaceae) collected in Indonesia, and their structures were identified by
spectroscopic methods.

Borreria latifolia is distributed widely throughout tropical Asia and is used as food or feed.² Among the
members of this genus, *B. articularis*³ is used to treat headache, gallstones and in a poultice upon the
abdomen of very small children with stomachache in China. *B. osimoides*³ heal wounds in Indonesia. As a
continuation of the studies on the traditional medicine (Jamu) and the medicinal resources in Indonesia,
we investigated the chemical constituents of *B. latifolia* collected there.

The methanol extract of the aerial parts of *B. latifolia* was passed over active charcoal packed in a column.
The resulting fractions were chromatographed repeatedly on silica gel, Sephadex LH-20, MCI-gel and
ODS column as described in the experimental section to isolate seven known iridoid glycosides (**1~7**),

together with acyclic diterpenoid, phytol (**8**). Here we report the isolation and identification of these compounds.

Compound (**1**) showed a brilliant reddish violet spot on the TLC upon heating with 10% sulfuric acid, before this spot subsequently changed to black, suggesting that **1** was an iridoid compound.⁴ In the ¹H-NMR spectrum, we observed the following signals: the doublet proton signal at δ 5.08 (1H, *J*=8.9 Hz) due to the presence of characteristic H-1 of the iridoid compound, two olefinic proton signals at δ 7.65 (1H, d, *J*=1.2 Hz) and δ 6.01 (1H, d, *J*=1.9 Hz), one hydroxymethylene proton signal at δ 4.95 and 4.80 (each, 1H, br d, *J*=15.6 Hz), hydroxymethine proton signal at δ 4.83 (1H, m), two methine proton signals at δ 3.02 (1H, ddd, *J*=7.8, 6.2, 1.2 Hz) and δ 2.63 (1H, dd, *J*=8.9, 7.8 Hz), and one acetyl proton signal at δ 2.10 (3H, s). In addition, in the ¹³C-NMR spectrum, we observed these signals: hydroxymethine carbon signal at δ 63.7, α,β-unsaturated carboxylic carbon at δ 172.5, double bond carbon signals at δ 155.2 and 108.4, the olefinic carbon signals at δ 131.8 and 145.7, and acetyl carbon signals at δ 172.4 and 20.8. Further, characteristic glucopyranosyl carbon signals and the coupling constant (*J*=7.8 Hz) of the anomeric proton signal at δ 4.74 (d) suggested glucose to be β-linked to the hydroxyl group at the C-1 in iridoid. Therefore, compound (**1**) was elucidated as asperulosidic acid according to values reported in the literature.⁵

Compounds (**2**) and (**3**) were identified by the comparison of the ¹H- and ¹³C-NMR spectra with those of **1** to be deacetylasperulosidic acid⁶ and scandoside,⁶ respectively. Compound (**4**) was identified by the comparison of the ¹H- and ¹³C-NMR spectra with those of **2** to be methyl deacetylasperulosidate.⁷

Compound (**5**) was identified by the comparison of the ¹H- and ¹³C-NMR spectra with those of **4** to be daphylloside.⁸ Compound (**6**) was identified by the comparison of the ¹H- and ¹³C-NMR spectra with those of **3** to be 6-*O*-acetylscandoside.

In the ¹H-NMR spectrum of compound (**7**), the characteristic proton signal at δ 5.15 (1H, d, *J*=6.1 Hz) due to the presence of hydroxymethine at H-1 in iridoid, and one olefinic proton signal at δ 7.45 (1H, d, *J*=1.4 Hz) were observed, but not another olefinic proton signal at C-7. Furthermore, proton signals at δ 2.27 (1H, ddd, *J*=13.4, 6.8, 1.6 Hz), δ 1.54 (1H, ddd, *J*=13.4, 9.6, 4.4 Hz) and δ 4.29 (1H, deformed t) were

detected. These findings indicated the existence of a hydroxyl group at C-7 as the loganin skeleton, which was supported by the ^{13}C -NMR spectral data. Based on the coupling constant ($W_{1/2}=3.1$) of the carbonyl methine at H-7, the hydroxyl group at C-7 was a β -configuration. In the comparison of ^{13}C -NMR of **7** with that of loganin, the existence of a carbon signal at δ 62.3 (t), because of hydroxymethylene at C-10 in **7**, indicated that compound (**7**) was 10-hydroxyloganin.⁹

Compound (**8**) was obtained as colorless oil. The IR spectrum showed absorption band (1668 cm^{-1}) due to double bond, together with those due to methyl, methylene groups and hydroxyl group. In the ^{13}C -NMR, double bond carbons (δ 136.9 and 126.3) and hydroxymethylene carbon (δ 59.0) were exhibited, besides signals of methyl (5), methylene (9), methine (3). In the ^1H -NMR, olefinic proton signal at δ 5.74 (1H, tq, $J=5.3, 1.3\text{ Hz}$), hydroxymethylene proton signals at δ 4.45 (2H, d, $J=6.5\text{ Hz}$), isopropyl proton signals at δ 1.51 (1H, sep, $J=6.8\text{ Hz}$) and 0.90, 0.88 (each 3H, d, $J=6.8\text{ Hz}$), methyl proton signal linked double bond at δ 1.69 (3H, br s), and two methyl proton signals at δ 0.88 and 0.87 (each 3H, s) were showed. From these data, compound (**8**) was considered as the acyclic diterpenoid bearing isopropyl group and hydroxymethyl group, to identify as phytol with literature values.¹⁰




Figure 1 Chemical Structures of Compounds (**1-8**)

Table 1 ^{13}C -NMR Spectral Data for Compounds (1-7) in CD_3OD

	1	2	3	4	5	6	7
C-1	101.2	101.5	98.9	101.6	101.3	93.4	99.0
C-3	155.2	155.4	153.9	155.3	155.3	150.2	152.9
C-4	108.4	108.5	111.0	108.3	108.1	106.2	113.0
C-5	42.3	42.8	47.1	42.7	42.4	37.5	33.5
C-6	75.3	75.4	82.5	75.4	75.3	86.3	43.1
C-7	131.8	130.0	130.0	129.9	131.9	129.0	73.3
C-8	145.7	151.4	147.4	151.5	145.9	144.2	42.2
C-9	46.2	45.9	46.1	45.9	46.3	45.3	49.9
C-10	63.7	61.7	61.1	61.7	63.7	61.9	62.3
C-11	172.5	170.8	171.9	169.4	169.3	172.5	169.4
C-1'	100.5	100.5	100.3	100.5	100.6	100.1	100.6
C-2'	74.8	75.0	74.7	75.0	74.9	74.6	74.8
C-3'	78.2	78.5	78.3	78.5	78.5	78.4	78.3
C-4'	71.4	71.7	71.5	71.7	71.6	71.6	71.5
C-5'	77.7	77.8	77.8	77.9	77.9	77.9	78.0
C-6'	62.8	62.8	62.6	62.9	63.0	62.8	62.3
CH_3O				51.7	51.8		51.7
CH_3CO	172.4				172.5	172.2	
CH_3CO	20.8				20.7	20.6	

EXPERIMENTAL

General Procedures and Plant Material Optical rotations were measured using a Jasco DIP-1000 digital polarimeter. IR and UV spectra were measured on a Shimadzu FT-IR 8300 infrared spectrometer and a Hitachi U-3000 spectrometer, respectively. The NMR spectra were recorded in CD_3OD or $\text{C}_5\text{D}_5\text{N}$ with TMS as internal standard on a Bruker DPX-400. Column chromatography was carried out on Kieselgel 60 (70-230 mesh, Merck Co.), Sephadex LH-20 (Pharmacia Co.). Medium pressure liquid chromatography (MPLC, micro pump KP-7, Kusano Scientific Co. Tokyo) was carried out on a CIG column [ODS (C-18)]. HPLC (JASCO HPLC system; pump:880-PL, detector:875 UV/VIS detector) was

carried out on a SiO₂ packed column (20 mm i. d. × 250 mm, YMC Co.).

Borreria latifolia K. Schum. was collected in July, 1992 at Dien in Indonesia and the plant was identified by Y. Saiki. A voucher specimen has been deposited in the Herbarium of Kobe Gakuin University.

Isolation of Compounds (1~8) The air-dried aerial parts of *B. latifolia* (860 g) was extracted 7 times with methanol (3 L) under 60 °C for 8 h. The combined methanol extracts were passed over active charcoal packed in a column (fraction I), and were eluted with methanol (fraction II) and 30% CHCl₃/ MeOH (fraction III), successively.

Fraction I (66.18 g) was subjected repeatedly to column chromatography on SiO₂ (CHCl₃-MeOH gradient solvent system), Sephadex LH-20 (MeOH : H₂O, 2 : 1) and ODS (Rp-18, MeOH : H₂O, 1 : 2) to yield compounds **(1)** (203.5 mg), **(2)** (30.0 mg), **(3)** (70.9 mg), **(4)** (37.2 mg), **(5)** (37.8 mg), **(6)** (24.0 mg), and **(7)** (41.3 mg). Fraction III (13.55 g) was subjected to repeat column chromatography on SiO₂ (CHCl₃-MeOH gradient solvent system) and Sephadex LH-20 (CHCl₃ : MeOH, 2 : 1), and finally was purified by HPLC (hexane : EtOAc, 9 : 1) to give compound **(8)** (95.7 mg).

Compound (1) (Asperulosidic acid) A white amorphous powder, anthrone / H₂SO₄ reaction positive, $[\alpha]_D^{26} +22.7^\circ$ (c=2.0, MeOH), IR $\nu_{\max}$ (KBr) cm⁻¹ : 3448, 1728, 1685, 1636; UV $\lambda_{\max}$ (MeOH) nm (log ϵ) : 201.5 (3.8), 239.1 (4.0); ¹H-NMR (δ , CD₃OD) : 5.08 (1H, d, *J*=8.9 Hz, H-1), 7.65 (1H, d, *J*=1.2 Hz, H-3), 3.02 (1H, ddd, *J*=7.8, 6.2, 1.2 Hz, H-5), 4.83 (1H, m, H-6), 6.01 (1H, d, *J*=1.9 Hz, H-7), 2.63 (1H, dd, *J*=8.9, 7.8 Hz, H-9), 4.95, 4.80 (each 1H, d, *J*=15.6 Hz, H-10), 2.10 (3H, s, COCH₃), 4.74 (1H, d, *J*=7.8 Hz, H-1'); ¹³C-NMR : Table1

Compound (2) (Deasperulosidic acid) A white amorphous powder, anthrone / H₂SO₄ reaction positive, $[\alpha]_D^{27} +4.13^\circ$ (c=3.0, MeOH). IR $\nu_{\max}$ (KBr) cm⁻¹ : 3432, 1685, 1670, 1636. UV $\lambda_{\max}$ (MeOH) nm (log ϵ) : 202.0 (3.7), 233.0 (3.8); ¹H-NMR (δ , CD₃OD): 5.05 (1H, d, *J*=8.9 Hz, H-1), 7.65 (1H, d, *J*=1.2 Hz, H-3), 3.00 (1H, ddd, *J*=7.7, 6.4, 1.2 Hz, H-5), 4.82 (1H, m, H-6), 6.02 (1H, d, *J*=1.9 Hz, H-7), 2.58 (1H, dd, *J*=8.9, 7.7 Hz, H-9), 4.46, 4.21 (each 1H, d, *J*=15.6 Hz, H-10), 4.72 (1H, d, *J*=7.8 Hz, H-1');

¹³C-NMR : Table I

Compound (3) (Scandoside) A white amorphous powder, anthrone / H₂SO₄ reaction positive, $[\alpha]_D^{26}$ -32.7 ° (c=0.7, MeOH). IR $\nu_{\max}$ (KBr) cm⁻¹: 3432, 1682, 1670, 1636; UV $\lambda_{\max}$ (MeOH) nm (log ϵ): 201.6 (3.8), 233.7 (3.9); ¹H-NMR (δ , CD₃OD): 5.09 (1H, d, J =6.7 Hz, H-1), 7.54 (1H, d, J =1.1 Hz, H-3), 5.82 (1H, d, J =1.0 Hz, H-7), 2.96 (1H, dd, J =9.0, 7.7 Hz, H-9), 4.35, 4.10 (each, 1H, d, J =15.4 Hz, H-10), 4.70 (1H, d, J =7.9 Hz, H-1'); ¹³C-NMR : Table 1.

Compound (4) (Methyl deacetylasperulosidate) A white amorphous powder, anthrone / H₂SO₄ reaction positive, $[\alpha]_D^{27}$ +17.5 ° (c=0.7, MeOH); IR $\nu_{\max}$ (KBr) cm⁻¹: 3424, 1696, 1670, 1634; UV $\lambda_{\max}$ (MeOH) nm (log ϵ): 201.1 (3.7), 234.9 (3.9); ¹H-NMR (δ , CD₃OD): 5.05 (1H, d, J =9.0 Hz, H-1), 7.65 (1H, d, J =1.6 Hz, H-3), 3.02 (1H, ddd, J =7.3, 6.0, 1.6 Hz, H-5), 6.02 (1H, d, J =2.1 Hz, H-7), 2.56 (1H, dd, J =9.0, 7.4 Hz, H-9), 4.45, 4.20 (each, 1H, d, J =15.6 Hz, H-10), 3.74 (3H, s, COCH₃), 4.71 (1H, d, J =4.71 Hz, H-1'); ¹³C-NMR : Table 1.

Compound (5) (Daphylloside) A white amorphous powder, anthrone / H₂SO₄ reaction positive, $[\alpha]_D^{32}$ +10.7 ° (c=2.0, MeOH); IR $\nu_{\max}$ (KBr) cm⁻¹: 3456, 1716, 1696, 1670, 1636; UV $\lambda_{\max}$ (MeOH) nm (log ϵ): 201.9 (3.7), 234.9 (4.0); ¹H-NMR (δ , CD₃OD): 5.06 (1H, d, J =9.0 Hz, H-1), 7.65 (1H, d, J =1.4 Hz, H-3), 3.03 (1H, ddd, J =7.4, 6.4, 1.4 Hz, H-5), 6.02 (1H, d, J =1.8 Hz, H-7), 2.64 (1H, dd, J =9.0, 7.4 Hz, H-9), 4.95, 4.80 (each, 1H, d, J =15.6 Hz, H-10), 2.09 (3H, s, COCH₃), 3.74 (3H, s, CH₃), 4.72 (1H, d, J =7.8 Hz, H-1'); ¹³C-NMR : Table 1.

Compound (6) (6-O-Acetylscandoside) A white amorphous powder, anthrone / H₂SO₄ reaction positive, $[\alpha]_D^{32}$ -15.6 ° (c=2.0, MeOH); IR $\nu_{\max}$ (KBr) cm⁻¹: 3392, 1738, 1702, 1664; UV $\lambda_{\max}$ (MeOH) nm (log ϵ): 202.4 (3.8), 234.5 (3.8); ¹H-NMR (δ , CD₃OD): 5.56 (1H, d, J =6.0 Hz, H-1), 7.30 (1H, d, J =2.2 Hz, H-3), 3.35 (1H, dd, J =6.1, 2.2 Hz, H-5), 5.95 (1H, d, J =1.4 Hz, H-7), 3.68 (1H, m, H-9), 4.77, 4.66 (each, 1H, d, J =14.3 Hz, H-10), 2.08 (3H, s, COCH₃), 4.68 (1H, d, J =7.9 Hz, H-1'); ¹³C-NMR : Table 1

Compound (7) (10-Hydroxyloganin) A white amorphous powder, anthrone / H₂SO₄ reaction positive, $[\alpha]_D^{27}$ -93.8 ° (c=1.4, MeOH); IR $\nu_{\max}$ (KBr) cm⁻¹: 3432, 1702, 1634; UV $\lambda_{\max}$ (MeOH) nm (log ϵ): 193.6 (3.5), 193.7 (3.5), 234.5 (4.0); ¹H-NMR (δ , CD₃OD): 5.15 (1H, d, J =6.1 Hz, H-1), 7.45 (1H, d, J =1.4 Hz,

H-3), 3.64 (1H, m, H-5), 2.27 (1H, ddd, $J=13.4, 9.6, 4.4$ Hz, H-6), 1.54 (1H, ddd, $J=13.4, 9.6, 4.4$ Hz, H-6), 4.29 (1H, t, $J=2.1$ Hz, H-7), 3.69 (3H, s, COCH₃), 4.56 (1H, d, $J=7.9$ Hz, H-1'); ¹³C-NMR : Table 1.

Compound (**8**) (Phytol) A colorless oil, $[\alpha]_D^{27} +0.2$ ($c=10$, CHCl₃), IR_{v_{max}} (CCl₄) cm⁻¹ : 3628, 2956, 2928, 2868, 1668, 1464, 1380; ¹H-NMR (δ , C₅D₅N) : 5.74 (1H, tq, $J=5.3, 1.3$ Hz, H-2), 4.45 (2H, d, $J=6.5$ Hz, H-1), 2.02 (1H, t, $J=7.1$ Hz, H-4), 1.69 (3H, br s, H-20), 1.51 (1H, sep, $J=6.8$ Hz, H-15), 0.90, 0.88 (each 3H, d, $J=6.8$ Hz, H-16 and H-17), 0.88, 0.87 (each 3H, s, H-18 and H-19); ¹³C-NMR (δ , C₅H₅N) : 136.9 (C-3), 126.3 (C-2), 59.0 (C-1), 40.3 (C-4), 39.7(C-14), 37.8 (C-8), 37.8 (C-10), 37.7 (C-12), 37.1 (C-6), 33.2 (C-7), 33.1 (C-11), 28.3 (C-15), 25.6 (C-5), 25.2 (C-13), 24.9 (C-9), 22.9 (C-17), 22.8 (C-16), 20.0 (C-18), 20.0 (C-19), 16.3 (C-20).

ACKNOWLEDGEMENTS

This paper is the contribution for special issue on the celebration of the 70th birthday of Professor James P. Kutney, Professor of the University of British Columbia.

REFERENCES

1. Part 5 in the series "Studies on Jamu and the medicinal resources in Indonesia". Part 4: *Chem. Pharm. Bull.*, 2001, **49**, 551.
2. E. J. H. Corner and K. Watanabe, "AN ILLUSTRATE OF TOROPICAL PLANTS", Vol. IV, Hirokawa Publishing Co., Tokyo, 1983, p. 677; C. A. Baker and R. C. B. V. D. Brink Jr., "FLORA OF JAVA", Vol. II, Wolters-Noordhoff N. V., the Netherlands, 1968, p. 354; S. Kasahara and S. Henmi, "MEDICAL FARB INDEX IN INDONESIA", P. T. Eisai Indonesia, Indonesia, 1986, p. 270.
3. I. H. Burkill, " A Dictionary of the Economic Products of the Malay Peninsula, " Publication Governments of Malaysia and Singapore, Kuala Lumpur, 1966, p. 356.
4. T. Endo and H. Taguchi, *Chem. Pharm. Bull.*, 1973, **21**, 2684 ; H. Inoue, Y. Takeda, S. Saito, H. Nishimura, and R. Sakuragi (nee Amano), *Yakugaku Zasshi*, 1974, **94**, 577.

5. A. Bianco, M. Guiso, C. Iavarone, P. Passacantilli, and C. Trogolo, *Gazz. Chim. Ital.*, 1978, **108**, 13;
A. Frank and H. Rimpler, *Phytochemistry*, 1987, **26**, 3015.
6. H. Inouye, S. Inouye, N. Shimokawa, and M. Okigawa, *Tetrahedron Lett.*, 1968, 683; R. K.
Chaudhuri, F. U. Afifi-Yazar, and O. Sticher, *Helv. Chim. Acta*, 1979, **62**, 1603; H. Inouye, Y. Takeda,
and H. Nishimura, *Phytochemistry*, 1974, **13**, 2219.
7. H. Inouye, Y. Takeda, S. Saitou, H. Nishimura, and R. Sakuragi, *Yakugaku Zasshi*, 1974, **94**, 577.
8. H. Inouye, M. Ueda, M. Hirabayashi, and N. Shimokawa, *Yakugaku Zasshi*, 1966, **86**, 943.
9. S. Uesato, M. Ueda, H. Inoue, H. Kuwajima, M. Yatsuzuka, and K. Takaishi, *Phytochemistry*, 1984,
23, 2535; S. Damtoft, S. Frederiksen, and R. Jensen, *Phytochemistry*, 1994, **35**, 1259.
10. G. D. Brown, *Phytochemistry*, 1994, **36**, 1553; T. Aoki, K. Takagi, T. Hirata, and T. Suga,
Phytochemistry, 1982, **21**, 1361; R. A. Goodman, E. Oldfield, and A. Allerhand, *J. Am. Chem. Soc.*,
1973, **95**, 7553.