


A NEW ROUTE TO (–)-APHANORPHINE USING A DIOXABICYCLO[3.2.1]OCTANE CHIRAL BUILDING BLOCK[†]

Adel S. ElAzab, Takahiko Taniguchi, and Kunio Ogasawara*

Pharmaceutical Institute, Tohoku University, Aobayama, Sendai 980-8578, Japan

Abstract — A new stereocontrolled route to (–)-aphanorphine, isolated from the fresh-water blue-green algae *Aphanizomenon flos-aquae* has been developed by using a chiral building block having a dioxabicyclo[3.2.1]octane framework and originally designed for the construction of the aldohexose molecules.


Recently, we designed a chiral building block (**1**) having a dioxabicyclo[3.2.1]octane framework for the diastereodivergent synthesis of all of the eight possible diastereomers of aldohexoses.¹ Owing to its biased structure, (–)-**1** allowed convex-face selective modification of its enone functionality leading to all of the eight L-aldohexose diastereomers in a diastereocontrolled manner.² We report here another use³ of the sugar building block for an alternative synthesis of (–)-aphanorphine^{4~6} (**2**), the only naturally occurring norbenzomorphan structure isolated from the fresh-water blue-green algae *Aphanizomenon flos-aquae* on the basis of the same methodology employed in the sugar synthesis (**Scheme 1**).



Scheme 1

Enantiopure [(–)-**1**] was reduced with diisobutylaluminum hydride (DIBAL) in the presence of copper (I) iodide in THF containing hexamethylphosphoric triamide (HMPA) to give the ketone (**3**), $[\alpha]_D^{30} +37.4^\circ$ (c1.1, CHCl_3). On reaction with iodomethane in the presence of lithium hexamethyldisilazide (Li-HMDS) in THF containing HMPA,⁷ **3** afforded a 2.5:1 mixture of the monomethylated ketone (**4**), which was refluxed with 4-methoxyphenylhydrazine hydrochloride in 90% aqueous pyridine^{3c,8} to give rise to the

[†]Dedicated to Prof. James P. Kutney, University of British Columbia, on the occasion of his 70th birthday.




Scheme 2

Reagents and conditions: (i) DIBAL, CuI, HMPA, THF, -78°C (89%). (ii) Li-HMDA, MeI, HMPA, THF, $-78\sim-30^\circ\text{C}$ (72%). (iii) 4-MeOC₆H₄NHNH₂-HCl, 90% aq. pyridine, reflux (92%). (iv) H₃PO₂, NaNO₂, Et₂O, AcOH, $0^\circ\text{C} \sim \text{rt.}$ (81%). (v) NaBH₄, MeOH, 0°C (100%). (vi) CS₂, MeI, NaH, THF (95%). (vii) Bu₃SnH, AIBN (cat.), benzene, reflux (95%). (viii) Raney Ni (W-2), EtOH, reflux (ix) MsCl, Et₃N, CH₂Cl₂, 0°C . (x) LiI, THF, reflux (94% from **12**).

carbinol amine (**8**), $[\alpha]_D^{29} -88.7^\circ$ (c 1.0, CHCl₃), as a single product. Apparently, the reaction proceeded through a convex-face selective 3,3-sigmatropic pathway via a diaza-1,5-diene intermediate **5** to give a transient imine (**6**) from which **8** was resulted *via* the amino-ketone (**7**) with hydrolytic loss of an ammonia under the conditions. This indicated that convex-face selectivity prevails even in the intramolecular reaction owing to the inherent steric nature of a bicyclo[3.2.1]octane system. To eliminate the extra aromatic amine functionality, **8** was exposed to sodium nitrite and hypophosphorus acid^{9,10} to initiate diazotization under reductive conditions. The expected reaction did occur to furnish the ketone (**9**), $[\alpha]_D^{28} -31.9^\circ$ (c 1.0, CHCl₃), in good yield. Since a single-step reduction of the carbonyl functionality of **9** under Wolff-Kishner conditions failed, **9** was first reduced with sodium borohydride to give the *endo*-alcohol (**10**), $[\alpha]_D^{29} -7.2^\circ$ (c 0.4, CHCl₃), which then was converted into the xanthate (**11**), $[\alpha]_D^{28} +32.1^\circ$ (c 1.0,

CHCl₃). On reflux with tributylstannane in benzene in the presence of a catalytic amount of azobisisobutyronitrile (AIBN),¹¹ (**11**) afforded the deoxygenation product (**12**), $[\alpha]_D^{30} +36.2^\circ$ (*c* 1.9, CHCl₃), excellently. To cleave the dioxolane functionality, which was found to be sturdy under standard acid-hydrolysis conditions, **12** was transformed into the iodide (**15**) through a sequential debenzylation with Raney nickel (W-2),¹² mesylation of the primary alcohol (**13**), $[\alpha]_D^{29} +66.3^\circ$ (*c* 1.1, CHCl₃), obtained, followed by substitution of the resulting mesylate **14** with lithium iodide to yield the iodide **15**, $[\alpha]_D^{30} +2.8^\circ$ (*c* 1.1, CHCl₃). Overall yield of **15** from the starting building block (**1**) was 41% in ten steps (**Scheme 2**).

The iodide (**15**) was then refluxed with zinc in ethanol containing acetic acid^{1,2} to initiate reductive cleavage to give the hemiacetal (**16**), as an epimeric mixture, which afforded the δ -lactone (**17**), $[\alpha]_D^{30} +35.7^\circ$ (*c* 1.2, CHCl₃), on oxidation with tetrapropylammonium perruthenate¹³ (TPAP) in the presence of 4-methylmorpholine *N*-oxide. Cleavage of the vinyl functionality of **17** was next carried out in a two-step sequence involving catalytic dihydroxylation and periodate cleavage¹⁴ to give the aldehyde (**18**). When **18** was refluxed with zinc in acetic acid intending to initiate cleavage of the α -oxygen bond of the formyl functionality,¹⁵ the reaction proceeded more easily than we anticipated. Gratifyingly, the product generated



Scheme 3

Reagents and conditions: (i) Zn, AcOH-EtOH(1:10), reflux. (ii) TPAP(cat.), NMO, THF (96% from **15**). (iii) OsO₄(cat.), NMO, 50% aq. THF, then NaIO₄, 50% aq. THF (83%). (iv) Zn, AcOH, reflux (69%). (v) (PhO)₂P(O)N₃, Et₃N, benzene, sealed tube, 140°C, 1 h, then MeOH, 4 h (92%).

was found to be the dihydronaphthalene (**20**), a more advanced intermediate, though not the initially expected formyl-acid (**19**). Upon heating with diphenylphosphoryl azide¹⁶ (DPPA) in benzene containing triethylamine in a sealed tube at 140 °C for one hour and for four hours at the same temperature after addition of methanol in the same sealed tube, **20** afforded the methyl carbamate¹⁷ (**22**), $[\alpha]_D^{28} +6.6^\circ$ (*c* 1.1, CHCl₃) {lit.,¹⁷ $[\alpha]_D^{30} +6.85^\circ$ (*c* 0.9, CHCl₃)}, through a formation of the isocyanate intermediate (**21**). Since we have previously developed a five-step transformation¹⁷ of [(+)-**22**] into (–)-aphanorphine (**2**), the present acquisition of (+)-**22** from the sugar building block [(–)-**1**] constitutes an alternative synthesis of the natural products in a formal sense. Overall yield of (+)-**22** from the iodide **15** was 51% in five isolated steps and, thus, 21% from the block [(–)-**1**] in 15 steps (**Scheme 3**).

In summary, we have demonstrated an alternative utilization of the chiral building block originally developed for the construction of the aldohexose molecules for a concise synthesis of (–)-aphanorphine, the only naturally occurring alkaloid known to have the norbenzomorphan framework.

ACKNOWLEDGEMENTS

We are grateful for an Egyptian Associate Channel System Program Scholarship (to A. S. E.).

REFERENCES AND NOTES

- (a) M. Takeuchi, T. Taniguchi, and K. Ogasawara, *Synthesis*, 1999, 341. (b) T. Taniguchi, M. Takeuchi, K. Kadota, A. S. ElAzab, and K. Ogasawara, *Synthesis*, 1999, 1325.
- (a) M. Takeuchi, T. Taniguchi, and K. Ogasawara, *Chirality*, 2000, **12**, 338. (b) M. Takeuchi, T. Taniguchi, and K. Ogasawara, *Tetrahedron Lett.*, 2000, **41**, 2609.
- Utilization of a dioxabicyclo[3.2.1]octane chiral building block for natural product synthesis, see: (a) the C28-C34 segment of FK-506: M. Takeuchi, T. Taniguchi, and K. Ogasawara, *Tetrahedron: Asymmetry*, 2000, **11**, 1601. (b) (–)-shikimic acid: M. Takeuchi, T. Taniguchi, and K. Ogasawara, *Synthesis*, 2000, 1375. (c) (–)-physostigmine and (–)-physovenine: A. S. ElAzab, T. Taniguchi, and K. Ogasawara, *Org. Lett.*, 2000, **2**, 2757. (d) (+)-febrifugine: M. Taniguchi and K. Ogasawara, *Org. Lett.*, 2000, **2**, 3193. (e) Pseudomonic acid key segment: M. Taniguchi and K. Ogasawara, *Tetrahedron Lett.*, in press.
- Isolation and relative structure: N. Gulavita, S. Hori, Y. Shimizu, P. Laszlo, and J. Clardy, *Tetrahedron Lett.*, 1988, **29**, 4381.
- Absolute configuration: S. Takano, K. Inomata, and K. Ogasawara, *J. Chem. Soc., Chem. Commun.*, 1990, 290.
- Enantiocontrolled synthesis: (a) S. Takano, K. Inomata, T. Sato, and K. Ogasawara, *J. Chem. Soc., Chem. Commun.*, 1990, 1591. (b) T. Honda, A. Yamamoto, Y. Cui, and M. Tsubuki, *J. Chem. Soc., Perkin Trans. 1*, 1992, 531. (c) A. I. Meyers, W. Schmidt, and B. Santiago, *Heterocycles*, 1995, **40**, 525. (d) A. N. Hulme, S. S. Henry, and A. I. Meyers, *J. Org. Chem.*, 1995, **60**, 1265. (e) A. Fadel and P. Arzel, *Tetrahedron: Asymmetry*, 1995, **6**, 893. (f) K. O. Hallinan and T. Honda, *Tetrahedron*, 1995, **51**, 12211. (g) S. Shiotani, H. Okada, K. Nakamata, T. Yamamoto, and F.

- Sekino, *Heterocycles*, 1996, **43**, 1031. (h) M. Node, H. Imazato, R. Kurosaki, Y. Kawano, T. Inoue, K. Nishide, and K. Fuji, *Heterocycles*, 1996, **42**, 811. (i) A. Fadel and P. Arzel, *Tetrahedron: Asymmetry*, 1997, **8**, 371. (j) K. Tanaka, Takahiko Taniguchi, and Kunio Ogasawara, *Tetrahedron Lett.*, 2001, **42**, 1049.
- 7 S. Takano, K. Inomata, and K. Ogasawara, *J. Chem. Soc., Chem. Commun.*, 1992, 169.
- 8 W. M. Welch, *Synthesis*, 1977, 645.
- 9 J. Sepiol, *Tetrahedron*, 1986, **42**, 609.
- 10 S. Takano, M. Moriya, and K. Ogasawara, *Tetrahedron Lett.*, 1992, **33**, 329.
- 11 D. H. R. Barton and S. W. McCombie, *J. Chem. Soc., Perkin Trans. 1*, 1975, 1574.
- 12 Y. Oikawa, T. Tanaka, K. Horita, and O. Yonemitsu, *Tetrahedron Lett.*, 1984, **25**, 5397.
- 13 S. V. Ley, J. Norman, W. P. Griffith, and S. P. Marsden, *Synthesis*, 1994, 639.
- 14 R. Pappo, D. S. Allen, Jr., R. V. Lemieux, and W. S. Johnson, *J. Org. Chem.*, 1956, **21**, 478.
- 15 S. Takano, S. Satoh, and K. Ogasawara, *J. Chem. Soc., Chem. Commun.*, 1988, 59.
- 16 T. Shioiri, K. Ninomiya, and S. Yamada, *J. Am. Chem. Soc.*, 1972, **94**, 6203.
- 17 M. Shimizu, T. Kamikubo, and K. Ogasawara, *Heterocycles*, 1997, **46**, 21.