

REACTIONS OF 1-METHOXY-3-(2-NITROVINYL)INDOLE WITH NUCLEOPHILES: AN INTERESTING SOLVENT EFFECT AND A NOVEL PREPARATION OF 3-SUBSTITUTED 1-METHOXYINDOLES¹

Koji Yamada, Fumio Yamada, and Masanori Somei*

Faculty of Pharmaceutical Sciences, Kanazawa University,
13-1 Takara-machi, Kanazawa 920-0934, Japan

Abstract — Nucleophiles react with 1-methoxy-3-(2-nitrovinyl)indole (**3**) at the 2-position regioselectively in a dipolar aprotic solvent (DMF or HMPA), while in THF they undergo Michael addition to the β -carbon of nitrovinyl side chain. Depending on bases employed, the resultant Michael addition products (**6d**, **10**, and **11**) are found to undergo interesting cyclizations to give novel 3-substituted 1-methoxyindoles (**7**, **9**, and **12**).

We found that 1-methoxyindoles (**1a–f**, Scheme 1) having an electron withdrawing group at the 3-position undergo nucleophilic substitution reaction² with various nucleophiles in DMF, culminating in a novel method for preparing 2-substituted indoles (**2a–f**), which are hardly available by the conventional electrophilic substitution reaction³ of indoles. In order to determine the scope of the method, we then selected 1-methoxy-3-(2-nitrovinyl)indole (**3**) as a substrate.

The compound (**3**), readily available from commercially available 2,3-dihydroindole in three steps in 49% overall yield,^{4a,b} reacted with NaOMe and NaOPr in DMF at 80–92°C to provide 2-methoxy- (**4a**) and 2-propoxy-3-(2-nitrovinyl)indole^{4a} (**4b**) in 85 and 55% yields, respectively, as expected. In sharp contrast, employing THF as a solvent instead of DMF, the same reactions at 0°C produced 1-methoxy-3-(1-methoxy-2-nitroethyl)indole (**6a**, Michael addition product) and **6b** in 90 and 92% yields, respectively. In these reactions, minor products were formed, but the absences of **4a** and **4b** among them were confirmed by taking ¹H-NMR spectrum for every fraction of products.

Solvents are known to play an important role in governing the reaction paths and products in some cases.⁵ Our above example suggested that dipolar aprotic solvent (DMF or HMPA) directs nucleophilic attack to the 2-position of **3** with concomitant liberation of the 1-methoxy group, while THF leads them to Michael addition at the β -carbon of nitrovinyl side chain.

In order to confirm the assumption, we further compared the reaction of **3** with a few additional nucleophiles. As a result, the reaction with sodium allyl oxide in DMF at 86°C provided **4c** and **5a** in 65 and 6% yields, respectively. However, the same reaction in THF at 0°C produced **6c** in 58% yield without formation of 2-substituted indoles (**4c** and **5a**).

Similarly, in the reaction with potassium 1,1-dimethylallyl oxide, **3** provided rearranged product (**5b**) via **4d** in 39% yield by the reaction at 88°C in HMPA. On the other hand, in THF at 0°C, **3** afforded **6d** in 81% yield. Interestingly, when the latter reaction was carried out in THF at reflux, a 55% yield of novel

Scheme 1






Figure 1

ORTEP Drawing of **8c** ($R = 0.069$)






Table 1



Entry	Base (mol eq.)	Reaction Conditions		Yield (%)		of Recovery
		Temp. (°C)	Time (min)	9	10	
1	KOtBu	rt	10	35	0	11
2	60% NaH	0	10	0	75	0

Scheme 2



cyclized product (**7**) was formed through the expected **6d**.

In order to determine its structure unequivocally by X-Ray single crystallographic analysis, an attempt was made to convert **7** to a crystalline compound. The reactions of *p*-bromo- and *p*-chlorobenzoyl chlorides with **7** afforded **8a** and **8b** in 63 and 82% yields, respectively, but they were not suitable crystals. *p*-Nitrobenzoyl chloride was finally found to give a 78% yield of **8c** as feasible prisms. Figure 1 shows ORTEP drawing of **8c**, and consequently the structure of **7** is proved as well.


Taking the formation of **7** into consideration, we planned to produce novel 3-substituted 1-methoxyindoles in one pot by causing Michael addition to take place first, and then let the resultant product react intramolecularly by selecting an appropriate base. Employing KO^tBu as a base, the reaction of 1,1-dimethylpropargylalcohol with **3** in THF at room temperature provided a novel cyclized product (**9**) in 35% yield as shown in Table 1 (entry 1), while the same reaction using NaH in THF at 0°C provided Michael addition product (**10**) in 75% yield (entry 2). Although the structure of **9** was determined based on its spectral data, further confirmation was obtained by the fact that **9** was produced in 60% yield from **10** by treatment with KO^tBu at 0°C for 30 min.

Sodium 3-butyne-1-oxide also reacted with **3** in THF to give **11** in 95% yield (Scheme 2). By the treatment of **11** with *n*-BuLi in THF at -16°C, a novel cyclization occurred to give 7-(1-methoxyindol-3-yl)-4,5-dihydro-7*H*-pyrano[3,4-*c*]isoxazole (**12**) in 48% yield together with unreacted **11** (11%). Extension of this reaction to aromatic and aliphatic nitrovinyl compounds is in progress.

A possible reaction mechanism for the formations of **7** and **12** is shown in Scheme 3. Base abstracts active methylene proton of the Michael product (**13**), giving nitronate intermediate (**14**). In the case that the R substituent is 1,1-dimethylallyl group, cycloaddition⁶ of nitronate to alkene side chain occurs to give **15**. Subsequent elimination of the alkoxy part⁷ produces **7** via **16**. On the other hand, if the R substituent of **14** is 1,1-dimethylpropargyl group, cycloaddition of nitronate to alkyne side chain gives **17**. The following *n*-BuLi abstraction of hydrogen at the carbon attached to nitrogen with concomitant liberation of lithium hydroxide results in the formation of stable aromatic isoxazole (**12**).

In summary, we have disclosed that **3** has two reaction sites for nucleophilic substitution, and the choice

Scheme 3



of the solvent governs the position to which the nucleophiles add. In addition, Michael addition products (**6d**, **10**, and **11**) undergo interesting cyclizations depending on the bases giving novel 3-substituted 1-methoxyindoles (**7**, **9**, and **12**). Attempts to determine the scope and limitations of these reactions are in progress.

REFERENCES AND NOTES

1. a) This report is Part 114 of a series entitled "The Chemistry of Indoles" and partly reported at the 120th Annual Meeting of Pharmaceutical Society of Japan, Book of Abstracts 2, Gifu, March 2000, p. 32. b) Part 113: K. Yamada, F. Yamada, T. Shiraishi, S. Tomioka, and M. Somei, *Heterocycles*, **58** (2002), in press. c) All new compounds gave satisfactory spectral and elemental analysis or high-resolution MS data for crystals or oils, respectively. **4a**) mp 209–212°C (decomp); **4b**) mp 191–198°C (decomp); **4c**) mp 124–129°C (decomp); **6a**) mp 101–102°C; **6b**) yellow viscous oil; **6c**) yellow viscous oil; **6d**) mp 85–86°C; **7**) yellow viscous oil; **8a**) mp 218–219°C (decomp); **8b**) mp 217–222°C (decomp); **8c**) mp 176.5–177.0°C; **9**) yellow viscous oil; **10**) mp 91.0–91.5°C; **11**) yellow viscous oil; **12**) yellow viscous oil.
2. Review: M. Somei, *Heterocycles*, 1999, **50**, 1157; M. Somei, N. Oshikiri, M. Hasegawa, and F. Yamada, *ibid.*, 1999, **51**, 1237; M. Somei, M. Nakajou, T. Teramoto, and F. Yamada, *ibid.*, 1999, **51**, 1949; M. Hasegawa, K. Yamada, Y. Nagahama, and M. Somei, *ibid.*, 1999, **51**, 2815; T. Kurauchi, Y. Nagahama, M. Hasegawa, K. Yamada, and M. Somei, *ibid.*, 2000, **53**, 1017; F. Yamada, A. Goto, and M. Somei, *ibid.*, 2000, **53**, 1255; M. Somei, Y. Fukui, M. Hasegawa, N. Oshikiri, and T. Hayashi, *ibid.*, 2000, **53**, 1725; M. Somei, F. Yamada, T. Kurauchi, Y. Nagahama, M. Hasegawa, K. Yamada, S. Teranishi, H. Sato, and C. Kaneko, *Chem. Pharm. Bull.*, 2001, **49**, 87; M. Somei, F. Yamada, T. Hayashi, A. Goto, and Y. Saga, *Heterocycles*, 2001, **55**, 457; K. Yamada, T. Kawasaki, Y. Fujita, and M. Somei, *ibid.*, 2001, **55**, 1151; T. Hayashi, W. Peng, Y. Nakai, K. Yamada, and M. Somei, *ibid.*, 2002, **56**, 421.
3. R. J. Sundberg, "The Chemistry of Indoles," Academic Press, New York, 1970; R. J. Sundberg, "Indoles," Academic Press, New York, 1996.
4. a) M. Somei, F. Yamada, T. Izumi, and M. Nakajou, *Heterocycles*, 1997, **45**, 2327; b) M. Somei, A. Tanimoto, H. Orita, F. Yamada, and T. Ohta, *ibid.*, 2001, **54**, 425. See also reference 1b.
5. C. Reichardt, "Solvent Effects in Organic Chemistry," 3, Verlag Chemie GmbH, Weinheim, 1979.
6. Related cycloadditions of nitronates: S. E. Denmark and A. Thorarensen, *Chem. Rev.*, 1996, **96**, 137; G. Broggini and G. Zecchi, *Synthesis*, 1999, 905 and references cited therein.
7. A. Hassner, O. Friedman, and W. Dehaen, *Liebigs Ann.*, 1997, 587.