

RHODIUM-CATALYZED ASYMMETRIC 1,4-ADDITION OF 3-THIOPHENEBORONIC ACID TO α,β -UNSATURATED CARBONYL COMPOUNDS

Kazuhiro Yoshida and Tamio Hayashi*

Department of Chemistry, Graduate School of Science, Kyoto University, Sakyo,
Kyoto 606-8502, Japan

E-mail: thayashi@kuchem.kyoto-u.ac.jp

Abstract - Asymmetric 1,4-addition of 3-thiopheneboronic acid to α,β -unsaturated carbonyl compounds proceeded with high enantioselectivity in the presence of 3 mol% (Rh) of $[\text{Rh}(\text{OH})((S)\text{-binap})]_2$ in dioxane/ H_2O (10/1) at 40 °C to give the corresponding optically active β -(3-thienyl) carbonyl compounds of over 94% ee.

Asymmetric 1,4-addition of organometallic reagents to α,β -unsaturated carbonyl compounds is a useful process giving enantiomerically enriched β -substituted carbonyl compounds.¹ As one of the most promising methods for the asymmetric 1,4-addition, the addition of aryl- and alkenylboronic acids under catalysis by chiral phosphine-rhodium complexes has been rapidly developed.²⁻⁴ However, to our best knowledge, heteroaromatic groups have not been introduced enantioselectively by the rhodium-catalyzed asymmetric 1,4-addition,⁵ probably due to the instability of the corresponding boronic acids under the reaction conditions, the reactions typically being carried out with $\text{Rh}(\text{acac})((S)\text{-binap})$ as a catalyst in dioxane/ H_2O at 100 °C.² During our recent studies on the catalytic cycle of the rhodium-catalyzed asymmetric 1,4-addition,⁶ we found that $[\text{Rh}(\text{OH})((S)\text{-binap})]_2$ is much more catalytically active, which


† Dedicated to Professor Yuichi Kanaoka on the occasion of his 75th birthday.

catalyzes the asymmetric 1,4-addition at a lower temperature with higher enantioselectivity. Here we wish to report that 3-thienyl group can be introduced on to the β -position of a variety of α,β -unsaturated carbonyl compounds with high enantioselectivity by use of the new catalyst $[\text{Rh}(\text{OH})((S)\text{-binap})]_2$ for the reaction of 3-thiopheneboronic acid.

RESULTS AND DISCUSSION

As a first set of experiment, several reaction conditions were examined for the addition of 3-thiopheneboronic acid (**2**) to 2-cyclohexenone (**1a**) (Scheme 1). Under our standard conditions used previously for the reaction of α,β -unsaturated ketones with phenylboronic acid,² that is, 3 mol % of the rhodium catalyst generated from $\text{Rh}(\text{acac})(\text{C}_2\text{H}_4)_2$ with (*S*)-binap⁷ in dioxane/ H_2O (10/1) at 100 °C for 5 h, the reaction of 2-cyclohexenone (**1a**) with 3-thiopheneboronic acid (**2**) (3 equiv to **1a**) gave only a poor yield (30%) of the 1,4-addition product, 3-thiophen-3-ylcyclohexanone (**3a**), whose enantiomeric purity is 97% (entry 1 in Table 1). Considering that the reaction of enone (**1a**) with phenylboronic acid gave a high yield (>90%) of the phenylation product under the same reaction conditions,² the low yield is ascribed to the use of 3-thiopheneboronic acid (**2**). In the reactions at this high temperature (100 °C), it has been sometimes observed^{2,6} that the hydrolysis of arylboronic acids ($\text{ArB}(\text{OH})_2$) giving arenes (ArH) is the main side reaction which is responsible for the low yields of the 1,4-addition products. The reaction carried out at 40 °C did not improve the yield of **3a** though the enantioselectivity was higher (99% ee) (entry 2). A higher yield of the 1,4-addition product (**3a**) was obtained by use of $[\text{Rh}(\text{OH})((S)\text{-binap})]_2$ as a catalyst. Thus, the reaction of **1a** with thiopheneboronic acid (**2**) (5 equiv.) in THF/ H_2O (10/1) in the presence of 3 mol % of $[\text{Rh}(\text{OH})((S)\text{-binap})]_2$ at 40 °C for 24 h gave 78% yield of (*S*)-3-thiophen-3-ylcyclohexanone (**3a**) with 99% ee (entry 3).

Scheme 1



Under similar reaction conditions, cyclic α,β -unsaturated ketones, 2-cyclopentenone (**1b**) and 2-cycloheptenone (**1c**), gave the corresponding thienylation products in 78% and 81% yields, respectively

Table 1. Asymmetric 1,4-Addition of 3-Thiopheneboronic Acid (**2**) to α,β -Unsaturated Carbonyl Compounds (**1**) Catalyzed by $[\text{Rh}(\text{OH})((S)\text{-binap})]_2$ ^a


entry	enone 1	reaction conditions	product 3 yield (%) ^d	% ee of 3 ^b (config)	$[\alpha]_D^{20c}$ of 3
1 ^e	1a	100 °C 5 h	3a 30	97 (<i>S</i>)	
2 ^e	1a	40 °C 24 h	3a 40	99 (<i>S</i>)	
3 ^f	1a	40 °C 24 h	3a 78	99 (<i>S</i>)	
4	1a	40 °C 24 h	3a 67	99 (<i>S</i>)	+11.1°
5	1b	40 °C 24 h	3b 78	97 (<i>S</i>)	-63.1°
6	1c	40 °C 24 h	3c 81	99 (<i>S</i>)	-34.7°
7	1d	40 °C 24 h	3d 79	99 (<i>S</i>)	+19.8°
8	1e	40 °C 24 h	3e 75	99 (<i>S</i>)	-25.8°
9	1f	40 °C 24 h	3f 82	94 (<i>R</i>)	-17.4°

^a The rhodium-catalyzed 1,4-addition was carried out with substrate (**1**) (0.20 mmol) and 3-thiopheneboronic acid (**2**) (0.60 mmol) in 0.55 mL of dioxane/H₂O (10/1) in the presence of 3 mol % (Rh) of $[\text{Rh}(\text{OH})((S)\text{-binap})]_2$ unless otherwise noted. ^b Determined by HPLC analysis of ketone (**3**) with chiral stationary phase columns (Chiralcel OD-H (**3a**, **3d**, **3e**, **3f**) and AD (**3b**, **3c**)). The configurations were assigned by consideration of the stereochemical reaction pathway (ref. 2a). ^c (c 1.00, chloroform). ^d Isolated yield by silica gel chromatography. ^e Reaction with 3 mol % of a rhodium catalyst generated from $\text{Rh}(\text{acac})(\text{C}_2\text{H}_4)_2$ and (*S*)-binap. ^f With 1.00 mmol of **2** in THF/H₂O (10/1).

(entries 5 and 6). The asymmetric thienylation of α,β -unsaturated ester, 5,6-dihydro-2*H*-pyran-2-one (**1d**), was also successful giving 79% yield of **3d** (entry 7). The enantiomeric excesses of **3b**, **3c**, and **3d** are all very high, ranging between 97% and 99% ee. The configurations of the 1,4-addition products (**3a-d**) are assigned to be *S* by consideration of the stereochemical reaction pathway.^{2a} The asymmetric 1,4-addition of thiopheneboronic acid also proceeded with high enantioselectivity for linear α,β -unsaturated ketones (**1e**) and (**1f**) (entries 8 and 9).

In summary, asymmetric 1,4-addition of 3-thiopheneboronic acid (**2**) to α,β -unsaturated carbonyl compounds (**1**) proceeded with high enantioselectivity in high yields by use of $[\text{Rh}(\text{OH})((S)\text{-binap})]_2$ as a catalyst. The enantiomerically enriched products substituted with thienyl group at the stereogenic carbon center are expected to be converted into a variety of useful compounds by further transformation.⁸

Scheme 2



EXPERIMENTAL SECTION

General. All anaerobic and/or moisture sensitive manipulations were carried out with standard Schlenk technique under predried nitrogen. Tetrahydrofuran and 1,4-dioxane were distilled from benzophenone-ketyl under nitrogen prior to use. Rhodium complex, $[\text{Rh}(\text{OH})((S)\text{-binap})]_2$ was prepared according to the reported procedure.⁶ 2-Cyclohexenone (**1a**), 2-cyclopentenone (**1b**), and 2-cycloheptenone (**1c**) were distilled before use. 5,6-Dihydro-2H-pyran-2-one (**1d**), 5-methyl-3-hexen-2-one (**1e**), 3-nonen-2-one (**1f**), and 3-thiopheneboronic acid (**2**) were used as received without further purification. NMR spectra were recorded on a JEOL JNM LA-500 spectrometer (500 MHz for ¹H and 125 MHz for ¹³C). Chemical shifts are reported in δ ppm referenced to an internal SiMe₄ standard for ¹H NMR and chloroform-*d* (δ 77.05) for ¹³C NMR. Optical rotations were measured on a JASCO DIP-370 polarimeter.

Rhodium-Catalyzed Asymmetric 1,4-Addition of 3-Thiopheneboronic Acid (2) to α,β -Unsaturated Carbonyl Compounds (1): General Procedure. To a solution of $[\text{Rh}(\text{OH})((S)\text{-binap})]_2$ (4.4 mg, 3.0 μmol) and 3-thiopheneboronic acid (**2**) (76.8 mg, 0.600 mmol) in dioxane (0.5 mL) was added α,β -unsaturated carbonyl compound (**1**) (0.200 mmol) and water (0.05 mL) at rt under nitrogen. The flask was immersed in a bath maintained at 40 °C for 24 h. The resulting mixture was extracted with ethyl acetate and the extract was washed with saturated sodium bicarbonate, before it was

dried over MgSO_4 and concentrated under reduced pressure. The residue was purified by PTLC on silica gel to give the desired 1,4-addition product (**3**). The results are summarized in Table 1. Spectroscopic data of the products are shown below:

3-Thiophen-3-ylcyclohexanone (3a): colorless liquid, purified by PTLC (hexane/AcOEt = 4/1); ^1H NMR (CDCl_3): δ 1.73-1.86 (m, 2H), 2.06-2.12 (m, 1H), 2.13-2.20 (m, 1H), 2.32-2.39 (m, 1H), 2.41-2.47 (m, 1H), 2.50 (ddd, $J = 14.1, 11.8, 1.3$ Hz, 1H), 2.68 (ddt, $J = 14.1, 4.1, 1.9$ Hz, 1H), 3.16 (tt, $J = 11.0, 4.1$ Hz, 1H), 6.98-6.99 (m, 2H), 7.29 (dd, $J = 4.7, 3.3$ Hz, 1H). ^{13}C NMR (CDCl_3): δ 25.04, 32.40, 39.85, 41.26, 48.52, 119.51, 125.95, 126.44, 145.41, 210.78. $[\alpha]^{20}_{\text{D}} + 11.1^\circ$ (c 1.00, CHCl_3). Anal. Calcd for $\text{C}_{10}\text{H}_{12}\text{OS}$: C, 66.63; H, 6.71. Found: C, 66.62; H, 6.69.

3-Thiophen-3-ylcyclopentanone (3b): white solid, purified by PTLC (hexane/AcOEt = 4/1); ^1H NMR (CDCl_3): δ 1.94-2.04 (m, 1H), 2.24-2.34 (m, 2H), 2.39-2.48 (m, 2H), 2.66 (dd, $J = 18.3, 7.7$ Hz, 1H), 3.49 (tt, $J = 10.2, 7.1$ Hz, 1H), 7.00-7.02 (m, 2H), 7.31 (t, $J = 3.9$ Hz, 1H). ^{13}C NMR (CDCl_3): δ 30.79, 37.71, 38.45, 45.68, 119.59, 126.18, 126.59, 144.28, 218.15. $[\alpha]^{20}_{\text{D}} - 63.1^\circ$ (c 1.00, CHCl_3). mp 33-35 $^\circ\text{C}$. Anal. Calcd for $\text{C}_9\text{H}_{10}\text{OS}$: C, 65.02; H, 6.06. Found: C, 65.14; H, 5.98.

3-Thiophen-3-ylcycloheptanone (3c): colorless liquid, purified by PTLC (hexane/AcOEt = 4/1); ^1H NMR (CDCl_3): δ 1.47-1.55 (m, 1H), 1.65-1.78 (m, 2H), 1.92-2.05 (m, 2H), 2.12-2.18 (m, 1H), 2.54-2.57 (m, 2H), 2.71 (ddd, $J = 14.6, 2.7, 1.8$ Hz, 1H), 2.87 (dd, $J = 14.5, 11.5$ Hz, 1H), 3.07 (tt, $J = 11.2, 2.6$ Hz, 1H), 6.95-6.97 (m, 2H), 7.26 (dd, $J = 4.9, 3.1$ Hz, 1H). ^{13}C NMR (CDCl_3): δ 24.09, 28.82, 37.78, 38.50, 43.95, 50.85, 118.92, 125.80, 126.53, 147.05, 213.27. $[\alpha]^{20}_{\text{D}} - 34.7^\circ$ (c 1.00, CHCl_3). Anal. Calcd for $\text{C}_{11}\text{H}_{14}\text{OS}$: C, 68.00; H, 7.26. Found: C, 68.08; H, 7.34.

4-Thiophen-3-yltetrahydropyran-2-one (3d): white solid, purified by PTLC (hexane/AcOEt = 1/2); ^1H NMR (CDCl_3): δ 2.00 (dtd, $J = 14.9, 10.0, 4.9$ Hz, 1H), 2.23 (dq, $J = 14.2, 4.3, 1.6$ Hz, 1H), 2.64 (dd, $J = 17.6, 10.2$ Hz, 1H), 2.96 (ddd, $J = 17.6, 6.0, 1.6$ Hz, 1H), 3.36 (tt, $J = 10.3, 5.4$ Hz, 1H), 4.38 (ddd, $J = 11.5, 10.0, 3.8$ Hz, 1H), 4.47 (dt, $J = 11.5, 4.8$ Hz, 1H), 6.98 (dd, $J = 5.1, 1.4$ Hz, 1H), 7.03-7.04 (m, 1H), 7.34 (dd, $J = 5.0, 2.8$ Hz, 1H). ^{13}C NMR (CDCl_3): δ 30.08, 32.92, 37.15, 68.39, 119.99, 126.00, 126.75, 143.65, 170.34. $[\alpha]^{20}_{\text{D}} + 19.8^\circ$ (c 1.00, CHCl_3). mp 49-51 $^\circ\text{C}$. Anal. Calcd for $\text{C}_9\text{H}_{10}\text{O}_2\text{S}$: C, 59.32; H, 5.53. Found: C, 59.54; H, 5.61.

5-Methyl-4-thiophene-3-ylhexan-2-one (3e): colorless liquid, purified by PTLC (hexane/AcOEt = 8/1); ^1H NMR (CDCl_3): δ 0.78 (d, $J = 6.7$ Hz, 3H), 0.90 (d, $J = 6.7$ Hz, 3H), 1.82 (octet, $J = 6.7$ Hz, 1H), 2.00 (s, 3H), 2.74 (d, $J = 7.3$ Hz, 2H), 3.12 (q, $J = 7.4$ Hz, 1H), 6.90-6.92 (m, 2H), 7.23 (dd, $J = 4.5, 3.4$ Hz, 1H). ^{13}C NMR (CDCl_3): δ 19.87, 20.65, 30.47, 32.94, 43.23, 47.53, 120.84, 125.08, 127.44, 143.71, 208.30. $[\alpha]^{20}_{\text{D}} - 25.8^\circ$ (c

1.00, CHCl₃). Anal. Calcd for C₁₁H₁₆OS: C, 67.30; H, 8.22. Found: C, 67.31; H, 8.03. **4-Thiophen-3-ylnonan-2-one (3f)**: colorless liquid, purified by PTLC (hexane/AcOEt = 8/1); ¹H NMR (CDCl₃): δ 0.84 (t, *J* = 7.1 Hz, 3H), 1.13-1.27 (m, 6H), 1.47-1.62 (m, 2H), 2.02 (s, 3H), 2.66 (dd, *J* = 16.1, 6.9 Hz, 1H), 2.69 (dd, *J* = 16.1, 7.5 Hz, 1H), 3.27 (dddd, *J* = 8.2, 7.5, 6.9, 6.4 Hz, 1H), 6.93-6.94 (m, 2H), 7.24 (dd, *J* = 4.8, 3.0 Hz, 1H). ¹³C NMR (CDCl₃): δ 14.01, 22.50, 26.97, 30.56, 31.70, 36.25, 36.59, 50.66, 120.15, 125.53, 126.55, 145.42, 207.98. [α]_D²⁰ -17.4° (*c* 1.00, CHCl₃). Anal. Calcd for C₁₃H₂₀OS: C, 69.59; H, 8.98. Found: C, 69.45; H, 8.78.

ACKNOWLEDGMENT

This work was supported in part by a Grant-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology, Japan. K. Y. thanks the Japan Society for the Promotion of Science for the award of a fellowship for graduate students.

REFERENCES AND NOTES

- For reviews on asymmetric 1,4-addition of organometallic reagents: (a) N. Krause and A. Hoffmann-Röder, *Synthesis*, 2001, 171; (b) M. P. Sibi and S. Manyem, *Tetrahedron*, 2000, **56**, 8033; (c) K. Tomioka and Y. Nagaoka, 'Comprehensive Asymmetric Catalysis,' Vol. 3, ed. by E. N. Jacobsen, A. Pfaltz, and H. Yamamoto, Springer, Berlin, 1999, Chapter 31.1.
- (a) Y. Takaya, M. Ogasawara, T. Hayashi, M. Sakai, and N. Miyaoura, *J. Am. Chem. Soc.*, 1998, **120**, 5579; (b) Y. Takaya, M. Ogasawara, and T. Hayashi, *Tetrahedron Lett.*, 1998, **39**, 8479; (c) Y. Takaya, T. Senda, H. Kurushima, M. Ogasawara, and T. Hayashi, *Tetrahedron: Asymmetry*, 1999, **10**, 4047; (d) Y. Takaya, M. Ogasawara, and T. Hayashi, *Tetrahedron Lett.*, 1999, **40**, 6957; (e) T. Hayashi, T. Senda, Y. Takaya, and M. Ogasawara, *J. Am. Chem. Soc.*, 1999, **121**, 11591; (f) Y. Takaya, M. Ogasawara, and T. Hayashi, *Chirality*, 2000, **12**, 469; (g) T. Hayashi, T. Senda, and M. Ogasawara, *J. Am. Chem. Soc.*, 2000, **122**, 10716; (h) T. Hayashi, *Synlett*, 2001, 879; (i) T. Senda, M. Ogasawara, and T. Hayashi, *J. Org. Chem.*, 2001, **66**, 6852; (j) T. Hayashi, M. Takahashi, Y. Takaya, and M. Ogasawara, *Org. Synth.*, 2002, **79**, 84.
- (a) S. Sakuma, M. Sakai, R. Itooka, and N. Miyaoura, *J. Org. Chem.*, 2000, **65**, 5951; (b) S. Sakuma and N. Miyaoura, *J. Org. Chem.*, 2001, **66**, 8944.
- (a) M. Kuriyama and K. Tomioka, *Tetrahedron Lett.*, 2001, **42**, 921; (b) M. T. Reetz, D. Moulin, and

- A. Gosberg, *Org. Lett.*, 2001, **3**, 4083; (c) M. Lautens, C. Dockendorff, K. Fagnou, and A. Malicki, *Org. Lett.*, 2002, **4**, 1311.
5. Non-asymmetric version of the 1,4-addition has been reported by use of potassium thienyltrifluoroborate: R. A. Batey, A. N. Thadani, and D. V. Smil, *Org. Lett.*, 1999, **1**, 1683.
6. T. Hayashi, M. Takahashi, Y. Takaya, and M. Ogasawara, *J. Am. Chem. Soc.*, 2002, **124**, 5052.
7. (a) H. Takaya, K. Mashima, K. Koyano, M. Yagi, H. Kumobayashi, T. Taketomi, S. Akutagawa, and R. Noyori, *J. Org. Chem.*, 1986, **51**, 629; (b) H. Takaya, S. Akutagawa, and R. Noyori, *Org. Synth.*, 1989, **67**, 20; (c) R. Noyori and H. Takaya, *Acc. Chem. Res.*, 1990, **23**, 345; (d) D. Cai, J. F. Payack, D. R. Bender, D. L. Hughes, T. R. Verhoeven, and P. J. Reider, *J. Org. Chem.*, 1994, **59**, 7180.
8. For reviews: (a) A. I. Meyers, 'Heterocycles in Organic Synthesis,' John Wiley & Sons, New York, 1974; (b) A. R. Katritzky and C. W. Rees, 'Comprehensive Heterocyclic Chemistry,' Pergamon, Amsterdam, 1984, Vol. 4; (c) A. R. Katritzky, C. W. Rees, and E. F. V. Scriven, 'Comprehensive Heterocyclic Chemistry II,' Pergamon, Amsterdam, 1996, Vol. 2.