

SYNTHESIS OF FURAZANE CONJUGATED NEW HETEROCYCLES

Renaud Beaudegnies* and Sebastian Wendeborn

Syngenta Crop Protection AG, Optimization Chemistry

CH-4002 Basel, Switzerland

E-mail: renaud.beaudegnies@syngenta.com

Abstract- An improved route for the synthesis of 3-(4-aminofurazan-3-yl)-1,2,4-oxadiazole heterocycles using micro-wave irradiation is reported. The preparation of novel 4-(thiazol-4-yl)furazan-3-ylamine, 4-(pyrimidin-4-yl)furazan-3-ylamine and, 4-(pyrazolidin-3-yl)furazan-3-ylamine heterocycles are described.

Furazanes (1,2,5-oxadiazoles) have been reported to exhibit a wide spectrum of biological properties.¹ In particular, agrochemical applications of furazanes and their derivatives as herbicides, plant-growth regulators and pesticides have been described.² Our biological screening revealed that 3-(4-aminofurazan-3-yl)-4H-[1,2,4]oxadiazol-5-one (**5**) exhibits herbicidal activity on broad leaf (*dicotyledones*) weeds. This prompted us to investigate the synthesis of novel 4-aminofurazane heterocycles.

In the first part of this communication we report the preparation of **5** as well as a new and efficient method for the high speed synthesis of [3-(4-aminofurazan-3-yl)-4H-[1,2,4]oxadiazol-5-yl]amine derivatives (**4**).

In the second part, we report the preparation of novel 4-aminofurazane rings conjugated to thiazole, pyrimidine and pyrazolidine heterocycles.

Few examples have been reported in the literature describing the synthesis of 3-(4-aminofurazan-3-yl)-[1,2,4]oxadiazole heterocycles.³ 4-(5-Trichloromethyl[1,2,4]oxadiazol-3-yl)-furazan-3-ylamine (**3**) appeared to be the intermediate of choice for the preparation of both 3-(4-aminofurazan-3-yl)-4H-[1,2,4]oxadiazol-5-one (**5**) and new [3-(4-aminofurazan-3-yl)-4H-[1,2,4]oxadiazol-5-yl]amine derivatives (**4a-p**) (Scheme 1). Indeed, it has been reported that the trichloromethyl group at the C-5 position of a 1,2,4-oxadiazole ring can be readily substituted by nucleophiles such as guanidines to yield 5-amino-1,2,4-oxadiazole derivatives.⁴


The sequence followed for the preparation of the amide oxime (**3**) is similar to the one previously published by Andrianov *et al.*^{3d} However, their one-pot procedure for the preparation of **3** directly from **1**⁵ could not be reproduced in our hands (lit.,^{3d} 58%). Our new stepwise pathway involves intermediate (**2**) and provides **3** more efficiently from **1** (77% yield two steps).

Treatment with *p*-toluidine was required for the selective removal of the trichloroacetyl group of the intermediate (**2**) which provides **3** in 81% yield. The ammonolysis of **3** in the presence of NH₄OH gave **5** (87% yield).

In order to access a number of structurally diverse analogs of **5**, a small library was prepared by treating **3** with a series of both primary and secondary amines. When the reaction was performed in EtOH/THF at 70°C (method A), the reaction was slow (2-3 days) and low yielding, but nonetheless provided a number of pure [3-(4-aminofurazan-3-yl)-4*H*-[1,2,4]oxadiazol-5-yl]amines after crystallisation.






















The efficiency of the process could be drastically improved when **3** and 1.2 eq. of amine were adsorbed together on neutral alumina and irradiated in the microwave oven at 400 W (method B). Under these conditions, the reactions were much faster (< 20 min) and substantially cleaner.

The compounds prepared by these two methods are summarized in Table 1.




Scheme 1. a) NaNO₂ (2 eq.), HCl (2N), rt, 5 h; b) NH₂OH.HCl (2.2 eq.), NaOH (pH=10), rt; c) reflux, 3 h; d) (CCl₃CO)₂O, 120°C, 5 h; e) *p*-toluidine (2 eq.), C₂H₅OH, 80°C, 19 h; f) (CCl₃CO)₂O, 60-80°C (lit.,^{2d} 58%); g) **method A** : amine (3-5 eq.), C₂H₅OH, 70°C, 2-3 d.; h) **method B** : amine (1.2 eq.) on neutral alumina, microwave (400W), 2x10 min.; i) 25% NH₄OH.

Table 1. Transformation of **3** into **4**


entry	amine	method	product	yield	entry	amine	method	product	yield
1		A	4a	<5%	12		B	4f	90%
2		B	4a	quant.	13		A	4g	44%
3		A	4b	15%	14		A	4h	81%
4		B	4b	quant.	15		A	4i	90%
5		A	4c	65%	16		A	4j	88%
6		B	4c	95%	17		A	4k	79%
7		A	4d	30%	18		A	4l	43%
8		B	4d	88%	19		A	4m	57%
9		A	4e	33%	20		A	4n	70%
10		B	4e	86%	21		B	4o	96%
11		A	4f	39%	22		A	4p	36%

We also report the preparation of novel 4-aminofurazanes bearing 5- or 6- membered ring heterocycles.⁶ Their syntheses use 1-(4-aminofurazan-3-yl)ethanone (**10**)^{7,8} as a key intermediate (Schemes 3 and 4). It has been reported that such 3-acylated furazane rings can be prepared through the reduction of the corresponding carbonyl-substituted furoxans.⁷ We report here a new synthesis of **10** as depicted in Scheme 2 in which the key step is the mild and regioselective cyclisation of **8** induced by the treatment with NaOAc in dioxane at 60°C, to give **9** in high yield.



Scheme 2. a) isopentyl nitrite, (C₂H₅)₂O, HCl; b) KCN (1.8 eq.), (C₂H₅)₂O, H₂O, 45°C ; c) H₂NOH.HCl, NaOH, H₂O, rt ; d) Ac₂O, 0°C-rt ; e) NaOAc (4 eq.), dioxane, 60°C; f) 20% HCl, H₂O, reflux, steam distillation.

1-(4-Aminofurazan-3-yl)ethanone (**10**) was brominated according to reported conditions.^{8b} Thiazole derivatives (**12** and **13**) were prepared in 26 to 99% yields through the Hantzsch reaction of bromide (**11**)^{8b} and the corresponding thioamides or thioureas (Scheme 3 and Table 2).⁹



Scheme 3. a) Br₂ (1.05 eq.), AcOH (100%), 55°C; b) **method A**: 1.1 eq. nucleophile, benzene, reflux, 2-3 h, Dean-Stark; c) **method B**: 1.1 eq. nucleophile, C₂H₅OH, reflux, 2-3 h.

Table 2. Synthesis of 4-(thiazol-4-yl)furazan-3-ylamine derivatives (**12**) and (**13**)


entry	reagent	product	yield	entry	reagent	product	yield
1		12a	60%	9		13e	70%
2		12b	26%	10		13f	79%
3		12c	34%	11		13g	91%
4		12d	93%	12		13h	98%
5		13a	87%	13		13i	94%
6		13b	75%	14		13j	85%
7		13c	99%	15		13k	90%
8		13d	88%	16		13l	96%

4-(Pyrimidin-4-yl)furazan-3-ylamine and 4-(pyrazolidin-3-yl)furazan-3-ylamine heterocycles (**18-23** and **24** to **31**) respectively are available in three steps from **10** via the reaction of **15** with ambident nucleophiles such as amidines, guanidines or hydrazines (Scheme 4 and Table 3).

As shown in the Scheme 4, triphenylphosphorane (**14**) did not react with aldehydes and only starting materials were recovered. In contrast, piperidine catalyzed aldol condensations between **10** and various aldehydes proceeded in satisfactory to excellent yields (Scheme 4).

The resulting α,β -unsaturated ketones (**15**, **16**, and **17**) were allowed to react with amidines and guanidines in the presence of oxygen to give 4-(pyrimidin-4-yl)furazan-3-ylamine derivatives (**24** to **31**) in low to moderate yields. Compounds (**15** and **17**) underwent reaction with hydrazine to give oxygen sensitive

4-(pyrazolidin-3-yl)furazan-3-ylamine heterocycles which were directly acylated with various acid chlorides in good yields, thus allowing the introduction of additional points of diversity.



Scheme 4. a) Br₂ (1.05 eq.), AcOH (100%), 55°C; b) R¹CHO (2.0 eq.), piperidine (0.1 eq.), C₂H₅OH /THF: 5/1, 50°C; c) PPh₃ (1.0 eq.), toluene, 100°C; d) NaOH (2N), CH₃OH, 0°C; e) R¹CHO (1.2 eq.), piperidine (0.1 eq.), 60°C; f) amidine or guanidine (1.1 eq.), *N,N*-dimethylacetamide, 60°C, O₂ (gas); g) PhNHNH₂ (2.0 eq.), THF/ C₂H₅OH: 1/20, rt; h) H₂NNH₂.H₂O (10 eq.), THF, rt; i) R³COCl (1.2 eq.), pyridine (3.0 eq.), CH₂Cl₂, rt.

Table 3. Synthesis of compounds (18 to 31)

entry	educt	R1	R2	product	yield	entry	educt	R1	R3	product	yield
1*	15	cyclopropyl	methyl	18	29%	8	15	cyclopropyl		25	92%
2	15	cyclopropyl	<i>p</i> -methoxyphenyl	19	14%	9	17	2-furyl		26	99%
3	15	cyclopropyl	phenyl	20	26%	10	25	cyclopropyl	methyl	27	74%
4	15	cyclopropyl	aniline	21	16%	11	26	2-furyl	2,4-dichlorophenyl	28	92%
5	15	cyclopropyl	benzylamine	22	25%	12	26	2-furyl	(4-chlorophenoxy)methyl	29	76%
6	16	<i>p</i> -methoxyphenyl	<i>m</i> -methylaniline	23	56%	13	26	2-furyl	2-furyl	30	73%
7	17	2-furyl		24	26%	14	26	2-furyl	cyclopropyl	31	79%

(*) Performed in the presence of 1.1 eq. of DBU.

CONCLUSION

We have developed a new and efficient method for the synthesis of [3-(4-aminofurazan-3-yl)-4H-[1,2,4]-oxadiazol-5-yl]amine derivatives. We have also enlarged the scope of the 4-aminofurazane chemistry to novel 4-(thiazol-4-yl)furazan-3-ylamine, 4-(pyrimidin-4-yl)furazan-3-ylamine and, 4-(pyrazolidin-3-yl)-furazan-3-ylamine heterocycles.

REFERENCES

1. R.M. Paton, *Comprehensive Heterocyclic Chemistry II*, ed. by A.R. Katritzky, C.W. Rees, and E.F.V. Scriven, Pergamon, Oxford, 1996, **4**, 229 and references cited therein.
2. (a) J.C. Bottaro, M.A. Petrie, P.E. Penwell, and D.C. Bomberger, *PCT Int. Appl.*, 2003, WO 0306371A1 (*Chem. Abstr.*, 2003, **138**, 126950); (b) H. Cerecetto, E. Dias, R. Di Maio, M.Gonzalez, S. Pacce, P. Saenz, G. Seoane, L. Suescun, A. Mombru, G. Fernandez, M. Lema, and J. Villalba, *J. Agric. Food Chem.*, 2000, **48**, 2995; (c) *Ger. Offen.*, 1985, DE 3326509A1 (*Chem. Abstr.*, 1985, **102**, 220879h); (d) R.W.W. Baldwin, *U.S. Patent*, 1970, US 3501285 (*Chem. Abstr.*, 1970, **72**, 111477s); (e) *French Patent*, 1969, FR 1576140 (*Chem. Abstr.*, 1970, **72**, 110178h); (f) W.E. Buting and C. Ainsworth, *U.S. Patent*, 1966, US 3279988 (*Chem. Abstr.*, 1967, **66**, 28776u).
3. (a) S.D. Shaposhnikov, N.V. Korobov, A.V. Sergievskii, S.V. Pirogov, S.F. Mel'nikova, and I.V. Tselinskii, *Russian J. Org. Chem.*, 2002, **38**, 1351; (b) S. Bailey, P.V. Fish, K. James, and G.A. Whitlock, *PCT Int. Appl.*, 2002, WO 0279200A1 (*Chem. Abstr.*, 2002, **137**, 294960); (c) V.N. Yarovenko, V.Z. Shirinyan, I.V. Zavarzin, and M.M. Krayushkin, *Izv. Akad. Nauk, Ser. Khim.* 1994, 118; (d) V.G. Andrianov, E.N. Rozhkov, and A.V. Ereemeev, *Khim. Geterotsikl. Soedin.*, 1994, 534; (e) V.G. Andrianov, V.G. Semenikhina, and A.V. Ereemeev, *Khim. Geterotsikl. Soedin.*, 1994, 539; f) O.A. Rakitin, V.A. Ogurtsov, E.A. Khaibullina, T.I. Godovikova, and L.I. Khmel'nitskii, *Khim. Geterotsikl. Soedin.*, 1993, 1283; (g) V.G. Andrianov, V.G. Semenikhina, and A.V. Ereemeev, *Khim. Geterotsikl. Soedin.*, 1992, 687; (h) V.N. Yarovenko, V.Z. Shirinyan, I.V. Zavarzin, and M.M. Krayushkin, *Izv. Akad. Nauk, Ser. Khim.* 1991, 2166; i) V.N. Yarovenko, V.K. Taralashvili, I.V. Zavarzin, and M.M. Krayushkin, *Tetrahedron*, 1990, **46**, 3941.
4. (a) I. Kumita, A. Niwa, C. Sano, and M. Takagi, *Jpn. Kokai Tokkyo Koho*, 1993, JP 05117255A2 (*Chem. Abstr.*, 1993, **119** : P271167h); (b) J.L. LaMattina and C.J. Mularski, *J. Org. Chem.*, 1984, **49**, 4800.
5. (a) V.G. Andrianov, A.V. Ereemeev, and Y.B. Sheremet, *Khim. Geterotsikl. Soedin.*, 1988, 856; (b) T. Ichikawa, T. Kato, and T. Takenishi, *J. Heterocycl. Chem.*, 1965, **2**, 253.
6. For reviews on Furazanes fused to 5-, 6-, and 7-membered rings see: (a) A.B. Sheremetev, *J. Heterocycl. Chem.*, 1995, **32**, 371; (b) A.B. Sheremetev, *Russian Chemical Reviews*, 1999, **68**, 137.
7. A.S. Kulikov, N.N. Makhova, T.I. Godovikova, S.P. Golova, and L.I. Khmel'nitskii, *Izv. Akad. Nauk, Ser. Khim.*, 1994, 679.
8. (a) T.S. Novikova, T. M. Mel'nikova, O.V. Kharitonova, V.O. Kulagina, N.S. Aleksandrova, A.B. Sheremetev, T.S. Pivina, L.I. Khmel'nitskii, and S.S. Novikov, *Mendeleev Commun.*, 1994, 138; (b) A.B. Sheremetev, A.S. Kulikov, and L.I. Khmel'nitskii, *Izv. Akad. Nauk, Ser. Khim.*, 1993, 744; (c) E. Borello, A. Zecchina, and E. Guglielminotti, *Gazz. Chim. Ital.*, 1965, **95**, 1465.
9. For similar reaction see : A.S. Kulikov and N.N. Makhova, *Izv. Akad. Nauk, Ser. Khim.*, 1998, 137.
10. Analytical and spectroscopic data are shown in the following. **1**: mp 190-192°C (H₂O), (lit.,^{5b} 190-191°C); ¹H NMR (300 MHz, CD₃OD) δ= 4.90 (br s); ¹³C NMR (75 MHz, CD₃OD) δ= 155.92, 145.80, 141.19; Anal. Calcd for C₃H₅N₅O₂ : C, 25.18; H, 3.52; N, 48.94. Found: C, 25.25; H, 3.48; N,

48.91; MS (electrospray): 142 (M-H⁺). **4m**: amorphous powder; ¹H NMR (300 MHz, DMSO d₆) δ= 9.00 (s, 1H), 6.35 (s, 2H), 3.50 (m, 2x2H), 3.28 (s, 3H); ¹³C NMR (75 MHz, DMSO d₆) δ= 171.61, 158.89, 155.31, 137.34, 69.94, 57.88, 42.85; MS (electrospray): 225 (M-H⁺), 261 (M+Cl⁻), 451 (2M-H⁺). **5**: mp 208-210°C (decomp), (isopropanol); ¹H NMR (300 MHz, DMSO d₆) δ= 7.30 (br s, 2H), 6.80 (br s, 1H); ¹³C NMR (75 MHz, DMSO d₆) δ= 172.46, 159.23, 155.29, 138.99; HRMS (EI): Calcd 169.0236 Found: 169.0233. **9**: mp 136-138°C (CH₃OH); ¹H NMR (300 MHz, CDCl₃) δ= 5.50 (br s, 2H), 2.51 (s, 3H), 2.30 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ= 166.87, 155.10, 154.38, 142.69, 19.28, 13.50. **10**: mp 94-95°C (CH₂Cl₂), (lit.,⁷ 96°C); ¹H NMR (300 MHz, CDCl₃) δ= 5.20 (s, 2H), 2.74 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ= 192.84, 155.17, 143.49, 28.16; MS (EI) 127 (M⁺). **11**: mp 162°C (CH₂Cl₂), (lit.,^{8b} 162-163°C); ¹H NMR (300 MHz, DMSO d₆) δ= 6.50 (br s, 2H), 4.87 (s, 2H); ¹³C NMR (75 MHz, DMSO d₆) δ= 184.72, 155.67, 141.91, 34.34; MS (EI) 207, 205 (M⁺). **13f**: amorphous powder; ¹H NMR (300 MHz, CD₃OD) δ= 7.20 (s, 1H), 4.80 (br s, 4H); ¹³C NMR (75 MHz, CD₃OD) δ= 171.70, 156.40, 142.38, 136.42, 109.07; MS (electrospray): 182 (M-H⁺), 228(M+HCOO⁻), 262, 264 (M+Br⁻); MS (electrospray): 184 (M+H⁺), 225 (M+CH₃CN+H⁺). **15**: mp 109°C (ethyl acetate); ¹H NMR (300 MHz, CDCl₃) δ= 7.10 (d, *J* = 15.6 Hz, 1H), 6.80 (dd, *J* = 15.6 and 11.3 Hz, 1H), 5.20 (br s, 1H), 1.75 (m, 1H), 1.15 (m, 2H), 0.85 (m, 2H); ¹³C NMR (75 MHz, CDCl₃) δ= 182.11, 158.81, 156.05, 143.32, 123.09, 15.98, 10.25; MS (electrospray): 180 (M+H⁺). **22**: mp 122-124°C (Hexane); ¹H NMR (300 MHz, CDCl₃) δ= 7.40-7.20 (m, 6H), 5.55 (br s, 1H), 5.20 (br s, 2H), 4.58 (d, *J* = 6.2 Hz, 2H), 1.95 (m, 1H), 1.15 (m, 2H), 1.05 (m, 2H); MS (electrospray): 309 (M+H⁺); MS (electrospray): 307 (M-H⁺), 343 (M+Cl⁻), 367 (M+CH₃COO⁻), 651 (2M+Cl⁻). **27**: amorphous powder; ¹H NMR (300 MHz, CDCl₃) δ= 7.32 (s, 1H), 6.38 (m, 1H), 6.34 (m, 1H), 5.71 (m, 1H), 5.20 (br s, 2H), 3.15 (m, 2H), 2.39 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ= 168.65, 154.27, 150.83, 142.52, 142.43, 140.24, 110.70, 108.53, 53.20, 38.32, 22.01.