

FACIAL SELECTIVITY OF THE SHARPLESS BROMINE CATALYZED AZIRIDINATION¹


Aaron C. Schmitt,² Catherine M. Smith, Eric A. Voight,² and George A. O'Doherty*

Department of Chemistry, West Virginia University, Morgantown, WV 26506, USA. George.ODoherty@mail.wvu.edu

Abstract – A bromine catalyzed aziridination reaction has been applied to 7,7-dibromo-3-norcarene, which resulted in a highly stereoselective route to *cis*-4,4-dibromo-8-(toluene-4-sulfonyl)-8-azatricyclo[5.1.0.0]octane in a 46% yield. In order to confirm the stereochemistry of the aziridination reaction *trans*-4,4-dibromo-8-(toluene-4-sulfonyl)-8-azatricyclo[5.1.0.0]octane and its diastereoisomer *cis*-4,4-dibromo-8-(toluene-4-sulfonyl)-8-azatricyclo[5.1.0.0]octane were prepared in a stereoselective manner from the corresponding *cis*- and *trans*-7,7-dibromo-3-norcarene epoxides (61 and 9.2% yields, respectively). These three routes offer stereochemical proof for the stereospecificity of the bromine-catalyzed aziridination of hindered alkenes.

INTRODUCTION


As part of a project directed toward the synthesis of a substituted cocaine ring system, we hoped to synthesize the 6-substituted tropane (**1**) from cycloheptane (**3**) (Scheme 1). In turn we envisioned preparing **3** from either *cis*- or *trans*-aziridine (**4a**) and (**4b**) by two ring opening reactions.^{3,4} In the course of our synthetic investigations of this route to the tropane ring system, we desired diastereoselective access to both the *cis*- and *trans*-aziridines (**4a**) and (**4b**). Thus we decided to investigate the various methods for the aziridination of cyclohexene (**5**). Herein, we describe two diastereoselective approaches for the selective synthesis of both *cis*- and *trans*-tosylaziridines (**4a**) and (**4b**). These results show the stereoselectivity of the bromine-catalyzed aziridination reaction and in turn lend credence to the mechanism proposed by Sharpless.^{5,6}



Scheme 1

RESULTS AND DISCUSSION


The starting 7,7-dibromo-3-norcarene (**5**) was prepared by the dibromocyclopropanation procedure of cyclohexa-1,4-diene reported by Winstein.⁷ Treatment of **5** with 1 equiv. chloramine-T and 10 mol % phenyltrimethylammonium perbromide⁸ provided a 3:1 mixture of oxidized products, the *N*-tosylaziridine (**4**) (35 %) and bromo sulfonamide (**7**) (11.3 %) (Scheme 2). The bromo sulfonamide (**7**) was cleanly converted into the aziridine (**4**) upon treatment with NaH under anhydrous conditions (98 %). The mixture of **4** and **7** could similarly be converted into **4** with NaH, which greatly simplified the chromatographic purification of **4**. While the ¹H NMR and ¹³C NMR spectra clearly indicated the formation of **4** as a single diastereoisomer, we were unable to determine whether the *cis* or *trans* stereoisomer was formed.⁹



Scheme 2


Sharpless has suggested that the perbromide-catalyzed aziridination of alkenes occurs through a bromonium ion (**8a**) or (**8b**) that is subsequently trapped by chloramine-T (forming **9a** or **9b**) (Scheme 3). The intermediate *N*-chlorosulfonamide (**9**) is then reduced by a halide ion and cyclized to form the corresponding tosylaziridine (**4**). In terms of the stereochemistry of aziridination, two mechanistic questions still remained unanswered. Are the two diastereotopic bromonium ions (**8a**) and (**8b**) rapidly inter-converting? Which of the two diastereotopic bromonium ions reacts faster with chloramine-T? Thus, at the outset it was not clear to us whether a facially biased alkene would react under these conditions to form the sterically more hindered *cis*-aziridine (**4a**) or sterically less hindered *trans*-aziridine (**4b**). Therefore, we decided to stereospecifically synthesize both the *cis*- and *trans*-*N*-

tosylaziridines (**4a**) and (**4b**) from stereo-defined starting materials and compare them to the material made from the Sharpless procedure.



Scheme 3


Following virtually identical conditions, the known *trans*-epoxide (**10**)¹⁰ and *cis*-epoxide (**11**)¹¹ were converted into diastereomeric azido-alcohols (**12**) and (**13**) (scheme 4).¹² Exposing the *trans*-epoxide (**10**) to excess NaN_3 (5 equiv.) and NH_4Cl (1 equiv.) in refluxing methanol for 3 days produced the azido-alcohol (**12**) in an 82 % yield. Not surprisingly the diastereomeric epoxide (**11**) reacted more quickly with NaN_3 requiring only 3 h to give the corresponding azide (**13**) in the same yield (82 %).



Scheme 4

Once again under similar conditions the diastereomeric azido-alcohols (**12**) and (**13**) were transformed to the corresponding *cis*- and *trans* *N*-tosylaziridines (**4a**) and (**4b**), respectively (Scheme 5).¹³ Exposure of **12** with triphenylphosphine in hot toluene (80 °C, 2 h) reductively cyclized **12** to an aziridine, which was protected *in situ* (TsCl /pyr, 0 °C) to form the *cis*-tosylaziridine (**4a**) (22 % yield). The low yield of **4a** is presumably due to the steric hindrance of the *endo*-azide in **12**. This is supported by the observation that exposure of the diastereomeric *exo*-azide (**13**) to the same PPh_3/TsCl conditions provided the corresponding *trans*-tosylaziridine (**4b**) in a much higher 84 % yield. With authentic samples of **4a** and **4b** in hand it was evident that the *N*-tosylaziridine prepared from the bromine-catalyzed aziridination of **5** gives exclusively the *cis*-aziridine (**4a**). The ^1H NMR and ^{13}C NMR of **4a** and **4b** were clearly distinguishable and the crude ^1H NMR spectra from the bromine-catalyzed aziridination showed no evidence of the *trans*-aziridine (**4b**).

In conclusion, two diastereoselective routes to the *cis*-*N*-tosylaziridines (**4a**) (46 and 9.2 % yields) and one diastereoselective route to the *trans*-*N*-tosylaziridines (**4b**) (61 % yield) have been developed. These three routes help establish the stereospecificity of the bromine-catalyzed aziridination of hindered alkenes. In particular, this heavily biased substrate demonstrates the high degree that the initially formed bromonium ion can be trapped. Further use of these *N*-tosylaziridines for the synthesis of the tropane ring system will be reported in due course.



Scheme 5

EXPERIMENTAL

General Methods: Unless otherwise stated, all reactions were carried out under an atmosphere of nitrogen using oven-dried glassware and standard syringe/septa techniques. Analytical TLC was performed using precoated glass-backed plates (Whatman K6F 60A, F₂₅₄) that were analyzed by fluorescence upon 254 nm irradiation or by staining with *p*-anisaldehyde, potassium permanganate, or phosphomolybdic acid stains. Liquid chromatography was performed using (flash chromatography) of the indicated solvent system on ICN reagent silica gel 60 (60-200 mesh). Ether and tetrahydrofuran were distilled from benzophenone and sodium metal. Dichloromethane and triethylamine were distilled from calcium hydride. Hexanes refers to the petroleum fraction bp 40-60 °C. Commercial reagents were used without purification unless otherwise noted. ¹H and ¹³C NMR spectra were recorded on Varian 300 and 500 MHz spectrometers. Chemical shifts are reported relative to CDCl₃ (δ 7.26 ppm) or internal tetramethylsilane (δ 0.00 ppm) for ¹H spectrum and CDCl₃ (δ 77.0 ppm) for ¹³C spectrum. Melting points are uncorrected. IR spectra were obtained on a Prospect MIDAC FT-IR spectrometer. HRMS spectrometric data was performed by the University of Minnesota Mass Spectrometry Laboratory. Combustion analyses were performed by M-H-W Laboratories, Phoenix, AZ.

***cis*-4,4-Dibromo-8-(toluene-4-sulfonyl)-8-azatricyclo[5.1.0.0^{3,5}]octane (4a) Method A:** To a solution of dibromide (**5**) (1.00 g, 3.96 mmol) dissolved in 20 mL of CH₃CN were added

phenyltrimethylammonium tribromide (PTAB) (0.142 g, 0.377 mmol) and Chloramine-T•3H₂O (1.23 g, 4.35 mmol) at rt. The reaction was stirred for 16 h, then diluted with CH₂Cl₂ (20 mL) and saturated NaHCO₃ solution (20 mL). The organic layer was separated, and the aqueous layer extracted with CH₂Cl₂ (2 x 20 mL). The combined organic layers were dried using Na₂SO₄ and the solvent was removed in vacuo. The crude material was first run through a plug of silica gel using 0 - 50 % ether/hexanes to remove starting material and excess tosylamine, and then a column was run using 5-20 % EtOAc/hexanes. Isolated were 0.204 g (20 %) of dibromide (**5**), 0.476 g (28 %) of aziridine as a white crystalline solid, and 0.182 g (9.1 %) of open tribromide (**7**) as an off-white crystalline solid. R_f 0.37 (20 % EtOAc/hexanes). mp 177 - 179 °C. ¹H NMR (500 MHz, CDCl₃) δ 7.77 (d, *J* = 8.5 Hz, 2H), 7.33 (d, *J* = 8.5 Hz, 2H), 2.96 (m, 2H), 2.63 (m, 2H), 2.44 (s, 3H), 1.79 (m, 2H), 1.12 (ddd, *J* = 17.0, 5.0, 4.0 Hz, 2H). ¹³C NMR (75 MHz, CDCl₃) δ 144.5, 135.1, 129.7, 127.7, 39.8, 37.0, 26.4, 21.7, 20.6. IR (neat): δ 3044, 2996, 2922, 1596, 1547, 1494, 1393, 1163 cm⁻¹. HRMS (CI) calcd for C₁₄H₁₅NO₂ Br₂S [M+H⁺] 419.9269, obsd 419.9279. Anal. Calcd for C₁₄H₁₅NO₂Br₂S: C, 39.93, H, 3.59. Found: C, 40.04, H, 3.77. Data for (+/-)-(1S*,3R*,4R*,6R*)-4-Methyl-N-(4,7,7-tribromobicyclo[4.1.0]hept-3-yl)-benzenesulfonamide (**7**). R_f 0.49 (20 % EtOAc/hexanes) ¹H NMR (300 MHz, CDCl₃) δ 7.76 (d, *J* = 8.1 Hz, 2H), 7.31 (d, *J* = 8.1 Hz, 2H), 4.76 (d, *J* = 5.1 Hz, 1H), 3.94 (ddd, *J* = 16.2, 9.9, 6.0 Hz, 1H), 3.15 (m, 1H), 2.85 (ddd, *J* = 15.6, 9.9, 6.0 Hz, 1H), 2.56 (ddd, *J* = 15.3, 6.0, 1.8 Hz, 1H), 2.43 (s, 3H), 1.92 (m, 2H), 1.42 (m, 2H). ¹³C NMR (75 MHz, CDCl₃) δ 143.9, 137.0, 129.8, 127.4, 54.1, 50.3, 36.8, 32.9, 29.0, 28.8, 26.9, 21.7. IR (neat): δ 3268, 3087, 2921, 1597, 1514, 1438, 1328, 1160, 1092 cm⁻¹. HRMS (CI) calcd for C₁₄H₁₆NO₂Br₃S [M+H⁺] 499.8530, obsd 499.8559, calcd for [M+NH₄⁺] 516.8796, obsd 516.8772.

cis-4,4-Dibromo-8-(toluene-4-sulfonyl)-8-azatricyclo[5.1.0.0^{3,5}]octane (4a) Method B: To a solution of tribromide (**7**) (143 mg, 0.286 mmol) in THF (2.8 mL) under nitrogen there was added 60 % NaH/mineral oil (24.3 mg, 0.632 mmol). After two hours the reaction was worked up by adding 10 % NaHCO₃ solution (10 mL) and was extracted with ethyl acetate (3 x 10 mL). The organic layers were combined and dried using Na₂SO₄, and the solvent was removed in vacuo. The crude product required no further purification, yielding 117 mg (98 %).

(+/-) -(1R*,3R*,4R*,6S*)-4-Azido-7,7-dibromobicyclo[4.1.0]heptan-3-ol (12): To a solution of epoxide (**10**) (302 mg, 1.12 mmol) in methanol (3 mL) there were added NaN₃ (366 mg, 5.63 mmol) and ammonium chloride (69.7 mg, 1.30 mmol). The mixture was then heated to 70 °C for 3 days. The reaction

was then diluted with ether (10 mL) and brine (10 mL). The layers were separated, and the aqueous layer was extracted with ether (3 x 10 mL). The organic layers were combined and dried over Na₂SO₄ and concentrated in vacuo. The remaining solid was purified by column chromatography (10 - 20 % EtOAc/hexanes) to yield 286 mg (82 %) of azido alcohol as a white solid, and 50 mg of epoxide starting material (17 %). R_f 0.49 (20 % EtOAc/hexanes) mp 57-59 °C. ¹H NMR (300 MHz, CDCl₃) δ 3.63 (ddd, *J* = 9.9, 9.9, 6.9 Hz, 1H), 3.23 (ddd, *J* = 12.5, 10.5, 5.4 Hz, 1H), 2.64 (ddd, *J* = 15.0, 9.6, 5.7 Hz, 1H), 2.45 (m, 1H), 2.25 (br s, 1H), 2.09-1.81 (m, 3H), 1.49 (ddd, *J* = 15.0, 12.6, 4.2, 1H). ¹³C NMR (75 MHz, CDCl₃) δ 69.4, 63.5, 37.5, 30.0, 29.3, 27.6, 26.9. HRMS calcd for C₇H₉N₃OBr₂ [M+NH₄⁺] 326.9456, obsd 326.9453. IR (neat): ν 3386, 3024, 2924, 2865, 2108, 1636, 1439, 1321, 1291, 1253, 1047 cm⁻¹. Anal. Calcd for C₇H₉N₃OBr₂: C, 27.04, H, 2.92. Found: C, 27.30, H, 3.12.

(+/-)-(1R*,3S*,4S*,6S*)-4-Azido-7,7-dibromobicyclo[4.1.0]heptan-3-ol (13): To a solution of epoxide (**11**) (302 mg, 1.12 mmol) in methanol (2.2 mL) were added NaN₃ (363 mg, 5.58 mmol) and ammonium chloride (59.8 mg, 1.12 mmol). The mixture was then heated to 70 °C and allowed to stir for 3 h. The reaction was then diluted with ether (5 mL) and brine (10 mL). The aqueous layer was extracted with ether (3 x 10 mL) and the combined organic layers were dried over Na₂SO₄ and concentrated in vacuo. The residue was purified by column chromatography (10 - 20 % EtOAc/hexanes) to yield 286 mg (82 %) of azide as a white solid. R_f 0.49 (20 % EtOAc/hexanes) mp 84 - 86 °C. ¹H NMR (300 MHz, CDCl₃) δ 3.37-3.52 (m, 2H), 2.57 (ddd, *J* = 15.6, 9.9, 5.4 Hz, 1H), 2.47 (dd, *J* = 13.8, 6.3 Hz, 1H), 2.21 (br s, 1H), 1.85-2.04 (m, 2H), 1.50 (ddd, *J* = 14.7, 11.4, 3.6 Hz, 2H). ¹³C NMR (75 MHz, CDCl₃) δ 70.0, 62.4, 37.3, 29.5, 28.0, 27.8, 27.1. IR (neat): ν 3387, 2924, 2865, 2108, 1290, 1047 cm⁻¹. HRMS calcd for C₇H₉N₃OBr₂ [M + NH₄⁺] 326.9456, obsd 326.9435. Anal. Calcd for C₇H₉N₃OBr₂: C, 27.04, H, 2.92. Found: C, 27.25, H, 3.17.

cis-4,4-Dibromo-8-(toluene-4-sulfonyl)-8-azatricyclo[5.1.0.0^{3,5}]octane (4a) Method C: To a solution of azide (**12**) (20.0 mg, 64.3 μ mol) in toluene (2 mL) was added triphenylphosphine (20.1 mg, 76.6 μ mol). The reaction was heated at 80 °C for 2 h until complete as judged by TLC analysis. The reaction was cooled to 0 °C and pyridine (10 μ L, 0.13 mmol) then tosyl chloride (18.6 mg, 97.5 μ mol) were added. The reaction was stirred overnight, then quenched with 10 % NaHCO₃ and extracted 3 times with CH₂Cl₂. The organic layers were dried using Na₂SO₄ and the solvent was removed in vacuo. The crude product was purified using column chromatography (10 - 20 % EtOAc/hexane). A white solid was isolated which had identical spectral data to compound (**4a**) yielding 6.0 mg of product (22 %).

***trans*-4,4-Dibromo-8-(toluene-4-sulfonyl)-8-azatricyclo[5.1.0.0^{3,5}]octane (4b):** To a solution of azide (**13**) (15.0 mg, 48.2 μ mol) in toluene (2 mL) was added triphenylphosphine (15.2 mg, 58.0 μ mol). The reaction was heated at 80 °C for 2 h until complete as judged by TLC analysis. The reaction was cooled to 0 °C and pyridine (10 μ L, 0.13 mmol) then tosyl chloride (18.4 mg, 96.5 μ mol) were added. The reaction was stirred overnight, then quenched with 10 % NaHCO₃ and extracted 3 times with CH₂Cl₂. The organic layers were dried using Na₂SO₄ and solvent was removed in vacuo. The crude product was purified using column chromatography (10 - 20 % EtOAc/hexane), yielding 17.0 mg of product as a white crystalline solid (84 %). *R*_f 0.50 (20 % EtOAc/hexanes). mp 165 -167 °C. ¹H NMR (500 MHz, CDCl₃) δ 7.79 (d, *J* = 8.5 Hz, 2H), 7.35 (d, *J* = 8.5 Hz, 2H), 2.88 (s, 2H), 2.46 (s, 3H), 2.33 (ddd, *J* = 16.5, 6.0, 2.5 Hz, 2H), 1.79 (dd, *J* = 16.0, 1.0 Hz, 2H), 1.70 (ddd, *J* = 5.0, 3.0, 1.0 Hz, 2H); ¹³C NMR (125 MHz, CDCl₃) δ 144.5, 135.3, 129.8, 127.7, 38.5, 36.9, 22.8, 21.7, 18.8. IR (film): ν 2920, 1596, 1428, 1323, 1159, 1091 cm⁻¹. HRMS calcd for C₁₄H₁₅NO₂Br₂S [M+H]⁺ 419.9269, obsd 419.9290.

ACKNOWLEDGMENTS

We thank the Arnold and Mabel Beckman Foundation, the American Cancer Society for an Institutional Research Grant (IRG-58-001-40-IRG-19), the American Chemical Society-Petroleum Research Fund (ACS-PRF#33953-G1) and the National Institute of General Medical Sciences (1R01 GM63150-01A1) for their generous support of our program. Funding by the National Science Foundation (NSF-EPSCoR award #0314742) for a Varian 600 MHz Inova Spectrometer and NMR facility is gratefully acknowledged.

REFERENCES AND NOTES

1. This paper is dedicated to Professor Leo A. Paquette on the occasion of his 70th birthday.
2. University of Minnesota Undergraduate Research Opportunity Program (UROP) Participant
3. For a review of dihalocyclopropane ring opening reactions, see: (a) M. G. Banwell and M. E. Reum, in 'Advances In Strain In Organic Chemistry,' Vol. 1, ed. by B. Halton, Jai Press, Inc. Greenwich, CT, 1991.

4. For Lewis acid catalyzed opening of aziridines see: B. A. B. Prasad, G. Sekar, and V. K. Singh, *Tetrahedron Lett.*, 2000, **41**, 4677 and references therein.
5. J. U. Jeong, B. Tao, I. Sagasser, H. Henniges, and K. B. Sharpless, *J. Am. Chem. Soc.*, 1998, **120**, 6844.
6. Upon completion of this work, similar stereoselectivity for this reaction were discovered in a cyclopentenol substrate, by O'Brien, see: D. Caine, P. O'Brien, and C. M. Rosser, *Org. Lett.*, 2002, **4**, 1923.
7. S. Winstein and J. Sonnenberg, *J. Am. Chem. Soc.*, 1961, **83**, 3235. W. von. E. Doering and A. K. Hoffmann, *J. Am. Chem. Soc.*, 1954, **76**, 6162.
8. We have found that freshly prepared tetrabutylammonium perbromide provided similarly good yields of **4** and **7**.
9. Due to the high degree of H-H *J*-coupling and the twist boat nature of **4** all nOe experiments were inconclusive in determining the relative stereochemistry between the cyclopropane and aziridine rings.
10. (a) J. Rudolph, K. L. Reddy, J. P. Chiang, and K. B. Sharpless, *J. Am. Chem. Soc.*, 1997, **119**, 6189.
(b) H. Adolfsson, A. Converso, and K. B. Sharpless, *Tetrahedron Lett.*, 1999, **40**, 3991.
11. L. A. Paquette, W. E. Fristad, C. A. Schuman, M. A. Beno, and C. G. Christoph, *J. Am. Chem. Soc.*, 1979, **101**, 4645.
12. K. Krajewski, Z. Ciunik, and I. Z. Siemion, *Tetrahedron: Asymmetry*, 1999, **10**, 4591.
13. J. Anman, T. Jarevang, and P. Somfai, *J. Org. Chem.*, 1996, **61**, 8148.