

FORMATION OF 3-PYRROLIN-2-ONES *VIA 5-ENDO-TRIG*


CYCLIZATION

Ruri Naitoh, Yayoi Nakamura, Emi Katano, Yohoko Nakamura, Emi Okada, and Morio Asaoka*

Department of Chemical and Biological Sciences, Faculty of Science, Japan Women's University, Mejirodai 2-8-1 Bunkyo, Tokyo 112-8681, Japan
E-mail: asaoka-m@fc.jwu.ac.jp


Abstract - Anionic cyclization of *N*-benzyl-3-phenylsulfanyl-2-propenamide derivatives gave the corresponding 3-pyrrolin-2-ones. Mechanistic investigation using deuterated starting materials revealed this cyclization proceeds *via* a 5-*endo-trig* process.

In the course of our studies on the chemistry of amide homoenolates,¹ we have examined the behavior of various *N*-benzyl-3-phenylsulfanyl-2-propenamide derivatives under basic conditions, and found that LDA treatment of (3-phenylsulfanyl)propenamide derivative (**1a**) at -20°C gave 3-pyrrolin-2-one (**2a**) in 80% yield. Having been interested in the mechanism, we looked into this reaction in detail.




Scheme 1

According to the Baldwin's rules, 5-*endo-dig* cyclizations are favorable processes whereas 5-*endo-trig* cyclizations are disfavored.² Therefore, we assumed that the above reaction should have proceeded *via* the 5-*endo-dig* process as shown below (Scheme 2).




Scheme 2

To confirm the assumption, deuterium was introduced into the starting material (**1a**). Thus, α -deuterated (**1b**) and β -deuterated compound (**1c**) were prepared and subjected to the cyclization. To our surprise, α -deuterium of **1b**, which should disappear during the *5-endo-dig* process, remained in the product (**2b**) and β -deuterium of **1c**, which should be untouched, disappeared (Scheme 3).



Scheme 3

On the bases of these results we concluded that this cyclization reaction proceeds *via* the *5-endo-trig* process as shown below (Scheme 4).



Scheme 4

Despite the disfavored nature of a *5-endo-trig* mode,² reports of examples continue to appear in the literature.³⁻⁵ However, to our best knowledge, most of the anionic *5-endo-trig* cyclizations reported so far include the nucleophilic addition of heteroatoms.^{3,4} Therefore, the above reaction is one of the rare examples which proceeds *via* the *5-endo-trig* process with an sp^3 carbon nucleophile.⁵

To assess the applicability of this type of cyclization, some 3-phenylsulfanyl-2-propenamide derivatives were subjected to the reaction. The results are summarized in Table 1.




Table 1. Cyclization of (3-phenylsulfanyl)propenamides.

entry	1	R	R'	time (min)	2	yield 2 (%)
1	(<i>S</i>)-1a	Me	Ph	60	(+)-2a	80
2	(<i>R</i>)-1a	Me	Ph	60	(-)-2a	78
3	1b (α -D)	Me	Ph	60	2b	76
4	1c (β -D)	Me	Ph	60	2a	72
5	1d	H	Ph	45	2d	46
6	1e	Me	<i>p</i> -Me-C ₆ H ₄	60	2e	64
7	1f	Me	1-naphthyl	45	2f	78
8	1g	Ph	Ph	60	2g	47
9	1h	Me	CO ₂ Et	30	2h	57

When **1d** (R=H) was used (entry 5), the yield was low and the product was contaminated with a small amount of impurities. Presumably side reaction occurs due to the high acidity of the γ -hydrogen of the product. In the other cases the corresponding pure products were isolated in good yields. Ester derivative (**1h**) also gave the corresponding cyclized product in good yield (entry 9).

It should be noteworthy that optically active starting materials [(*S*)-**1a** and (*R*)-**1a**] gave the corresponding optically active products [(+)-**2a**: $[\alpha]_D +278^\circ$ (*c* 2.8, CHCl₃) and (-)-**2a**: $[\alpha]_D -273^\circ$ (*c* 1.5, CHCl₃)] (entries 1 and 2).⁶ The optical purity was determined by HPLC on CHIRALCEL OJ-R and found to be >98% ee.

A typical procedure: To a cooled (-20°C) THF solution (20 mL) of LDA (1.3 mmol) was added a THF solution (5 mL) of **1a** (1 mmol), and then the mixture was stirred at that temperature for 60 min. Usual workup followed by purification by flash column chromatography (hexane:AcOEt=1:1) gave **2a** in 80% yield.

Further studies to explore the scope and limitations of the above reaction are currently in progress.

REFERENCES

1. a) T. Shinozuka, Y. Kikori, M. Asaoka, and H. Takei, *J. Chem. Soc., Perkin Trans. 1*, 1996, 119. b) T. Shinozuka, T. Utsumi, M. Inagaki, and M. Asaoka, *J. Chem. Soc., Perkin Trans. 1*, 1999, 2433. c) N. Sadayori, A. Kawatoh, R. Sakurazawa, T. Utsumi, M. Inagaki, and M. Asaoka, *Heterocycles*, 2002, **58**, 181.
2. a) J. E. Baldwin, *J. Chem. Soc., Chem. Commun.*, 1976, 734. b) J. E. Baldwin, J. Cutting, W. Dupont, L. Kruse, L. Silberman, and R. C. Thomas, *J. Chem. Soc., Chem. Commun.*, 1976, 736. c) J. E. Baldwin, R. C. Thomas, L. Kruse, and L. Silberman, *J. Org. Chem.*, 1977, **42**, 3846.
3. For some examples of pyrrolidine formation *via* anionic 5-*endo* cyclization, see: a) J. Ichikawa, M. Fujiwara, Y. Wada, T. Okauchi, and T. Minami, *Chem. Commun.*, 2000, 1887. b) J. Ichikawa, Y. Wada, T. Okauchi, and T. Minami, *Chem. Commun.*, 1997, 1537. c) D. Craig, P. S. Jones, and G. J. Rowlands, *Synlett*, 1997, 1423. d) D. W. Knight, A. L. Redfern, and J. Gilmore, *Tetrahedron Lett.*, 1998, **39**, 8909. e) M. B. Berry, D. Craig, P. S. Jones, and G. J. Rowlands, *Chem. Commun.*, 1997, 2141. f) S.-S. P. Chou and T.-M. Yuan, *Synthesis*, 1991, 171. g) A. D. Jones and D. W. Knight, *Chem. Commun.*, 1996, 915. h) A. Padwa and B. H. Norman, *J. Org. Chem.*, 1990, **55**, 4801.
4. For some examples of tetrahydrofuran formation *via* 5-*endo* cyclization, see: a) D. Craig, J. N. Ikin, N. Mathews, and A. M. Smith, *Tetrahedron Lett.*, 1995, **36**, 7531. b) S. P. Bew, J. M. Barks, D. W. Knight, and R. J. Middleton, *Tetrahedron Lett.*, 2000, **41**, 4447.
5. a) J. Clayden, D. W. Watson, M. Helliwell, and M. Chambers, *Chem. Commun.*, 2003, 2582. b) J. Ichikawa, K. Sakoda, and Y. Wada, *Chem. Lett.*, 2002, **31**, 282 and references cited therein.
6. Dearomatizing anionic cyclizations *via* a closely related mechanism are reported: a) R. A. Bragg and J. Clayden, *Tetrahedron Lett.*, 1999, **40**, 8323 and 8327. b) J. Clayden, J. C. Menet, and J. D. Mansfield, *Org. Lett.*, 2000, **2**, 4229. c) A. Ahmed, J. Clayden, and M. Rowley, *Tetrahedron Lett.*, 1998, **39**, 6103. d) J. Clayden and J. M. Rowley, *Chem. Commun.*, 1998, 297. e) J. Clayden and S. A. Yasin, *Chem. Commun.*, 1999, 231. f) J. Clayden, F. E. Knowles, and C. J. Menet, *Synlett*, 2003, 1701 and references cited therein.