


INTRAMOLECULAR PAUSON-KHAND REACTION OF 3-ALKYNYL-1-ALKYLIDENECYCLES: A CONVENIENT SYNTHESIS OF [5.n.1.0^{1,5}] TRICYCLIC COMPOUNDS

Miyuki Ishizaki,* Mina Masamoto, Osamu Hoshino, Kiyoshi Nishitani, and Hiroshi Hara*

Faculty of Pharmaceutical Sciences, Tokyo University of Science, 2641, Yamazaki, Noda-shi, Chiba 278-8510, Japan

Abstract – An intramolecular Pauson-Khand reaction of various 3-alkynyl-1-alkylidenecyclic compounds was performed to give the corresponding [5.n.1.0^{1,5}] tricyclic compounds ($n = 2-4$). A facile construction of the core structure in cedrene terpenoids was also investigated in detail and formal total synthesis of α - and β -cedrenes was accomplished.


The Pauson-Khand reaction¹ is known as one of the attractive methods for the construction of polycyclic compounds, including cyclopentenone structures. The usefulness of the reaction has been demonstrated in its application to the total synthesis of natural products.² Recent work in our laboratory has revealed a facile construction of bi- and tri-cyclic compounds by the Pauson-Khand reaction of *exo*-cyclic enynes.³ Especially, the intramolecular Pauson-Khand reaction of 2-alkynyl-1-methylenecyclic compounds (**A**) (*exo*-cyclic enynes) formed angular type tricycles (**B**) in good yields (Scheme 1).^{3a-e} We envisage that the reaction would be applicable to the synthesis of the [5.3.1.0^{1,5}] tricyclic skeleton (**D**), which is the core structure of the widely known cedrene terpenoids, from 3-propynyl-1-alkylidenecyclic compounds (**C**)



Scheme 1

(*exo*-cyclic enynes). In this paper, we wish to describe our investigation on the generality of the synthesis of [5.*n*.1.0^{1,5}] tricyclic skeleton (**D**) (*n* = 2-4) by the intramolecular Pauson-Khand reaction of various 3-alkynyl-1-alkylidenecyclic compounds (**C**) (Scheme 1) and its application to construct various functionalized core structures of cedrene terpenoids.^{4,5}

Various Pauson-Khand precursors (**1a-f**) bearing the *exo*-methylene group were synthesized as shown in Scheme 2. Thus, Wittig olefination of ketones (**3a-c**)⁶ afforded enynes (**1a-c**) in 64-69% yields. An *N*-Cbz piperidine derivative (**1d**) was obtained by propynylation⁶ of the enone (**4**)⁷ followed by Wittig olefination. Reduction of **1d** furnished *N*-methyl piperidine (**1e**). A pyrane derivative (**1f**) was synthesized from the enone (**5**)⁸ in four steps.















Scheme 2

With enynes in hand, the intramolecular Pauson-Khand reaction of enynes (**1a-f**) was examined by the following three procedures: refluxing in toluene (Method A); heating at 83°C with BuSMe (3.5 eq.)^{9a} in CHCl₂CHCl₂ (Method B); oxidative treatment with *N*-methylmorpholine *N*-oxide^{9b} (9-12 eq.) at room temperature in CH₂Cl₂ (Method C). The results are shown in Table 1. The reaction of methylenecyclohexane (**1b**) gave the corresponding [5.3.1.0^{1,5}] tricyclic compound (**2b**) in moderate to good yields (57-79%) by Methods A-C. A similar reaction of methylenecycloheptane (**1c**) afforded [5.4.1.0^{1,5}] tricycle (**2b**) in moderate yields (55-67%) by Methods A and B, whereas the reaction by Method C furnished **2b** in 21% yield. Unexpectedly, the reaction of methylenecyclopentane (**1a**) resulted in the formation of **2a** in low yields despite employment of Methods A-C. The result would be due to a

highly strained tricyclic system in **2a**. In the case of aza and oxa derivatives (**1d-f**), although the reaction of benzyloxy carbamate (**1d**) afforded a tricyclic compound (**2d**) in moderate yields (47-60%), a similar reaction of *N*-methylpiperidine (**1e**) and pyrane (**1f**) formed an intractable mixture and **2e** and **2f** were attained in low yields (8-16%). It was found that generally reactions under thermal conditions (Methods A and B) except for **1a** were superior to an oxidative condition (Method C).


Much attention has been focused on naturally occurring cedrene-type terpenoids, as represented by α - and β -cedrene (**9a,b**), because of their intriguing [5.3.1.0^{1,5}] tricyclic skeleton and much synthetic work on

Table 1. Intramolecular Pauson-Khand reaction of various enynes (**1a-f**).

Entry	Substrate	Method ^a	Time (h)	Product	Yield (%)
1	 1a	A	4	 2a	4
2		B	8		12
3		C	2		13
4	 1b	A	4	 2b	73
5		B	10		79
6		C	2		57
7	 1c	A	4	 2c	55
8		B	10		67
9		C	2		21
10	 1d	A	4	 2d	54
11		B	8		60
12		C	2		47
13	 1e	A	4	 2e	16
14		B	10		18
15		C	2		8
16	 1f	A	4	 2f	14
17		B	10		15
18		C	2		10















a) Method A; refluxing in toluene. Method B; heating at 83°C with BuSMe (3.5 eq.) in CHCl₂CHCl₂. Method C; oxidative treatment with *N*-methylmorpholine *N*-oxide (9 eq. for **1a-c**, **e** and 12 eq. for **1d**, **f**) at room temperature in CH₂Cl₂.

them has been achieved.^{5,10} As mentioned above, [5.n.1.0^{1.5}] tricycles could be obtained by the intramolecular Pauson-Khand reaction of 3-alkynyl-1-methylenecycles. Thus, we turned our attention to synthesize functionalized [5.3.1.0^{1.5}] tricyclic compounds by the reaction of 3-alkynyl-1-alkylidenecyclohexanes. For short-step synthesis, we adopted the Nicholas reaction¹¹ to obtain Pauson-Khand precursors bearing the carbonyl group on a cyclohexane ring (Scheme 3). Thus, 4-alkylidenecyclohexanones (**10a,b**)¹² were converted to trimethylsilyl enol ethers (**11a,b**), of which the Nicholas reaction reported by Schreiber¹³ with the alkyne-cobalt complex (**12a**) afforded the expected Pauson-Khand reaction precursors (**13a,b**) in good yields. Similar reaction of **11a,b** with **12b** produced alkyne-cobalt complexes (**13c,d**) in moderate yields. Because **11b** was obtained as a mixture of regio-isomers (5 : 4 by ¹H-NMR spectrometry), alkyne-cobalt complexes (**13b,d**) were also attained as a mixture of diastereomers. The ratio of diastereomers was estimated by ¹H-NMR spectroscopic analysis (52 : 48 for **13b** and 59 : 41 for **13d**), although the geometry of the olefinic part could not be determined. The Pauson-Khand reaction of synthesized alkyne-cobalt complexes bearing alkylidene group was examined. As shown in Table 2, an expected [5.3.1.0^{1.5}] tricyclic compound (**15a**) was obtained by the Pauson-Khand reaction of **13a** in 52-60% yields under thermal conditions (Methods A and B), whereas a low yield was observed under an oxidative condition (Method C). Unfortunately, a similar reaction of **13b** bearing tri-substituted olefinic moiety produced **15b** in low yields despite Methods A-C, which is thought to be attributable to spatially small carbonyl groups in **13a,b**, because Pauson *et al.* have reported that a similar reaction of the acetal substrate gave a tricyclic compound in high yield.⁵ Therefore, transformation¹⁴ of ketones (**13a,b**) to acetals (**14a,b**) might cause favorable conformation in a transition state based on a reactive rotamer effect (Scheme 4).¹⁵ As expected, yields of the Pauson-Khand reaction of




Scheme 2

Table 2. Intramolecular Pauson-Khand reaction of various alkyne-cobalt complexes (**13a-d** and **14a-c**).

Entry	Substrate	Method ^a	Time (h)	Product	Yield (%)
1	 13a	A	4	 15a	52
2		B	8		60
3		C	2		19
4	 13b	A	4	 15b	12 ^b
5		B	10		30 ^b
6		C	2		4 ^b
7	 13c	A	4	 15c	62
8		B	10		72
9		C	2		26
10	 13d	A	4	 15da : R = α -Me 15db : R = β -Me	14+16 ^c
11		B	10		25+25 ^c
12		C	2		3+4 ^c
13	 14a	A	4	 16a	78
14		B	10		85
15		C	2		48
16	 14b	A	4	 16b	42
17		B	8		60
18		C	2		22
16	 14c	A	4	 16c	62
17		B	8		80
18		C	2		42


a) Method A; refluxing in toluene. Method B; heating at 83°C with BuSMe (3.5 eq.) in CHCl₂CHCl₂. Method C; oxidative treatment with *N*-methylmorpholine *N*-oxide (12 eq.) at room temperature in CH₂Cl₂. b) Obtained as a mixture of diastereomers. c) Isolated yield of **15da** and **15db**.



Scheme 4

14a,b under thermal conditions remarkably improved to 78-85% for **16a** and 42-60% for **16b**. We also synthesized a chiral acetal (**14c**) from **13a**. However, bulky 1,2-diphenylethanol prevented complete conversion to a chiral acetal (**14c**) and **14c** was obtained as an inseparable 54 : 46 mixture (^1H -NMR spectrometry). The Pauson-Khand reaction of **14c** smoothly proceeded by Method B to afford **16c** in 80% yield as a 52 : 48 mixture of diastereomers.

Next, the Pauson-Khand reaction of dimethyl derivatives (**13c,d**) was investigated. Although, unfortunately, acetalization of ketones (**13c,d**) did not occur, the reaction of **13c** gave rise to a tricyclic product (**15c**) in moderate to good yields (62-72%) by Methods A and B. A similar reaction of the tri-substituted olefin (**13d**) produced a *ca.* 1 : 1 ratio of desired products (**15da,db**) in 50% combined yield by Method B. The results suggested that the bulky dimethyl group in **13c,d** would set a favorable conformation in a transition state due to the reactive rotamer effect¹⁵ without acetalization. The stereochemistry of **15da** and **15db** was determined by conversion to known compounds (**17a,b**).⁵ Since Pauson *et al.* have reported formal total synthesis of α - and β -cedrenes (**9a,b**) from **17a**,⁵ we also accomplished formal total synthesis of α - and β -cedrenes (Scheme 5).



Scheme 5

In conclusion, we have investigated the intramolecular Pauson-Khand reaction of various 3-alkynyl-1-alkylidenecyclic derivatives to give corresponding $[5.n.1.0^{1,5}]$ tricyclic compounds ($n = 2-4$). A

straightforward synthesis of the alkyne-cobalt complex using Schreiber's method (Nicholas reaction), and subsequently the Pauson-Khand reaction, offered short-step access to the functionalized core structure of cedrene terpenoids. The usefulness of this approach was proved by a five-step synthesis of the known precursor (**17a**) for the total synthesis of α - and β -cedrenes.

ACKNOWLEDGEMENTS

The authors are grateful to Mrs. F. Hasegawa of this faculty for her MS spectral measurements.

EXPERIMENTAL

General. All melting points were measured on a Yanagimoto melting point apparatus and are uncorrected. IR spectra were obtained with a JASCO FT/IR-400. ^1H and ^{13}C NMR spectra were taken with a JEOL JNM AL-300 (300 MHz for ^1H NMR and 75 MHz for ^{13}C NMR) spectrometer in a CDCl_3 solution with tetramethylsilane as the internal standard. MS spectra were measured on a Hitachi M-80 or a JEOL JMS D-300 spectrometer. Column chromatography was performed over silica gel (Merck Kiegesellschaft 60). Preparative TLCs were run on a Merck 5744 plate. Organic extracts were dried over MgSO_4 , unless otherwise noted.

General Procedure for Wittig Reaction of Ketones (3a-d)

To a stirred suspension of methyltriphenylphosphonium bromide (2.5 eq.) and *t*-BuOK (2.4 eq.) in THF was added a solution of ketone (1 eq.) in THF. After being stirred for 2 h, the reaction was quenched with water. The mixture was extracted with Et_2O . The organic extracts were washed with brine, dried, and evaporated under reduced pressure to give a residue, which was taken up in pentane. The precipitate was filtered by suction. The filtrate was evaporated under reduced pressure to give an oily residue, which was purified by bulb-to-bulb distillation to afford corresponding olefins (**1a-c**). In the case of **1d**, column purification (AcOEt : hexane = 1 : 10) was performed.

1-Methylene-3-(2-propynyl)cyclopentane (1a); from **3a** (0.381 g, 3.12 mmol), **1a** (0.240 g, 64.0%) was obtained as a colorless oil; bp 100-110°C/70 mmHg; ^1H NMR δ 4.83-4.86 (2H, m), 2.03-2.52 (7H, m), 1.94 (1H, t, J = 2.6 Hz), 1.70-1.73 (1H, m), 1.39-1.73 (1H, m); IR 3304, 2942, 2117, 1658, 1431 cm^{-1} ; EI MS m/z 120 (M^+); HRMS m/z calcd for C_9H_{12} (M^+) 120.0939, found: 120.0930.

1-Methylene-3-(2-propynyl)cyclohexane (1b); from **3b** (0.669 g, 4.9 mmol), **1b** (0.461 g, 64.6%) was obtained as a colorless oil; bp 80-100°C/30 mmHg; ^1H NMR δ 4.64 (2H, s), 2.13-2.42 (4H, m), 1.98 (1H, t, J = 2.6 Hz), 1.50-1.96 (4H, m), 1.14-1.44 (3H, m); IR 3307, 2929, 2116, 1645 cm^{-1} ; EI MS m/z 134 (M^+); HRMS m/z calcd for $\text{C}_{10}\text{H}_{14}$ (M^+) 134.1096, found: 134.1097.

1-Methylene-3-(2-propynyl)cycloheptane (1c); from **3c** (0.580 g, 3.87 mmol), **1c** (0.396 g, 69.2%) was obtained as a colorless oil; bp 100-110°C/30 mmHg; ^1H NMR δ 4.84 (2H, s), 2.12-2.61 (6H, m), 2.07 (1H,

t, $J = 2.6$ Hz), 1.75-1.98 (4H, m), 1.34-1.66 (3H, m); IR 3308, 2924, 2853, 2118, 1637, 1445 cm^{-1} ; EI MS m/z 147 ($\text{M}^+ - 1$); HRMS m/z calcd for $\text{C}_{11}\text{H}_{15}$ ($\text{M}^+ - 1$) 147.1172, found: 147.1158.

***N*-Benzyloxycarbonyl 4-methylene-2-(2-propynyl)piperidine (1d)**; from **3d** (0.542 g, 2.0 mmol), **1d** (0.460 g, 85.5%) was obtained as a colorless oil; ^1H NMR δ 7.28-7.37 (5H, m), 5.15 (2H, s), 4.89, 4.82 (each 1H, s), 4.57-4.70 (1H, m), 4.18 (1H, br d, $J = 11.5$ Hz), 2.80-2.95 (1H, m), 2.30-2.46 (4H, m), 2.15-2.29 (2H, m), 1.92 (1H, t, $J = 2.6$ Hz); IR 3298, 2118, 1698, 1420 cm^{-1} ; EI MS m/z 269 (M^+); HRMS m/z calcd for $\text{C}_{17}\text{H}_{19}\text{NO}_2$ (M^+) 269.1416, found: 269.1427.

***N*-Benzyloxycarbonyl 4-oxo-2-(2-propynyl)piperidine (3d)**. To a stirred solution of **4** (2.000 g, 8.66 mmol) and allenyltriphenyltin (6.698 g, 17.2 mmol) in CH_2Cl_2 (50 mL) at -40°C was added a 1M solution of TiCl_4 in CH_2Cl_2 (10.4 mL, 10.4 mmol) over a period of 5 min. After being stirred for 4 h, the reaction was quenched with water. The mixture was extracted with CHCl_3 . The organic extracts were washed with brine, dried, and evaporated under reduced pressure to give a residue, which was taken up in Et_2O and treated with 10% aqueous KF for 1 h. The precipitate was filtered by suction. The filtrate was extracted with Et_2O . The organic extracts were washed with brine, dried, and evaporated under reduced pressure to give a residue, which was purified by column chromatography (AcOEt : hexane = 1 : 10 then 1 : 2) to afford **3d** (1.322 g, 56.3%) and **4'** (0.806 g, 40.3%) as colorless oils; ^1H NMR δ 7.37 (5H, s), 5.18 (2H, s), 4.67-4.76 (1H, m), 4.30-4.45 (1H, m), 4.40-3.55 (1H, m), 2.66 (2H, d, $J = 5.6$ Hz), 2.36-2.60 (4H, m), 1.99 (1H, t, $J = 2.6$ Hz); IR 3288, 2110, 1726, 1697, 1417 cm^{-1} ; EI MS m/z 271 (M^+); HRMS m/z calcd for $\text{C}_{16}\text{H}_{17}\text{NO}_3$ (M^+) 271.1208, found: 271.1206.

***N*-Methyl-4-methylene-3-(2-propynyl)piperidine (1e)**. To a stirred solution of **3d** (0.269 g, 1.0 mmol) in THF (5 mL) at rt was added LiAlH_4 (0.110 g, 2.9 mmol). After being stirred for 0.5 h, the reaction was quenched with saturated an aqueous Na_2SO_4 solution. The precipitates were filtered off and the filtrate was evaporated under reduced pressure to give an oily residue, which was taken up in Et_2O . The ether layer was extracted with 1M HCl. The extracts were washed with Et_2O and made alkaline with 1M NaOH. The aqueous layer was extracted with Et_2O . The organic extracts were dried (K_2CO_3) and evaporated *in vacuo* to afford **1e** (0.141 g, 94.6%) as a colorless oil; ^1H NMR δ 4.71 (2H, s), 2.85 (1H, dt, $J = 4.0, 10.1$ Hz), 2.14-2.42 (8H, m), 2.34 (3H, s), 2.04 (1H, t, $J = 2.6$ Hz); IR 3306, 2941, 2118, 1655 cm^{-1} ; EI MS m/z 149 (M^+); HRMS m/z calcd for $\text{C}_{10}\text{H}_{15}\text{N}$ (M^+) 149.1203, found: 149.1201.

2-(2-Hydroxyethyl)tetrahydropyran-4-one (6). A mixture of **5** (2.88 g, 12.4 mmol) and 10% Pd/C (0.8 g) in AcOEt (25 mL) was stirred for 22 h at rt. After the mixture was filtered, the filtrate was evaporated under reduced pressure to give a residue, which was purified by column chromatography (AcOEt : hexane = 1 : 3 then AcOEt) to afford **6** (1.16 g, 64.9%) as a colorless oil; ^1H NMR δ 4.28 (1H, dd, $J = 7.6, 11.6$ Hz), 3.75-3.86 (3H, m), 3.67 (1H, ddd, $J = 3.0, 11.6, 12.2$ Hz), 2.57 (1H, ddd, $J = 7.6, 12.2, 14.2$ Hz), 2.28-2.64 (4H, m), 1.69-1.93 (2H, m); IR 3420, 1715 cm^{-1} ; EI MS m/z 144 (M^+); HRMS m/z calcd for

C₇H₁₂O₃ (M⁺) 144.0786, found: 144.0782.

2-(2-Hydroxyethyl)-4-methylenetetrahydropyran (7). To a stirred suspension of methyltriphenylphosphonium bromide (7.50 g, 21 mmol) and *t*-BuOK (2.22 g, 19.8 mmol) in THF (100 mL) at rt was added a solution of **6** (1.139 g, 7.9 mmol) in THF (20 mL). After being stirred for 1 h, the reaction was quenched with water. The mixture was extracted with Et₂O. The organic extracts were washed with brine, dried, and evaporated under reduced pressure to give a residue, which was purified by column chromatography (AcOEt : hexane = 1 : 3 then 1 : 1) to afford **7** (0.960 g, 85.5%) as a colorless oil; ¹H NMR δ 4.77, 4.76 (each 1H, d, *J* = 2.0 Hz), 4.11 (1H, ddd, *J* = 1.3, 5.6, 10.9 Hz), 3.82 (2H, t, *J* = 5.4 Hz), 3.38-3.58 (2H, m), 2.07-2.40 (5H, m), 1.71-1.91 (2H, m); IR 3389, 1652 cm⁻¹; EI MS *m/z* 142 (M⁺); HRMS *m/z* calcd for C₈H₁₄O₂ (M⁺) 142.0994, found: 142.0991.

2-Formylmethyl-4-methylenetetrahydropyran (8). To a stirred solution **7** (0.955 g, 6.7 mmol) in CH₂Cl₂ (80 mL) at rt was added Dess-Martin periodinane (3.50 g, 8.2 mmol). After being stirred for 2 h, the reaction was quenched with 10% aqueous Na₂S₂O₃. The mixture was extracted with CHCl₃. The organic extracts were washed with saturated aqueous NaHCO₃ and brine, dried, and evaporated under reduced pressure to give a residue, which was purified by column chromatography (AcOEt : hexane = 1 : 10) to afford **8** (0.765 g, 81.4%) as a colorless oil; ¹H NMR δ 9.87 (1H, dd, *J* = 2.0, 2.6 Hz), 4.76 (2H, t, *J* = 1.8 Hz), 4.65 (1H, ddd, *J* = 1.7, 5.3, 10.9 Hz), 3.77-3.86 (1H, m), 3.27 (1H, ddd, *J* = 3.0, 10.9, 13.3 Hz), 2.64 (1H, ddd, *J* = 2.6, 7.9, 16.3 Hz), 2.51 (1H, ddd, *J* = 1.7, 4.6, 16.3 Hz), 2.02-2.36 (4H, m); IR 2941, 2851, 1726, 1654 cm⁻¹; EI MS *m/z* 140 (M⁺); HRMS *m/z* calcd for C₈H₁₂O₂ (M⁺) 140.0837, found: 140.0842.

4-Methylene-2-(2-propynyl)tetrahydropyran (1f). To a stirred suspension of PPh₃ (5.294 g, 20.2 mmol) in CH₂Cl₂ (15 mL) at 0°C was added a solution of CBr₄ (3.349 g, 10 mmol) in CH₂Cl₂ (15 mL). After being stirred for 10 min, a solution of **8** (0.707 g, 5.1 mmol) in CH₂Cl₂ (10 mL) was added. The mixture was further stirred for 2 h at rt and evaporated under reduced pressure to give a residue. Et₂O was added and precipitates were filtered off. The filtrate was evaporated under reduced pressure to give a residue, which was purified by column chromatography (AcOEt : hexane = 1 : 5) to afford 4-methylene-2-(3,3-dibromo-2-propenyl)tetrahydropyran (1.208 g, 80.8%) as a colorless oil; ¹H NMR δ 6.52 (1H, t, *J* = 6.9 Hz), 4.75, 4.74 (each 1H, s), 4.05 (1H, ddd, *J* = 4, 5.6, 9.6 Hz), 3.31-3.42 (2H, m), 1.98-2.34 (6H, m); IR 2896, 2848, 1653 cm⁻¹.

To a stirred solution of the dibromide (1.186 g, 4.01 mmol) in Et₂O (10 mL) at 0°C was added 1M MeLi in Et₂O (10 mL, 10 mmol). After being stirred for 1 h, the reaction was quenched with water. The mixture was extracted with Et₂O. The organic extracts were washed with brine, dried, and evaporated under reduced pressure to give a residue, which was purified by bulb-to-bulb distillation to afford **1f** (0.487 g, 89.4%) as a colorless oil; bp 130°C/75 mmHg; ¹H NMR δ 4.76 (2H, s), 4.10 (1H, ddd, *J* = 1.7, 5.7, 10.9

Hz), 3.36-3.47 (2H, m), 2.28-2.47 (4H, m), 2.10-2.25 (2H, m), 2.04 (1H, t, $J = 2.6$ Hz); IR 3299, 2942, 2853, 2121, 1654 cm^{-1} ; EI MS m/z 136 (M^+); HRMS m/z calcd for $\text{C}_9\text{H}_{12}\text{O}$ (M^+) 136.0888, found: 136.0894.

General Procedures for Pauson-Khand Reaction of *exo*-Cyclic Enynes (1a-f). A mixture of enyne (1 eq.) and $\text{Co}_2(\text{CO})_8$ (1.15 eq.) in toluene, $\text{CH}_2\text{ClCH}_2\text{Cl}$ or CH_2Cl_2 , was stirred for 1 h at rt. Method A: refluxing in toluene; Method B: heating at 83°C with BuSMe (3.5 eq.) in $\text{CHCl}_2\text{CHCl}_2$; Method C: oxidative treatment with *N*-methylmorpholine *N*-oxide (9-12 eq.) at rt in CH_2Cl_2 . The solvent was removed under reduced pressure to give a residue, which was diluted with Et_2O . The precipitate was removed by suction filtration through a Celite 545 short pad. Purification of the products obtained by Methods A-C was carried out on preparative TLC.

4,5,6,7-Tetrahydro-3a,6-methanoinden-2-one (2a); oil; ^1H NMR δ 5.69 (1H, t, $J = 1.7$ Hz), 2.45, 2.31 (each 1H, d, $J = 18.2$ Hz), 2.25-2.70 (3H, m), 1.52-1.93 (3H, m), 1.22-1.50 (3H, m); ^{13}C NMR δ 210.4, 194.3, 56.3, 42.5, 38.8, 35.2, 32.0, 30.0, 29.5; IR 2954, 2869, 1711, 1697, 1634 cm^{-1} ; EI MS m/z 148 (M^+); HRMS m/z calcd for $\text{C}_{10}\text{H}_{12}\text{O}$ (M^+) 148.0887, found: 148.0882.

5,6,7,8-Tetrahydro-4H-3a,7-methanoazulen-2-one (2b); oil; ^1H NMR δ 5.81 (1H, s), 2.49-2.64 (3H, m), 2.22 (2H, s), 1.79-1.90 (1H, m), 1.72 (1H, d, $J = 12.2$ Hz), 1.50-1.66 (5H, m), 1.35-1.41 (1H, m); ^{13}C NMR δ 210.4, 193.8, 123.7, 53.2, 49.9, 42.8, 39.3, 36.2, 31.9, 30.7, 20.0; IR 2931, 2856, 1708, 1692, 1624 cm^{-1} ; EI MS m/z 162 (M^+); HRMS m/z calcd for $\text{C}_{11}\text{H}_{14}\text{O}$ (M^+) 162.1044, found: 162.1044.

4,4,6,7,8,9-Hexahydro-3a,8-methanocyclopentacycloocten-2-one (2c); oil; ^1H NMR δ 5.79 (1H, t, $J = 1.7$ Hz), 2.96 (1H, dd, $J = 9.2, 19.6$ Hz), 2.63-2.78 (1H, m), 2.43 (1H, d, $J = 19.6$ Hz), 2.39, 2.82 (each 1H, d, $J = 16.8$ Hz), 2.02 (1H, d, $J = 12.5$ Hz), 1.23-1.95 (9H, m); ^{13}C NMR δ 210.5, 195.0, 124.1, 55.0, 53.7, 42.5, 38.4, 37.0, 36.8, 33.9, 25.3, 25.0; IR 2920, 2855, 1703, 1624 cm^{-1} ; EI MS m/z 176 (M^+); HRMS m/z calcd for $\text{C}_{12}\text{H}_{16}\text{O}$ (M^+) 176.1200, found: 176.1208.

***N*-Benzyloxycarbonyl 3-oxo-8-azatricyclo[5.3.1.0^{1,5}]undec-4-ene (2d);** oil; ^1H NMR δ 7.36 (5H, s), 5.93 (1H, s), 5.15 (2H, s), 4.95-5.20 (1H, m), 4.09 (1H, br s), 2.96-3.15 (1H, m), 2.70-2.87 (2H, m), 2.32 (2H, s), 1.97-2.09 (1H, m), 1.89 (1H, d, $J = 11.2$ Hz), 1.50-1.65 (1H, m), 1.30-1.45 (1H, m); ^{13}C NMR δ 208.8, 188.4, 136.4, 128.4, 128.2, 128.0, 127.8, 124.9, 67.1, 54.8, 52.2, 48.7, 42.2, 38.8, 36.9, 32.8; IR 2954, 1697, 1624 cm^{-1} ; EI MS m/z 297 (M^+); HRMS m/z calcd for $\text{C}_{18}\text{H}_{19}\text{NO}_3$ (M^+) 297.1364, found: 297.1364.

***N*-Methyl-8-azatricyclo[5.3.1.0^{1,5}]undec-4-en-3-one (2e);** oil; ^1H NMR δ 5.85 (1H, s), 3.45 (1H, t, $J = 4.8$ Hz), 2.89 (1H, d, $J = 19.6$ Hz), 2.77 (1H, dd, $J = 4.8, 11.3$ Hz), 2.29 (3H, s), 2.00-2.36 (6H, m), 1.54 (1H, ddd, $J = 2.9, 4.4, 13.9$ Hz), 1.24-1.33 (1H, m); ^{13}C NMR δ 210.2, 190.7, 124.1, 62.8, 51.8, 48.5, 48.4, 43.1, 43.0, 37.2, 24.5; IR 2986, 2852, 1696, 1624 cm^{-1} ; EI MS m/z 177 (M^+); HRMS m/z calcd for $\text{C}_{11}\text{H}_{15}\text{NO}$ (M^+) 177.1152, found: 177.1152.

8-Oxatricyclo[5.3.1.0^{1,5}]undec-4-en-3-one (2f); mp 61-62°C (AcOEt-hexane); ¹H NMR δ 5.92 (1H, s), 4.65-4.68 (1H, m), 3.86 (1H, dd, *J* = 6.9, 12.2 Hz), 3.73 (1H, dd, *J* = 4.3, 12.2 Hz), 2.98 (1H, d, *J* = 20.1 Hz), 2.68 (1H, dd, *J* = 4.6, 20.1 Hz), 2.30 (2H, s), 2.18 (1H, dt, *J* = 6.9, 12.5 Hz), 2.06 (1H, dd, *J* = 2.3, 11.5 Hz), 1.43-1.49 (1H, m), 1.25-1.32 (1H, m); ¹³C NMR δ 209.3, 188.8, 124.6, 76.7, 60.6, 51.9, 48.8, 43.0, 39.0, 32.2; IR 2960, 2857, 1699, 1635 cm⁻¹; EI MS *m/z* 164 (M⁺); HRMS *m/z* calcd for C₁₀H₁₂NO₂ (M⁺) 164.0837, found: 164.0822.

General Procedure for Synthesis of Silyl Enol Ethers (11a,b). To a stirred solution of ketone (**10a** or **10b**) (1 eq.) and Et₃N (1.2 eq.) in CH₂Cl₂ (3 mL per 1 mmol of ketone) at 0°C under argon was added TMSOTf (1.1 eq.). After being stirred for 2 h, the reaction was quenched with saturated aqueous NaHCO₃. The organic layer was washed with brine, dried, and evaporated under reduced pressure to give a residue, which was purified by bulb-to-bulb distillation.

4-Methylene-1-trimethylsilyloxycyclohexene (11a); from **10a** (0.550 g, 5.0 mmol), **11a** (0.890 g, 97.8%) was obtained as a colorless oil; bp 120°C/40 mmHg; ¹H NMR δ 4.78-4.81 (1H, m), 4.74, 4.72 (each 1H, s), 2.73-2.80 (2H, m), 2.33 (2H, t, *J* = 6.6 Hz), 2.10-2.15 (2H, m), 0.17 (9H, s); EI MS *m/z* 182 (M⁺); HRMS *m/z* calcd for C₁₀H₁₈OSi (M⁺) 182.1127, found: 182.1128.

4-Ethylidene-1-trimethylsilyloxycyclohexene (11b); from **10b** (0.992 g, 8.0 mmol), **11b** (1.156 g, 99.9%) was obtained as a colorless oil; bp 140°C/40 mmHg; ¹H NMR δ 5.18-5.32 (1H, m), 4.79-4.82 (1H, m), 2.66-2.73 (2H, m), 2.25-2.55 (2H, m), 2.02-2.10 (2H, m), 1.55-1.65 (3H, m), 0.17 (5H, s), 0.16 (4H, s); EI MS *m/z* 196 (M⁺); HRMS *m/z* calcd for C₁₁H₂₀OSi (M⁺) 196.1283, found: 196.1281.

General Procedure for Synthesis of Pauson-Khand Precursors (13a-d). To a stirred solution of the silyl enol ether (**11a** or **11b**) (5.0 mmol) and the cobalt complex (**12a** or **12b**) (6.0 mmol) in CH₂Cl₂ (60 mL) at -78°C under argon was added a 0.95M hexane solution of Et₂AlCl (5.9 mmol). The mixture was warmed up to 0°C for 24 h. After the reaction was quenched with water, the mixture was extracted with CHCl₃. The organic extracts were washed with brine, dried, and evaporated under reduced pressure to give a residue, which was purified by column chromatography (benzene : hexane = 1 : 2 then 1 : 1) to afford the corresponding alkyne-cobalt complex (**13a-d**).

13a; from **12a**, **13a** (1.584 g, 72.9%) was obtained as black mass solid; ¹H NMR δ 6.09 (1H, s), 5.00 (2H, s), 3.54 (1H, d, *J* = 15.3 Hz), 2.25-3.05 (8H, m); IR 2090, 2050, 1993, 1704 cm⁻¹; EI MS *m/z* 406 (M⁺-CO); HRMS *m/z* calcd for C₁₅H₁₂O₆Co₂ (M⁺-CO) 406.9298, found: 406.9296.

13b; from **12b**, **13b** (1.576 g, 70.4%) was obtained as a mixture of regio-isomers as a black tar; ¹H NMR δ 6.06 (0.52H, s), 6.02 (0.48H, s), 5.30-5.50 (1H, m), 3.38-3.54 (1H, m), 1.99-3.08 (8H, m), 1.66 (3H, d, *J* = 6.3 Hz); IR 2091, 2050, 1998, 1714 cm⁻¹; EI MS *m/z* 420 (M⁺-CO); HRMS *m/z* calcd for C₁₆H₁₄O₆Co₂ (M⁺-CO) 419.9454, found: 419.9450.

13c; from **12a**, **13c** (1.255 g, 54.4%) was obtained as a black mass solid; ¹H NMR δ 6.02 (0.52H, s), 6.00

(0.48H, s), 5.44 (1H, br s), 3.40-3.53 (1H, m), 1.90-3.10 (8H, m), 1.66 (3H, d, $J = 6.7$ Hz); EI MS m/z 433 ($M^+ - C_2H_4$); HRMS m/z calcd for $C_{16}H_{12}O_7Co_2$ ($M^+ - C_2H_4$) 433.9247, found: 433.9250.

13d; from **12b**, **13d** (1.238 g, 52.9%) was obtained as a mixture of regio-isomers as a black tar; 1H NMR δ 6.13 (1H, s), 5.39 (1H, br s), 2.10-3.20 (6H, m), 1.50-2.67 (4H, m), 1.41 (1.23H, s), 1.44 (1.77H, s), 1.34 (1.23H, s), 1.31 (1.77H, s); IR 2974 cm^{-1} ; EI MS m/z 448 ($M^+ - C_2H_4$); HRMS m/z calcd for $C_{17}H_{14}O_7Co_2$ ($M^+ - C_2H_4$) 447.9408, found: 447.9404.

Convesion of Ketones (13a,b) to Acetals (14a,b). A solution of the ketone (**13a** or **13b**) (1 eq.), trimethyl orthoformate (2 eq.), ethylene glycol (10 eq.), and p -TsOH $\cdot$ H $_2$ O (0.4 eq.) in CH_2Cl_2 (10 mL per 1 mmol of **13a,b**) at rt was stirred for 22 h. The mixture was washed with water and brine, successively. The organic layer was dried and evaporated under reduced pressure to give the corresponding acetal (**14a** or **14b**).

(**14a**); from **13a** (0.679 g, 1.57 mmol), **14a** (0.724 g, 96.7%) was obtained as a black oil; 1H NMR δ 6.06 (1H, s), 4.73 (1H, s), 4.69 (1H, s), 3.99 (4H, br s), 3.18 (1H, d, $J = 15.3$ Hz), 2.71 (1H, dd, $J = 10.0, 15.3$ Hz), 2.55 (1H, dd, $J = 4.5, 13.3$ Hz), 2.19-2.30 (3H, m), 1.77-1.86 (2H, m), 1.50-1.59 (1H, m); IR 2091, 2050, 2016 cm^{-1} ; EI MS m/z 450 ($M^+ - CO$); HRMS m/z calcd for $C_{17}H_{16}O_7Co_2$ ($M^+ - CO$) 449.9560, found: 449.9559.

(**14b**); from **13b** (0.538 g, 1.20 mmol), **14b** (0.558 g, 94.5%) was obtained as a mixture of regio-isomers as a black oil; 1H NMR δ 6.09 (0.53H, s), 6.06 (0.47H, s), 5.20-5.30 (1H, m), 3.99 (4H, br s), 3.12-3.22 (1H, m), 2.10-2.73 (5H, m), 1.41-1.89 (6H, m); IR 2090, 2050, 2015, 1736 cm^{-1} ; EI MS m/z 464 ($M^+ - CO$); HRMS m/z calcd for $C_{18}H_{18}O_7Co_2$ ($M^+ - CO$) 463.9717, found: 463.9733.

Convesion of Ketone (13a) to Acetal (14c). A solution of **13a** (0.100 g, 0.23 mmol), trimethyl orthoformate (30 μ L, 0.27 mmol), (1*S*,2*S*)-hydrobenzoin (0.060 g, 0.28 mmol), and p -TsOH $\cdot$ H $_2$ O (0.018 g, 0.09 mmol) in CH_2Cl_2 (2 mL) at rt was stirred for 22 h. The mixture was washed with water and brine, successively. The organic layer was dried and evaporated under reduced pressure to give a residue, which was purified by TLC (hexane : AcOEt = 8 : 1) to afford **14c** (0.073 g, 50.4%) as a mixture of regio-isomers as a black oil and **13a** (0.042 g, 42.2%). 1H NMR δ 7.18-7.31 (10H, m), 6.10 (0.54H, s), 6.09 (0.46H, s), 4.65-4.88 (4H, m), 3.57 (0.54H, d, $J = 14.8$ Hz), 3.46 (0.46H, d, $J = 14.5$ Hz), 3.01 (0.54H, dd, $J = 10.2, 14.8$ Hz), 2.87 (0.46H, dd, $J = 10.7, 14.5$ Hz), 2.55-2.74 (1H, m), 2.30-2.50 (3H, m), 1.80-2.20 (3H, m); IR 2090, 2053, 2018 cm^{-1} ; EI MS m/z 602 ($M^+ - CO$); HRMS m/z calcd for $C_{29}H_{24}O_7Co_2$ ($M^+ - CO$) 602.0816, found: 602.0811.

General Procedures for Pauson-Khand Reaction of *exo*-Cyclic Enynes (13a-d, 14a-c). The reaction of enyne (1 eq.) in toluene, CH_2ClCH_2Cl or CH_2Cl_2 , was performed. Method A: refluxing in toluene; Method B: heating at 83°C with BuSMe (3.5 eq.) in $CHCl_2CHCl_2$; Method C: oxidative treatment with *N*-methylmorpholine *N*-oxide (12 eq.) at rt in CH_2Cl_2 . The solvent was removed under reduced pressure to give a residue, which was diluted with Et $_2$ O. The precipitate was removed by suction filtration through a

Celite 545 short pad. Purification of the products obtained by Methods A-C was carried out on preparative TLC.

4,5,7,8-Tetrahydro-3a,7-methanoazulene-2,6-dione (15a); mp 73-74°C (AcOEt-hexane); ¹H NMR δ 6.00 (1H, t, *J* = 1.6 Hz), 3.04 (1H, t, *J* = 5.4 Hz), 2.86-2.90 (2H, m), 2.33-2.59 (2H, m), 2.43 (2H, s), 2.21, 2.13 (each 1H, d, *J* = 11.4 Hz), 1.88-2.04 (1H, m), 1.68-1.78 (1H, m); ¹³C NMR δ 210.2, 208.8, 188.6, 125.7, 52.3, 51.2, 48.4, 40.5, 36.7, 34.7, 31.5; IR 2960, 2869, 1716, 1691, 1624 cm⁻¹; EI MS *m/z* 176 (M⁺); HRMS *m/z* calcd for C₁₁H₁₂O₂ (M⁺) 176.0837, found: 176.0847.

4,5,7,8-Tetrahydro-3-methyl-3a,7-methanoazulene-2,6-dione (15b); oil; ¹H NMR δ 6.04 (0.25H, t, *J* = 1.7 Hz, olefinic H), 5.93 (0.75H, t, *J* = 1.7 Hz, olefinic H), 2.75-3.10 (3H, m), 2.30-2.60 (3H, m), 1.41-2.23 (4H, m), 1.16 (0.75H, d, *J* = 7.3 Hz), 1.09 (2.25H, d, *J* = 7.6 Hz); ¹³C NMR δ 212.8, 210.7, 187.3, 124.6, 123.4, 56.1, 53.1, 51.6, 50.9, 50.8, 40.9, 38.1, 35.9, 35.0, 34.6, 33.3, 31.8, 31.7, 29.7, 14.3, 9.3; IR 2964, 2932, 2872, 1707, 1696, 1633 cm⁻¹; EI MS *m/z* 190 (M⁺); HRMS *m/z* calcd for C₁₂H₁₄O₂ (M⁺) 190.0994, found: 190.0991.

4,5,7,8-Tetrahydro-8,8-dimethyl-3a,7-methanoazulene-2,6-dione (15c); mp 134-135°C (AcOEt-hexane); ¹H NMR δ 5.92 (1H, s), 2.61 (1H, d, *J* = 3.6 Hz), 2.53, 2.40 (each 1H, d, *J* = 17 Hz), 1.95-2.45 (5H, m), 1.60-1.73 (1H, m), 1.33, 1.29 (each 3H, s); ¹³C NMR δ 210.4, 208.5, 198.9, 123.1, 63.7, 52.8, 50.4, 42.4, 36.9, 35.8, 35.6, 31.3, 22.5; IR 2969, 2872, 1700, 1620 cm⁻¹; EI MS *m/z* 204 (M⁺); HRMS *m/z* calcd for C₁₃H₁₆O₂ (M⁺) 204.1150, found: 204.1142.

4,5,7,8-Tetrahydro-3,8,8-trimethyl-3a,7-methanoazulene-2,6-dione (15da); mp 167-168°C (AcOEt-hexane); ¹H NMR δ 5.96 (1H, s), 2.63 (1H, d, *J* = 3.9 Hz), 2.56 (1H, dd, *J* = 9.0, 18.6 Hz), 2.54 (1H, q, *J* = 7.2 Hz), 2.36 (1H, dt, *J* = 9.8, 18.6 Hz), 2.10, 2.03 (each 1H, d, *J* = 10.8 Hz), 2.05-2.12 (1H, m), 1.43 (1H, t, *J* = 11.4 Hz), 1.34, 1.27 (each 3H, s), 1.15 (3H, d, *J* = 7.2 Hz); ¹³C NMR δ 210.4, 197.2, 172.8, 122.3, 64.2, 56.4, 54.9, 42.7, 37.5, 35.7, 32.8, 31.2, 22.2, 9.3; IR 2967, 2932, 2871, 1700, 1621 cm⁻¹; EI MS *m/z* 218 (M⁺); HRMS *m/z* calcd for C₁₄H₁₈O₂ (M⁺) 218.1305, found: 218.1303.

4,5,7,8-Tetrahydro-3,8,8-trimethyl-3a,7-methanoazulene-2,6-dione (15db); mp 156-157°C (AcOEt-hexane); ¹H NMR δ 5.84 (1H, s), 2.61 (1H, d, *J* = 4.9 Hz), 2.54 (1H, dd, *J* = 9.0, 18.6 Hz), 2.36 (1H, q, *J* = 7.6 Hz), 2.33-2.43 (1H, m), 2.16-2.24 (2H, m), 1.83 (1H, d, *J* = 12.4 Hz), 1.63-1.70 (1H, m), 1.33, 1.26 (each 3H, s), 1.11 (3H, d, *J* = 7.6 Hz); ¹³C NMR δ 212.7, 210.6, 197.8, 120.9, 63.4, 56.8, 52.8, 42.7, 37.8, 36.0, 32.4, 30.8, 22.3, 14.8; IR 2973, 2956, 2937, 2868, 1701, 1621 cm⁻¹; EI MS *m/z* 218 (M⁺); HRMS *m/z* calcd for C₁₄H₁₈O₂ (M⁺) 218.1305, found: 218.1312.

6,6-Ethylenedioxy-4,5,7,8-tetrahydro-3a,7-methanoazulene-2-one (16a); mp 113-114°C (AcOEt-hexane); ¹H NMR δ 5.87 (1H, s), 3.85-4.08 (4H, m), 2.76 (1H, d, *J* = 19.8 Hz), 2.65 (1H, dd, *J* = 1.0, 18.1 Hz), 2.44 (1H, t, *J* = 5.8 Hz), 2.28 (2H, s), 2.21 (1H, dd, *J* = 1.7, 5.8 Hz), 1.91-2.02 (1H, m), 1.66-1.72 (2H, m), 1.50 (1H, ddd, *J* = 3.0, 5.6, 11.6 Hz), 1.31-1.39 (1H, m); ¹³C NMR δ 209.9, 191.7, 124.7, 109.6,

64.7, 64.1, 52.0, 48.7, 44.9, 38.7, 35.7, 30.7, 30.2; IR 1696, 1621 cm^{-1} ; EI MS m/z 220 (M^+); HRMS m/z calcd for $\text{C}_{13}\text{H}_{16}\text{O}_3$ (M^+) 220.1097, found: 220.1097.

6,6-Ethylenedioxy-4,5,7,8-tetrahydro-3-methyl-3a,7-methanoazulene-2-dione (16b); mp 113–118°C; ^1H NMR δ 5.90 (0.26H, s), 5.80 (0.74H, s), 3.85–4.08 (4H, m), 2.79 (0.26H, s), 2.72 (0.74H, s), 2.42–2.63 (2H, m), 2.17–2.28 (1H, m), 1.26–1.84 (6H, m), 1.10 (0.78H, d, $J = 7.3$ Hz), 1.02 (2.22H, d, $J = 7.6$ Hz); ^{13}C NMR δ 213.9, 190.4, 189.7, 123.5, 122.5, 110.1, 109.9, 64.8, 64.2, 55.8, 55.3, 53.3, 51.1, 45.2, 44.5, 39.0, 37.2, 34.0, 31.9, 31.0, 30.9, 30.4, 29.9, 29.6, 14.1; IR 1691, 1627 cm^{-1} ; EI MS m/z 234 (M^+); HRMS m/z calcd for $\text{C}_{14}\text{H}_{18}\text{O}_3$ (M^+) 234.1255, found: 234.1264.

6,6-(1,2-Diphenylethylenedioxy)-4,5,7,8-tetrahydro-3a,7-methanoazulene-2-one (16c); amorphous solid; ^1H NMR δ 7.15–7.43 (10H, m), 5.93 (0.52H, s), 5.92 (0.48H, s), 4.67–4.84 (2H, m), 2.61–3.04 (3H, m), 1.90–2.52 (6H, m), 1.40–1.75 (2H, m); ^{13}C NMR δ 209.9, 191.6, 191.5, 137.0, 136.8, 136.4, 136.1, 128.5, 128.4, 128.3, 126.8, 126.7, 126.6, 126.5, 125.0, 124.9, 110.6, 110.2, 85.7, 84.8, 52.4, 52.0, 48.8, 48.7, 46.4, 39.1, 38.4, 35.8, 35.6, 32.1, 31.4, 30.9, 30.6; IR 1696, 1626 cm^{-1} ; EI MS m/z 372 (M^+); HRMS m/z calcd for $\text{C}_{25}\text{H}_{24}\text{O}_3$ (M^+) 372.1725, found: 372.1720.


6,6-Ethylenedioxy-4,5,7,8-tetrahydro-3,8,8-trimethyl-3a,7-methanoazulene-2,6-dione (17a); mp 86°C (hexane); ^1H NMR δ 5.83 (1H, s), 3.80–4.06 (4H, m), 2.30 (1H, q, $J = 7.2$ Hz), 2.23 (1H, dd, $J = 11.7$ Hz), 1.98 (1H, d, $J = 5.1$ Hz), 1.69–1.87 (4H, m), 1.48, 1.20 (each 3H, s), 1.14–1.25 (1H, m), 1.09 (3H, d, $J = 7.2$ Hz); ^{13}C NMR δ 211.5, 200.3, 120.9, 110.0, 64.8, 63.5, 56.5, 56.3, 55.0, 42.3, 37.8, 33.7, 32.8, 30.9, 22.6, 8.8; IR 2969, 2932, 2883, 1706, 1623 cm^{-1} ; EI MS m/z 262 (M^+); HRMS m/z calcd for $\text{C}_{16}\text{H}_{22}\text{O}_3$ (M^+) 262.1569, found: 262.1570.

6,6-Ethylenedioxy-4,5,7,8-tetrahydro-3,8,8-trimethyl-3a,7-methanoazulene-2,6-dione (17b); mp 84°C (hexane); ^1H NMR δ 5.77 (1H, s), 3.79–4.05 (4H, m), 2.24 (1H, q, $J = 7.5$ Hz), 1.90–2.03 (3H, m), 1.69–1.79 (3H, m), 1.48, 1.19 (each 3H, s), 1.22–1.34 (1H, m), 1.01 (3H, d, $J = 7.5$ Hz); ^{13}C NMR δ 213.8, 201.2, 119.6, 110.1, 64.8, 63.5, 57.0, 55.6, 52.5, 42.3, 39.1, 32.7, 32.5, 31.6, 22.6, 14.8; IR 2968, 2926, 2876, 1699, 1623 cm^{-1} ; EI MS m/z 262 (M^+); HRMS m/z calcd for $\text{C}_{16}\text{H}_{22}\text{O}_3$ (M^+) 262.1569, found: 262.1574.

REFERENCES AND NOTES

1. For reviews, (a) J. Blanco-Urgoiti, L. Anorbe, L. Perez-Serrano, G. Dominguez, and J. Perez-Castelles, *Chem. Soc. Rev.*, 2004, **33**, 32. (b) K. M. Brummond and J. L. Kent, *Tetrahedron*, 2000, **56**, 3263. (c) A. J. Fletcher and S. D. R. Christie, *J. Chem. Soc., Perkin Trans. 1*, 2000, 1657.
2. For some recent examples; (a) D. A. Ockey, M. A. Lewis, and N. E. Shore, *Tetrahedron*, 2003, **59**, 5377. (b) I. Nomura and C. Mukai, *Org. Lett.*, 2002, **4**, 4304. (c) C. Mukai, M. Kobayashi, I. J. Kim, and M. Hanaoka, *Tetrahedron*, 2002, **58**, 5335. (d) J. Cassayne, F. Gagosz, and S. Z. Zard, *Angew.*

- Chem., Int. Ed.*, 2002, **41**, 1783. (e) J. Marco-Contelles, J. Ruiz-Caro, E. Mainetti, P. Devin, L. Fensterbank, and M. Malacria, *Tetrahedron*, 2002, **58**, 1147. (f) M. E. Krafft, Y. Y. Cheung, and K. A. Abboud, *J. Org. Chem.*, 2001, **66**, 7443. (g) J. Cassayre and S. Z. Zard, *J. Organometal. Chem.*, 2001, **624**, 316. (h) M. E. Krafft, Y. Y. Cheung, and C. A. Juliano-Capucão, *Synthesis*, 2000, 1020.
3. (a) M. Ishizaki, K. Iwahara, K. Kyoumura, and O. Hoshino, *Synlett*, 1999, 587. (b) M. Ishizaki, K. Iwahara, Y. Niimi, H. Satoh, and O. Hoshino, *Tetrahedron*, 2001, **57**, 2729. (c) M. Ishizaki, Y. Niimi, and O. Hoshino, *Chem. Lett.*, 2001, 546. (d) M. Ishizaki, Y. Niimi, and O. Hoshino, *Tetrahedron Lett.*, 2003, **44**, 6029. (e) M. Ishizaki and O. Hoshino, *J. Org. Synth. Chem. Jpn.*, 2003, **61**, 1166. (f) M. Ishizaki, Y. Kasama, M. Zyo, Y. Niimi, and O. Hoshino, *Heterocycles*, 2001, **55**, 1439. (g) M. Ishizaki, M. Zyo, Y. Kasama, Y. Niimi, O. Hoshino, K. Nishitani, and H. Hara, *Heterocycles*, 2003, **60**, 2259.
4. Preliminary communication; M. Ishizaki, M. Masamoto, and O. Hoshino, *Heterocycles*, 2002, **57**, 1409.
5. During development of this work, formal total synthesis of α - and β -cedrenes using the intramolecular Pauson-Khand reaction of *exo*-cyclic enyne (**18**) was reported (Scheme 6); W. J. Kerr, M. McLaughlin, A. J. Morrison, and P. L. Pauson, *Org. Lett.*, 2001, **3**, 2945.



6. J.-i. Haruta, K. Nishi, S. Matsuda, S. Akai, Y. Tamura, and Y. Kita, *J. Org. Chem.*, 1990, **55**, 4853.
7. H. Yamada, S. Aoyagi, and C. Kibayashi, *Tetrahedron Lett.*, 1996, **37**, 8787.
8. I. Paterson, J. D. Smith, and R. A. Ward, *Tetrahedron*, 1995, **51**, 9413.
9. (a) T. Sugihara, M. Yamada, M. Yamaguchi, and M. Nishizawa, *Synlett*, 1999, 771. (b) S. Shambayati, W. E. Crowe, and S. L. Schreiber, *Tetrahedron Lett.*, 1990, **31**, 5289.
10. (a) H.-Y. Lee, S. Lee, D. Kim, K. Byung, J. S. Bahn, and S. Kim, *Tetrahedron Lett.*, 1998, **39**, 7713. (b) S. Das, A. Pal, and D. Mukherjee, *Tetrahedron Lett.*, 1996, **37**, 4421. (c) Y. J. Chen, C. M. Chen, and W. Y. Lin, *Tetrahedron Lett.*, 1993, **34**, 2961. (d) A. Steinmyer, W. Schwede, and F. Bohlmann, *Ann.*, 1988, 925. (e) H. Irie, S. Tanaka, A. Yamamura, Y. Mizuno, H. Tomimatsu, K. Ashizawa, and T. Taga, *Chem. Pharm. Bull.*, 1982, **32**, 2886. (f) K. Pramod and G. S. R. S. Rao, *Chem. Commun.*, 1982, 762. (g) P. A. Wender and J. J. Howbert, *J. Am. Chem. Soc.*, 1981, **103**, 688. (h) E. G. Breitholle and A. G. Fallis, *J. Org. Chem.*, 1978, **43**, 1964. (i) E. G. Breitholle and A. G. Fallis, *Can. J. Chem.*, 1976, **91**, 1557. (j) E. J. Corey, N. N. Girotra, and C. T. Mathew, *J. Am. Chem. Soc.*, 1969, **91**, 1507. (k) G. Stork and F. H. Clarke Jr., *J. Am. Chem. Soc.*, 1955, **77**, 1072.

11. For reviews, (a) B. J. Teobald, *Tetrahedron*, 2002, **58**, 4133. (b) J. R. Green, *Curr. Org. Chem.*, 2001, **5**, 809.
12. A. J. Frontier, S. I. Danishefsky, G. A. Koppel, and D. Meng, *Tetrahedron*, 1998, **54**, 12721.
13. S. L. Schreiber, T. Sammakia, and W. E. Crowe, *J. Am. Chem. Soc.*, 1986, **108**, 3128.
14. X. Su, H. Gao, L. Hang, and Z. Li, *Synth. Commun.*, 1995, **25**, 2807.
15. M. E. Jung and J. Gervay, *J. Am. Chem. Soc.*, 1991, **113**, 224.