

HETEROCYCLES, Vol. 65, No. 12, 2005, pp. 2861 - 2870

Received, 11th July, 2005, Accepted, 13th October, 2005, Published online, 14th October, 2005

REACTIONS OF 4-PENTENOIC ACID WITH SULFENYL CATIONS
GENERATED ELECTROCHEMICALLY FROM BISQUINOLINYL AND
BISPYRIDINYL DISULFIDES [#]

Krzysztof Marciniak, ^{a)} Andrzej Maślankiewicz, ^{a)} * Maria J. Maślankiewicz, ^{b)}
and Zbigniew Ciunik ^{c)}

^{a)} Department of Organic Chemistry, The Medical University of Silesia
Jagiellońska 4, 41-200 Sosnowiec, Poland. ^{b)} Institute of Chemistry, University of
Silesia, Szkolna Str. 9, 40-006 Katowice, Poland. ^{c)} Institute of Chemistry,
University of Wrocław, F. Joliot-Curie 14, 50-383 Wrocław, Poland

E-mail: maslankiewicz@slam.katowice.pl

Abstract - Lactonization of 4-pentenoic acid to 5-(azinythio)methyloxolan-2-ones (**8**) was performed by addition of electrochemically generated azinylsulfenyl cations starting from the respective 3,3'-bis(pyridinyl or quinolinyl) disulfides (**2**) or (**5**) and 4,4'-bis(7-chloroquinolinyl or 3-methylthioquinolinyl) disulfides (**6b**), (**6c**) and using a bromide ion as a redox catalyst. Sulfenyl cations generated in the same manner from 2,2'-bispyridinyl and 2,2'-bisquinolinyl disulfides (**1**) or (**4**) reacted with 4-pentenoic acid to form thiazolo[3,2-*a*]pyridinio- or quinoliniopropanoate (**9a**) or (**9b**). In the case of 8,8'-bisquinolinyl disulfide (**7**), oxolan-2-one (**8e**) was accompanied by 2,3-dihydro-1,4-thiazinequinolinio derivative (**10**). 4,4'-Bis(pyridinyl nor quinolinyl) disulfides (**3**) and (**6a**) did not give such products.


INTRODUCTION

Electrochemically generated phenylsulfenyl cation reacts with 4-pentenoic acid to form 5-(phenylthio)methyloxolan-2-one. ¹ The same reaction course was observed when 3,3'-bis(4-substituted quinolinyl) disulfides were used as a source of quinolinylsulfenyl cation. ² To evaluate the scope of this reaction, we selected further diazinyl disulfides including 2,2'-, 3,3'-, 4,4'-, 4,4'-bis(3-, or 7-substituted quinolinyl)-, and 8,8'-bisquinolinyl disulfides (**4**), (**5**), (**6a,b,c**) and (**7**), as well as 2,2'-, 3,3'-, 4,4'-bispyridinyl disulfides (**1-3**). Our study showed that in the case of 2,2'- and 8,8'- isomers (**1**), (**4**) and (**7**) the neighbor endocyclic nitrogen atom affects the reaction course and induces the cyclization of the reaction mixture components to thiazolo- and thiazinequinolinium or pyridinium betaines (**9**) or (**10**).










RESULTS AND DISCUSSION

We started with the reaction of bis(γ -quinoliny) disulfides (**6**). In the case of bis(3-methylthio) and bis(7-chloro) derivatives (**6b**) and (**6c**), the reaction of electrochemically generated sulfenyl cations with 4-pentenoic acid proceeds *via* expected pathway and gives the respective 5-(4-quinolinythio)methyl-oxolan-2-ones (**8**) in good yields. (Table, Entries 7 and 8).

Scheme 1




Table

Entry	Substrate	Working Potential [V]	Product
1	 (1)	1.25	 (9a) x HBr, 74 % [a]
2	 (2)	1.55	 (8a), (27 %)
3	 (3)	1.91	Passivation of working electrode (anode). Non-consumed substrate (3) (<i>ca.</i> 90 %) was recovered.
4	 (4)	1.80	 (9b) x HBr, 84 % [a]
5	 (5)	1.75	 (8b), 34 %

6		1.85	Passivation of working electrode (anode). Non-consumed substrate (6a), (ca. 90 %) was recovered.
7		1.85	 8c (80 %)
8		1.81	 8d (87 %)
9		1.65	 (8e) (64%) (10) (11 %)

[a] Crude products (**9a**) and (**9b**) contained varied (up to 0.5 mol equiv.) amounts of bromine, therefore the betaines were characterized as hydrogen bromides (**9a**) x HBr and (**9b**) x HBr.

4,4'-Bispyridinyl disulfide (**3**) nor 4,4'-bisquinoliny disulfide (**6a**) did not give products of oxolanone type, although the oxidation potentials of compounds (**6a**) and (**6b**) are identical (see Table). Most probably, the sulfenyl cation (**64⁺**) formed from 3-methylthio derivative (**6b**) is stabilized by interaction with the *ortho*-methylthio group.³




Low yields in the preparation of oxolan-2-ones (**8a**) and (**8b**) from 3,3'-bispyridinyl disulfide (**2**) and 3,3'-bisquinoliny disulfide (**5**) may be due to the low stability of compounds (**8a**) and (**8b**) as compared to the properties of 4-substituted derivatives of **8b**.² All oxolan-2-ones (**8a,b,c,b,e**) showed the positive hydroxyamic test⁴ for the lactone moiety.

Sulfenyl cations generated electrochemically from 2,2'-bispyridinyl and 2,2'-bisquinoliny disulfides (**1**) and (**4**) readily reacted with 4-pentenoic acid to form the products which did not contain a lactone function, and also their other properties (e.g. higher melting points, *R_f* values) differ from those of oxolanones (**8**). Considering the formation of the above mentioned oxolan-2-ones (**8**) in terms of

alkylation of thioazine species, literature review was performed. It revealed that the addition of bromine to 2-allylthiopyridine led either to 3-bromomethyl-2,3-dihydrothiazolo[3,2-*a*]pyridinium bromide (**9e**) (at 0-5 °C) or to 2-bromomethyl-2,3-dihydrothiazolo[3,2-*a*]pyridinium bromide (**9d**) (at room temperature).⁵ Furthermore, 2-bromo-3-(2-pyridinylthio)butyric acid underwent cyclization to 2-methyl-2,3-dihydrothiazolo[3,2-*a*]pyridinium-3-carboxylate.⁶ It suggests that the interaction of α -pyridine – or α -quinolinesulphenyl cations with pentenoic acid may lead to the formation of 2,3-dihydrothiazolo[3,2-*a*]pyridinium species of type (**9**). Furthermore, if the sulfenylation of alkenes follows the Markovnikov rule,¹ the reactions with α,α' -diazinyl disulfides (**1**) and (**4**) should give products with S-CH₂ groups of type (**9c**). However, the chemical shift values δ_{H} of S-CH and N⁺-CH₂ protons in ¹H NMR spectra of our products (**9a**) and (**9b**) fit well the respective data reported⁵ for isomeric salt (**9d**). (Scheme 2)

Scheme 2

The chemical shift values δ_{H} [ppm] for protons from S- and N-methylene or methine groups in salts (**9**) and (**10**)




To prove this supposition a total analysis of the ¹H and ¹³C NMR spectra of products (**9a**, **9b** and **10**) was performed using 1D and 2D NMR spectrometry (including HSQC and HMBC). The crucial data in the structure assignment of **9a**, **9b** and **10** come from the long-range proton-carbon correlations deduced from HMBC spectra. They show the connectivity links between N-methylene protons and both α -azinyl carbons as well as S-CH protons and C_{arom.}-S carbons. (see Scheme 3) Due to the folded shape of the dihydrothiazole and dihydrothiazine moiety in compounds (**9a**) and (**10**), respectively, two signals of C-H protons (from N-CH₂ groups) were observed. However, they exhibited the same one-bond and long-range proton carbon correlation. This effect was observed previously by Cox *et al.*⁹


The formation of **9a** and **9b** could be explained taking into account the well documented fact that alkenes react with sulphenyl halides *via* thiiranium intermediates.^{7,8} Therefore, the primarily formed azinyl cation should interact with 4-pentenoic acid to form thiiranium species (**T⁺**). (Scheme 4)

Scheme 3

Selected three-bond proton-carbon correlations deduced from HMBC spectra of **9a**, **9b** and **10**



Scheme 4



These species formed from 3-pyridine and 3- or 4-quinoline derivatives (**2**, **5**, **6b**, **6c**) would rearrange to oxolanones (**8**) (Scheme 4, route *a*). However, ring enlargement of thiiranium species (**T2**⁺) formed from α -azinylsulfenyl cations to thiazolium moiety in betaines (**9a**) or (**9b**) should start with less hindered site as nucleophilic displacement of methylene group by azine *endocyclic* nitrogen atom. Thus, as suggested by Kim⁵ for the formation of **9d**, thiiranium species (**T**⁺) formed from α -azinylsulfenyl cations underwent transformation to *anti*-Markovnikov products (**9a**) or (**9b**) containing *N*-methylene group (Scheme 4, route *b*). In the case of 8-quinolinyl isomer (**7**), the competition between the transformations of *S*-(8-quinolinyl)thiiranium salt to oxolanone (**8e**) and thiazinium betaine (**10**) was observed.

CONCLUSIONS

Two types of products, *i.e.* azinylthiooxolanones (**8**) and thiazolium betaines (**9**) or thiazinium betaine (**10**) were obtained after the electrolysis of bispyridinyl and bisquinolinyl disulfides (**1-7**) in the presence of 4-pentenoic acid. Their formation could be rationalized assuming the interaction of azinylsulfenyl cation with 4-pentenoic acid, which leads to the formation of azinylthiiranium species (**T**⁺). The latter should undergo final transformation either by nucleophilic cleavage of thiirane ring with carboxylic oxygen to form oxolanones (**8**) or by *endocyclic* nitrogen to give betaines (**9**) and (**10**).

EXPERIMENTAL

All melting points are uncorrected. All NMR spectra were recorded on a Bruker AVANS 400 spectrometer operating at 400.22 MHz and 100.64 MHz for ¹H and ¹³C nuclei, respectively, in deuteriochloroform or in hexadeuterodimethyl sulfoxide solutions with tetramethylsilane (δ 0.0 ppm) as internal standard. Two-dimensional ¹H-¹³C HSQC and HMBC experiments were performed using standard Bruker software HSQCGP and HMBCGP, respectively, and the following parameters: the spectral widths in *F*₂ and *F*₁ were *ca* 5 kHz for ¹H and 16.7 kHz for ¹³C, the relaxation delay was 1.5 s, the refocusing in the HSQC experiment was 1.7 ms and the delay for long-range evolutions was 50 ms in ¹H / ¹³C HMBC. 2D spectra were acquired as 2048 x 1024 hypercomplex files, with 1-4 transients. EIMS spectra were determined on a Finnigan MAT 95 spectrometer at 70 eV.

TLC analyses were performed employing Merck's silicagel 60 F₂₅₄ plates using a mixture of methylene chloride - ethanol (10:1, v/v) as an eluent.

2,2'- and 4,4'-Bispyridinyl disulfides (**1**) and (**3**) were commercial products. 3,3'-Bispyridinyl disulfide (**2**) and 3,3'-bisquinolinyl disulfide (**5**) were prepared as described previously.¹⁰ 4,4'-Bisquinolinyl disulfides (**6a**), (**6b**), (**6c**) were prepared by oxidation of the respective 4-quinolinethiones in alkaline milieu with potassium ferricyanide.¹¹ 8,8'-Bisquinolinyl disulfide (**7**) was prepared from 8-chlorosulfonylquinoline.¹²

General procedure for preparative electrolysis:

Electrolysis was carried out under controlled potential in three-compartment H-cell equipped with a platinum working electrode (area 10 cm²), a carbon rod as counter electrode and a saturated calomel electrode as a reference. Electrodes were connected to an Atlas Sollich 9833 potentiostat in combination with Atlas DC 9933 computer program. Values of oxidation potentials (determined in 0.1 M solution of tetraethylammonium perchlorate in acetonitrile) were used as working potentials for electrolysis.

A solution of disulfide (**1-7**) (1 mmol) in 100 mL of 0.1 M solution of tetrabutylammonium bromide in methylene chloride was electrolyzed at working potential as shown in Table. After electric current consumption of 10⁻⁴ F, 4-pentenoic acid (0.2 g, 2 mmol) in 2 mL of methylene chloride was added to the reaction mixture and the electrolysis was continued up to complete consumption of the starting disulfide (as monitored by TLC).

Isolation of electrolysis products:

i) The reaction mixture after the electrolysis of disulfides (**2**), (**5**), (**6b**) and (**6c**) was evaporated up to the volume of 30 mL, washed with water (3 x 30 mL), dried with anhydrous sodium sulfate. The solvent was stripped off. The residue was purified by column chromatography on silica gel 60 (Merck) using a mixture of methylene chloride - ethanol (10:1, v/v) as an eluent. The solid samples of **8c** and **8d** were recrystallized from acetone. Oxolanones (**8a**) and (**8b**) were obtained as thick oils.

ii) In the case of disulfide (**7**) the reaction mixture was filtered off to give 0.06 g (11 %) of thiazinium betaine (**10**), which was then recrystallized from methanol containing 3 drops of 48 % aqueous hydrogen bromide to give pure hydrogen bromide of salt (**10**). The filtrate was then worked up as above to give oxolanone (**8e**).

iii) The reaction mixture after the electrolysis of disulfides (**1**) or (**4**) was filtered off to give crude thiazolium derivatives (**9a**) or (**9b**), respectively. The sample of **9a** was recrystallized from methanol to afford crystals of semi bromine complex of **9a**. The samples of crude **9a** and **9b** were recrystallized from methanol containing 3 drops of 48 % aqueous hydrogen bromide to give pure hydrogen bromides of **9a** and **9b**.

ix) In the case of disulfides (**3**) and (**6a**) the consumption of electric current has stopped within 20 min. It may be due to the passivation of the working electrode (anode) by precipitation of a thin film of polymeric material on the surface of this electrode. The electrolysis was continued for additional 2 h. The mixture was then treated as above (procedure i) to afford only the non-consumed starting disulfide (**3**) or (**6a**), *ca.* 90 %.

5-(3-Pyridinylthio)methyloxolan-2-one (**8a**)

An oil. EIMS (70eV) (m/z): 209 (69, M⁺), 110 (100). ¹H NMR (CDCl₃), δ: 1.45-1.51 (m, 2H,

CHCH₂CH₂), 1.67-1.73 (m, 2H, CH CH₂CH₂), 3.33 (dd, 2H, $J=14.0$ Hz, $J=8.0$ Hz, SCH₂), 4.65-4.75 (m, 1H, CHO), 7.21 (dd, 1H, $J=7.8$ Hz, $J=4.8$ Hz, H5), 7.76 (ddd, 1H, $J=7.8$ Hz, $J=2.0$ Hz, $J=1.9$ Hz, H4), 8.44 (dd, 1H, $J=4.8$ Hz, $J=1.9$ Hz, H6), 8.61 (d, 1H, $J=2.0$ Hz, H2). *Anal.* Calcd for C₁₀H₁₁NO₂S: C 57.39; H 5.30; N 6.69. Found: C 57.12; H 5.23; N 6.47.

5-(3-Quinolinythio)methyloxolan-2-one (8b)

An oil. EIMS (70eV) (m/z): 259 (82, M⁺), 160 (100). ¹H NMR (CDCl₃), δ : 2.01-2.11 (m, 1H, CHCH₂CH₂), 2.39-2.48 (m, 1H, CHCH₂CH₂CO), 2.50-2.67 (m, 2H, CHCH₂CO), 3.19 (dd, 1H, $J=13.9$ Hz, $J=6.9$ Hz, SCH₂), 3.40 (dd, 1H, $J=13.9$ Hz, $J=5.2$ Hz, SCH₂), 4.64-4.71 (m, 1H, CHO), 7.57 (ddd, 1H, $J=8.1$ Hz, $J=7.0$ Hz, $J=1.1$ Hz, H6), 7.71 (ddd, 1H, $J=8.4$ Hz, $J=7.0$ Hz, $J=1.4$ Hz, H7), 7.77 (dd, 1H, $J=8.1$ Hz, $J=1.4$ Hz, H5), 8.08 (d, 1H, $J=8.4$ Hz, H8), 8.20 (d, 1H, $J=2.2$ Hz, H4), 8.89 (d, 1H, $J=2.2$ Hz, H2). *Anal.* Calcd for C₁₄H₁₃NO₂S: C 64.84; H 5.05; N 5.40. Found: 64.51; H 4.98; N 5.31.

5-(3-Methylthio-4-quinolinythio)methyloxolan-2-one (8c)

mp 81-82 °C (acetone). EIMS (70 eV) m/z : 305 (100, M⁺). ¹H NMR (CDCl₃), δ : 1.78-1.83 (m, 2H, CHCH₂CH₂), 2.15-2.19 (m, 2H, CH₂CH₂CO), 2.71 (s, 3H, SCH₃), 3.22 (d, $J=7.3$ Hz, 2H, SCH₂), 4.43-4.48 (m, 1H, CHO), 7.68-7.75 (m, 2H, 2 x H_{arom}), 8.02-8.05 (m, 1H, H_{arom}), 8.40-8.43 (m, 1H, H_{arom}), 8.86 (s, 1H, H2). *Anal.* Calcd for C₁₅H₁₅NO₂S₂: C 58.99; H 4.90; N 4.59; S 20.99. Found: C 58.78; H 4.80; N 4.65; S 20.23.

5-(7-Chloro-3-quinolinythio)methyloxolan-2-one (8d)

mp 93-94 °C (acetone). EIMS (70 eV) m/z : 293 (37, M⁺), 295 (13.5, M+2). 85 (100). ¹H NMR (CDCl₃), δ : 2.09-2.18 (m, 1H, CHCH₂CH₂), 2.45-2.54 (m, 1H, CHCH₂CH₂), 2.55-2.70 (m, 2H, CH₂CH₂CO) 3.34 (dd, 1H, $J=13.7$ Hz, $J=6.6$ Hz, SCH₂), 3.52 (dd, 1H, $J=13.7$ Hz, $J=5.4$ Hz, SCH₂), 4.78-4.84 (m, 1H, CHO), 7.28 (d, 1H, $J=4.8$ Hz, H3), 7.53 (dd, 1H, $J=9.1$ Hz, $J=2.1$ Hz, H6), 8.08 (d, 1H, $J=9.1$ Hz, H5), 8.09 (d, 1H, $J=2.1$ Hz, H8), 8.76 (d, 1H, $J=4.8$ Hz, H2). *Anal.* Calcd for C₁₄H₁₂NO₂ClS: C 57.24; H 4.12; N 4.77; S 10.92. Found: C 57.12; H 4.10; N 4.70; S 10.82.

5-(8-Quinolinythio)methyloxolan-2-one (8e)

An oil. EIMS (70eV) (m/z): 259 (20, M⁺), 161 (100). ¹H NMR (CDCl₃), δ : 2.45-2.65 (m, 4H, CHCH₂CH₂CO), 3.22-3.28 (m, 2H, SCH₂), 4.78-4.80 (m, 1H, CHO), 7.49 (dd, 1H, $J=8.0$ Hz, $J=4.4$ Hz, H3), 7.50 (dd, 1H, $J=8.0$ Hz, $J=7.0$ Hz, H6), 7.64 (dd, 1H, $J=7.0$ Hz, $J=1.2$ Hz, H7), 7.67 (dd, 1H, $J=8.0$ Hz, $J=1.2$ Hz, H5), 8.18 (dd, 1H, $J=8.0$ Hz, $J=1.6$ Hz, H4), 8.98 (dd, 1H, $J=4.4$ Hz, $J=1.6$ Hz, H2). *Anal.* Calcd for C₁₄H₁₃NO₂S: C 64.84; H 5.05; N 5.40. Found: C 64.59; H 4.99; N 5.27.

3-(2,3-Dihydrothiazolo[3,2-a]pyridinio-2-yl)propanoic acid bromide (9a) x HBr

mp 222-224 °C (methanol). ¹H NMR (DMSO) δ [δ_C for carbons from single bond and / long range proton-carbon correlations]: two independent signals of H3 protons were observed: 1.99 (m, 1H) and 2.08

(m, 1H), both with the same proton-carbon correlations [28.9 (C3) / 63.9 (C3'), 173.3 (C1)], 2.40 [(m, 2H, H2); 30.8 (C2) / 46.5 (C2')], 4.41 [(quintet, 1H, $J=7.6$ Hz, $J=7.6$ Hz, $J=7.5$ Hz, $J=4.9$ Hz, H2'); 46.5 (C2') / 30.8 (C2), 158.4 (C8a')], two independent signals of H3' protons were observed: 5.07 (dd, 1H, $J=13.6$ Hz, $J=4.9$ Hz) and 5.17 (dd, 1H, $J=13.6$ Hz, $J=7.5$ Hz) both with the same proton-carbon correlations [63.9 (C3') / 28.9 (C3), 143.0 [(C5'), 158.4 (C8a')], 7.76 [(ddd, 1H, $J=7.6$ Hz, $J=6.4$ Hz, $J=1.0$ Hz, H6'); 122.6 (C6') / 123.2 (C8')], 8.16 [(dd, 1H, $J=8.4$ Hz, $J=1.0$ Hz, H8'); 123.2 (C8') / 122.6 (C6')], 8.34 [(ddd, 1H, $J=8.4$ Hz, $J=7.6$ Hz, $J=1.0$ Hz, H7'); 144.6 (C7') / 143.0 (C5'), 158.4 (C8a')], 8.92 [(dd, 1H, $J=6.4$ Hz, $J=1.0$ Hz, H5'); 143.0 (C5') / 63.9 (C3'), 144.6 (C7'), 158.4 (C8a')], 12.34 [(s, 1H, COOH) [173.3 (C1)]. *Anal.* Calcd for C₁₀H₁₂NO₂BrS: C 41.39; H 4.17; N 4.83. Found: C 41.34; H 4.10; N 4.62.

3-(2,3-Dihydrothiazolo[3,2-a]quinolinio-2-yl)propanoic acid bromide (9b) x HBr

mp 198-200 °C (acetone). ¹H NMR (DMSO), δ : [δ_C for carbons from single bond and / long range proton-carbon correlations]: 2.16 [(m, 2H, H3); 29.7 (C3) / 60.9 (C3'), 173.4 (C1)], 2.48 [(m, 2H, H2); 30.8 (C2) / 45.6 (C2')], 4.59 (m, 1H, H2'); 45.6 (C2') / 30.8 (C2), 164.4 (C10a')], 5.39 (m, 2H, H3'); 60.9 (C3') / 29.7 (C3), 138.1 (C4a'), 164.4 (C10a')], 7.89 [(ddd, 1H, $J=8.0$ Hz, $J=7.2$ Hz, $J=1.0$ Hz H7'); 128.2 (C7') / 118.7 (C5'), 126.6 (C8a')], 8.15 [(ddd, 1H, $J=8.0$ Hz, $J=7.2$ Hz, $J=1.2$ Hz, H6'); 134.8 (C6') / 130.7 (C8'), 138.1 (C4a')], 8.19 [(d, 1H, $J=8.9$ Hz, H10'); 118.8 (C10') / 128.6 (C8a')], 8.24 [(dd, 1H, $J=8.4$ Hz, $J=1.0$ Hz, H5'); 118.7 (C5') / 126.6 (C8a'), 128.2 (C7')], 8.33 [(dd, 1H, $J=8.0$ Hz, $J=1.2$ Hz, H8'); 130.7 (C8') / 134.8 (C6'), 138.1 (C4a'), 147.3 (C9')], 8.93 [(d, 1H, $J=8.9$ Hz, H9'); 147.3 (C9') / 130.7 (C8'), 138.1 (C4a'), 164.4 (C10a')], 12.35 [(s, 1H, C1); 173.4 (C1)]. *Anal.* Calcd for C₁₄H₁₄NO₂BrS: C 49.42; H 4.15; N 4.12. Found: C 49.21; H 4.05; N 3.83.

3-(2,3-Dihydro-1,4-thiazine[2,3,4-*i,j*]quinolinio-2-yl)propanoate (10) ¹³

mp 242-244 °C (acetone). EIMS (70eV) (m/z): 259 (61, M⁺), 174 (100). ¹H NMR (DMSO), δ [δ_C for carbons from single bond and / long range proton-carbon correlations]: two independent signals of H3 protons were observed: 1.77-1.84 (m, 1H) and 2.03-2.10 (m, 1H) both with the same proton-carbon correlations [26.7 (C3) / 61.6 (C3'), 173.5 (C1)], ca. 2.50 [(m, 2H, H2); 30.7 (C2) / 35.7 (C2')], 3.98 [(m, 1H, H2'); 35.7 (C2') / 125.8 (C10a')], 5.10 [(dd, 1H, $J=14.3$ Hz, $J=7.4$ Hz, H3'); 61.6 (C3') / 35.7 (C2'), 133.0 (C10b'), 150.4 (C5')], 5.37 [(dd, 1H, $J=14.3$ Hz, $J=2.2$ Hz, H3'); 61.6 (C3') / 26.7 (C3), 133.0 (C10b'), 150.4 (C5')], 7.90 [(dd, 1H, $J=7.9$ Hz, $J=7.9$ Hz, H9'); 129.4 (C9') / 125.8 (C10a'), 130.5 (C7a')], 8.12 [(d, 1H, $J=7.9$ Hz, H10'); 132.7 (C10') / 127.0 (C8'), 133.0 (C10b')], 8.19 [(dd, 1H, $J=8.4$ Hz, $J=5.8$ Hz, H6'); 122.3 (C6') / 130.5 (C7a'), 150.4 (C5')], 8.21 [(d, 1H, $J=7.9$ Hz, H8'); 127.0 (C8') / 132.7 (C10'), 133.0 (C10b'), 148.7 (C7')], 9.29 [(d, 1H, $J=8.4$ Hz, H7'); 148.7 (C7') / 133.0 (C10b'), 150.4 (C5')], 9.41 [(d, 1H, $J=5.8$ Hz H5'); 150.4 (C5') / 61.6 (C3'), 122.3 (C6'), 133.0 (C10b'), 148.7 (C7')], and 173.5 (C1). *Anal.* Calcd for C₁₄H₁₃NO₂S: C 64.84; H 5.05; N 5.40; S 12.36. Found: C 64.75; H 4.99; N 5.24.

REFERENCES

Part LXXXVI in the Series of Azinyl Sulfides

1. S. Töteberg-Kaulen and E. Steckhan, *Tetrahedron*, 1988, **44**, 4389.
2. K. Marciniak, A. Maślankiewicz, and S. Boryczka, *Heterocycles*, 2002, **57**, 1279.
3. T. Tobien, H. Hungerbühler, and K-D. Asmus, *Phosphorus, Sulfur and Silicon*, 1994, **95-96**, 249.
4. a) L. M. Harwood, C. J. Moody, and J. M. Percy, *Experimental Organic Chemistry*, II ed., Blackwell Science, Oxford, 1999, p. 238. b) A. Vogel, *Textbook of practical organic chemistry including qualitative organic analysis*, IV ed., Longman, London and New York, 1978, p. 1075.
5. D. G. Kim, *Khim. Geterotsikl. Soedin.*, 1999, 334.
6. K. Undheim and R. Lie, *Acta Chem. Scand.* 1973, **27**, 19.
7. I. V. Koval, *Usp. Khim. (Russ. Chem. Rev.)*, 1995, **64**, 150.
8. I. V. Koval, *Usp. Khim. (Russ. Chem. Rev.)*, 1995, **64**, 781.
9. O. Cox, J. A. Prieto, L. Ramirez, M. Rodriguez, and J. R. Martinez, *J. Heterocycl. Chem.* 1999, **36**, 937.
10. A. Albert and G. B. Barlin, *J. Chem. Soc.*, 1959, 2384.
11. M. Konishi, N. Fukuda, Y. Oku, H. Yamazaki, K. Imazumi, and H. Kobayashi, Japan PCT Int. Appl. WO 96 23, 774 (1995) (*Chem. Abstr.*, 1996, **125**, 2476331h).
12. Yu. A. Bankovskii, A. F. Yevinsh, and E. A. Luksha, *Zh. Obshch. Khim.*, 1958, **28**, 2273.
13. The Autonom name: 2,3-dihydro-1-thia-3a-azoniaphenalene was used for this heterocyclic system as well (*Chem. Abstr.*, 1998, **128**, 294757a, for the paper of D. G. Kim, *Khim. Geterotsikl. Soedin.*, 1997, 1133).