

HETEROCYCLES, Vol. 65, No. 11, 2005, pp. 2771 - 2776

Received, 25th July, 2005, Accepted, 26th August, 2005, Published online, 26th August, 2005

NEW VERMISTATIN DERIVATIVES ISOLATED FROM *PENICILLIUM SIMPLICISSIMUM*

Shin-ichirou Komai,^a Tomoo Hosoe,^a Takeshi Itabashi,^a Koohei Nozawa,^b
Takashi Yaguchi,^c Kazutaka Fukushima,^c and Ken-ichi Kawai^{a,*}

^aFaculty of Pharmaceutical Sciences, Hoshi University, Ebara 2-4-41, Shinagawa-ku, Tokyo 142-8501, Japan. e-mail: kawai@hoshi.ac.jp ^bSchool of Pharmaceutical Sciences, Ohu University, 31-1 Misumido, Tomita-machi, Koriyama, Fukushima 963-8611, Japan. ^cResearch Center for Pathogenic Fungi and Microbial Toxicoses, Chiba University, Inohana 1-8-1, Chuo-ku, Chiba 260-8673, Japan

Abstract – Four new vermistatin derivatives, dihydrovermistatin (**1**), acetoxyl-dihydrovermistatin (**2**), hydroxydihydrovermistatin (**3**), and penisimplicissin (**4**) were isolated along with vermistatin (**5**), penicillide, and funicone from the extract of *Penicillium simplicissimum* IFM 53375. The structures of **1** – **4** were determined by spectroscopic and chemical methods.

INTRODUCTION

During our search for new antifungal compounds against pathogenic filamentous fungi, *Aspergillus fumigatus* and *A. niger*, and/or pathogenic yeasts, *Candida albicans* and *Cryptococcus neoformans* from fungal sources,¹ we found that the extract of *Penicillium simplicissimum* IFM 53375 showed characteristic antifungal activity against *A. fumigatus*. Recently new meroterpenoids, penisimplicins A and B, were isolated from the above extract.² Further fractionation of the extract led to the isolation of four new vermistatin derivatives designed as dihydrovermistatin (**1**), acetoxyl-dihydrovermistatin (**2**), hydroxydihydrovermistatin (**3**), and penisimplicissin (**4**) along with penicillide³, vermistatin (**5**)⁴, and funicone.⁵ In this paper, we report the structural determination of four new vermistatin derivatives (**1** – **4**).

RESULTS AND DISCUSSION

The molecular formulae of **1**– **4** were determined by HREI-MS spectrum as C₁₈H₁₈O₆, C₂₀H₂₀O₈, C₁₈H₁₈O₇ and C₁₆H₁₄O₆, respectively. The UV spectra of **1**–**4** were almost superimposable with those of vermistatin (**5**).⁴ The ¹H-NMR spectra of **1**–**4** (Table 1) showed commonly two methoxy signals, a methine proton bearing a hydroxyl group, two aromatic or olefinic protons, and a pair of *meta*-coupled

doublet aromatic protons as same as **5**, except for the side chain. The above results indicated that **1-4** were the derivative of vermistatin (**5**).

Table 1. ^1H - and ^{13}C -NMR spectral data for **1-5** in CDCl_3 (δ in ppm)

No	1		2		3		4		5	
	^{13}C	^1H	^{13}C	^1H	^{13}C	^1H	^{13}C	^1H	^{13}C	^1H
1	170.0		170.0		170.2		170.0		170.0	
2	129.3		129.3		129.4		129.3		129.3	
3	99.0	6.98 d(1.9)	99.0	6.98 d(2.0)	99.0	6.97 d(2.0)	99.0	6.98 d(1.8)	99.0	6.99 d
4	163.1		163.1		163.1		163.1		163.1	
5	105.1	6.69 d(1.9)	105.2	6.69 d(2.0)	105.0	6.66 d(2.0)	105.1	6.68 d(1.8)	105.1	6.68 d
6	154.9		154.9		154.8		154.8		154.7	
7	127.7		127.5		127.5		127.7		127.6	
8	73.5	6.46 s	73.4	6.44 (s)	74.2	6.32 s	73.5	6.45 s	73.5	6.46 s
9	123.4		123.7		122.8		123.4		123.6	
10	177.1		176.7		177.3		176.9		177.0	
11	114.4	6.20 s	116.3	6.22 (s)	116.0	6.21 s	115.1	6.19 s	112.9	6.16 s
12	169.4		165.1		167.2		165.9		162.1	
14	154.5	7.44 s	154.6	7.45 (s)	155.3	7.64 s	154.4	7.42 s	153.8	7.43 s
15	35.3	2.48 t(7.6)	39.9	2.70 dd(14.6,5.2) 2.77 dd(14.6,7.3)	43.3	2.57 dd(14.6,8.2) 2.64 dd(14.6,4.0)	19.7	2.26 s	123.1	6.07 d
16	19.9	1.67 qt(7.4,7.6)	67.4	5.19 qdd(6.4,7.3,5.2)	65.3	4.18 qdd(6.4,8.2,4.0)			135.9	6.60 dq
17	13.4	0.98 t(7.4)	19.8	1.30 d(6.4)	23.4	1.28 d(6.4)			18.5	1.92 d
4-OCH ₃	56.0	3.88 s	56.0	3.88 s	56.0	3.87 s	56.0	3.88 s	56.0	3.88 s
6-OCH ₃	55.8	3.79 s	55.8	3.79 s	55.8	3.78 s	55.8	3.79 s	55.8	3.79 s
COCH ₃			21.1	2.02 s						
$\underline{\text{COCH}}_3$			170.2							

The side chain of **1**, **2**, **3**, and **4** were confirmed as propyl, 2-acetyloxypropyl, 2-hydroxypropyl, and methyl groups, respectively, from the analysis of ^1H -NMR spectra. The ^1H - and ^{13}C -NMR spectral assignments of **1-4** (Table 1) were determined from the detailed analyses of the ^1H - ^1H COSY, HMQC, and HMBC spectra. Hydrogenation of vermistatin (**5**) using H_2 gas under the presence of 5% Pd-C led to dihydrovermistatin (**1**), whereas acetylation of hydroxydihydrovermistatin (**3**) gave acetoxydihydrovermistatin (**2**). From the CD spectra of **1-4** compared with that of vermistatin (**5**), the stereochemistry at C-8 in **1-4** was confirmed as the same as that of vermistatin (**5**).

In order to confirm the absolute configuration of the secondary alcohol in the side chain of **2** and **3**, the advanced Mosher's method⁶ was applied to **3**. The (*R*)- and (*S*)- α -methoxy- α -trifluoromethyl-phenylacetic acid (MTPA) esters of **3** were synthesized and the values of the chemical shift differences between the (*S*)- and (*R*)-MTPA esters [$\Delta\delta = \delta_{\text{S}} - \delta_{\text{R}}$ in Hz (500 MHz)] were calculated. From the results (Figure 1), the secondary hydroxyl group in **3** was assigned as *R*-configuration. Therefore, the absolute configuration of the acetoxyl group in **2** was determined as *R*-configuration. The absolute structures of four vermistatin derivatives were consequently confirmed as depicted in **1-4**.





Figure 1. Differences of chemical shifts ($\Delta\delta$ in Hz) between the (*S*)- and (*R*)-MTPA esters of **3**

Vermistatin (**5**) was originally isolated as the cytotoxic compound for tumor cells from *Penicillium vermiculatum*,⁷ which is the synonym for an anamorph of *Talaromyces flavus*.⁸ Vermistatin (**5**) was also isolated as fijiensin, phytotoxic on various banana cultivars, from *Mycosphaerella fijiensis*, the causal agent of Black Sigatoka disease of bananas.⁹ Recently Arai *et al.* reported⁸ the isolation of vermistatin (**5**) along with funicone and its derivatives from *T. flavus*. On the taxonomy, the teleomorphic states of *Penicillium* are broadly divided into two genera, *Talaromyces* and *Eupenicillium*. The place of *P. simplicissimum*, which is a member of the group including *Eupenicillium*, is phylogenetically different from that of *P. vermiculatum* in the genus *Penicillium*. From this point of view, it is taxonomically interesting that *P. simplicissimum* produced vermistatin (**5**) and its derivatives (**1-4**) along with funicone. Since **1 – 5** showed almost no antifungal activity, further investigation for the antifungal substances in *P. simplicissimum* will be attempted.

EXPERIMENTAL

General Experimental Procedures. General experimental procedures were described in the previous paper.¹

Isolation of the metabolite from *P. simplicissimum*. *P. simplicissimum* IFM 53375 was cultured at 25 °C for 21 days in 16 Roux flasks containing 150 g of moist rice in each flask. The fermented rice (2.5 kg) was extracted with CH_2Cl_2 -MeOH (1:1) (8 L) at rt for 16 h and the organic layer was evaporated *in vacuo*. The resultant extract (56 g) was suspended in H_2O and extracted with EtOAc, and then the organic layer was evaporated *in vacuo*, respectively. The EtOAc extract (40 g), which showed the strong antifungal activity against *A. fumigatus*, was separated by column chromatography on silica gel (600 g) into six fractions: CH_2Cl_2 (50:1) (8 g), CH_2Cl_2 -EtOH (20:1) (14 g), CH_2Cl_2 -EtOH (10:1) (4 g), CH_2Cl_2 -EtOH (5:1) (4 g), CH_2Cl_2 -EtOH (1:1) (5 g), and EtOH (1.5 g). The 2nd fraction [CH_2Cl_2 -EtOH (20:1)] was purified by LPLC on a silica gel column using benzene-acetone (15:1) followed by the further

purification of HPLC on silica gel column [CH_2Cl_2 -acetone (15:1)] to give penicillide³ (35 mg), dihydrovermistatin (**1**) (20mg), vermistatin⁴ (**5**) (1.0 g), funicone⁵ (8 mg), penisimplicissin (**4**) (15 mg), and acetoxydihydrovermistatin (**2**) (20 mg). The 3rd fraction [CH_2Cl_2 -EtOH (10:1)] was also purified with LPLC on a silica gel column using CH_2Cl_2 -EtOH (20:1), followed by the purification with HPLC on a silica gel column using CH_2Cl_2 -acetone (2:1) to give hydroxydihydrovermistatin (**3**) (32 mg). Penicillide, funicone, and **5** were identified by the comparison with the published data.³⁻⁵

Dihydrovermistatin (**1**) : Colorless needles (from MeOH); mp 143-145°C; $[\alpha]_{\text{D}}^{19}$ -108° ($c = 0.10$, CHCl_3); IR (KBr) ν_{max} : 1770(CO_2), 1665 (CO) cm^{-1} ; UV (MeOH) λ_{max} (log ϵ): 210 (4.6), 245 (4.1), 303 (3.6); EI-MS m/z : 330.1094 (M^+ , 330.1103 for $\text{C}_{18}\text{H}_{18}\text{O}_6$, 90), 287 (17), 217 (100); CD($\Delta\epsilon$): 210 (+13.9), 222 (-16.4), 250 (+2.2), 303 (-1.0) nm. The assignments of ^1H - and ^{13}C -NMR signals are summarized in Table 1.

Acetoxydihydrovermistatin (**2**) : White amorphous powder; $[\alpha]_{\text{D}}^{19}$ -80° ($c = 0.20$, CHCl_3); IR (KBr) ν_{max} : 1770(CO_2), 1660 (CO) cm^{-1} ; UV (MeOH) λ_{max} (log ϵ): 210 (4.7), 246 (4.2), 305 (3.7); EI-MS m/z : 388.1160 (M^+ , 388.1158 for $\text{C}_{20}\text{H}_{20}\text{O}_8$, 63), 328 (100), 217 (33); CD($\Delta\epsilon$): 212 (+18.6), 222 (-42.5), 250 (+3.6), 303 (-3.8) nm. The assignments of ^1H - and ^{13}C -NMR signals are summarized in Table 1.

Hydroxydihydrovermistatin (**3**) : Colorless needles (from MeOH); mp 184-185°C; $[\alpha]_{\text{D}}^{19}$ -85° ($c = 0.13$, CHCl_3); IR (KBr) ν_{max} : 3380(OH), 1770(CO_2), 1660 (CO) cm^{-1} ; UV (MeOH) λ_{max} (log ϵ): 210 (4.6), 248 (4.2), 306 (3.6) nm; EI-MS m/z : 346.1064 (M^+ , 346.1053 for $\text{C}_{18}\text{H}_{18}\text{O}_7$, 100), 287 (23), 217 (42); CD($\Delta\epsilon$): 210 (+21.1), 222 (-22.6), 249 (+3.16), 301 (-0.8), 325 (+0.5) nm. The assignments of ^1H - and ^{13}C -NMR signals are summarized in Table 1.

Penisimplicissin (**4**) : Colorless needles (from acetone); mp 185-186°C; $[\alpha]_{\text{D}}^{19}$ -119° ($c = 0.10$, CHCl_3); IR (KBr) ν_{max} : 1770(CO_2), 1665 (CO) cm^{-1} ; UV (MeOH) λ_{max} (log ϵ): 210 (4.6), 247 (4.1), 305 (3.5); EI-MS m/z : 302.0793 (M^+ , 302.0790 for $\text{C}_{16}\text{H}_{14}\text{O}_6$, 100), 287 (M-Me, 10), 259 (46), 217 (38); CD($\Delta\epsilon$): 210 (+15.4), 222 (-21.2), 250 (+2.4), 303 (-1.5) nm. The assignments of ^1H - and ^{13}C -NMR signals are summarized in Table 1.

Synthesis of (S)- and (R)-MTPA esters of hydroxydihydrovermistatin (**3**).

Dicyclohexylcarbodiimide (16 mg), 4-dimethylaminopyridine (6 mg), and (S)- or (R)-MTPA (16 mg) were added to a solution of hydroxydihydrovermistatin (**3**) (3.5 mg) in CH_2Cl_2 (1 mL). The reaction mixture was kept at 40°C for 1.5 h, and then washed with 0.5M HCl, saturated NaHCO_3 , and water successively, and dried over Na_2SO_4 . After removal of the solvent by evaporation, the residue was purified by HPLC (silica gel) (CH_2Cl_2 -acetone 15:1) to afford the (R)- or (S)-MTPA ester of **3** [1.1 mg for (S), 2.6mg for (R)].

(S)-MTPA ester of **3**: White amorphous powder. ^1H -NMR (CDCl_3 , 270 MHz; except for the phenyl signals): δ 7.30 (1H, s, H-14), 6.99 (1H, d, $J = 1.8$ Hz, H-3), 6.69 (1H, d, $J = 1.8$ Hz, H-5), 6.47 (1H, s,

H-8), 6.06 (1H, s, H-11), 5.50 (1H, qt, $J=6.4, 6.4$ Hz, H-16), 3.89 (3H, s, 4-OCH₃), 3.75 (3H, s, 6-OCH₃), 2.76 (2H, d, $J=6.4$ Hz, H-15), 1.44 (3H, d, $J=6.4$ Hz, H-17).

(*R*)-MTPA ester of **3**: White amorphous powder. ¹H-NMR (CDCl₃, 270 MHz; except for the phenyl signals): δ 7.39 (1H, s, H-14), 6.99 (1H, d, $J=1.8$ Hz, H-3), 6.67 (1H, d, $J=1.8$ Hz, H-5), 6.45 (1H, s, H-8), 6.19 (1H, s, H-11), 5.46 (1H, qt, $J=6.4, 6.4$ Hz, H-16), 3.89 (3H, s, 4-OCH₃), 3.72 (3H, s, 6-OCH₃), 2.81 (2H, d, $J=6.4$ Hz, H-15), 1.37 (3H, d, $J=6.4$ Hz, H-17).

Acetylation of hydroxydihydrovermistatin (3). A mixture of **3** (5 mg), acetic anhydride (0.6 mL), and pyridine (0.1 mL) was stirred at rt for 16 h. Work-up in the usual manner provided a viscous oil, which was purified by HPLC on silica gel with CH₂Cl₂-acetone (10:1) to give an acetate (4 mg). This acetate was identical to acetoxidyhydrovermistatin (**2**) based on the spectroscopic data including the optical rotation and CD curve.

Hydrogenation of vermistatin (5). 5% Pd-C (20 mg) was suspended in a solution of vermistatin (**5**) (20 mg) in MeOH (4 mL) and the mixture stirred at rt in a hydrogen atmosphere for 10 min. After removal of the catalysts, the solvent was evaporated in vacuo. The residue was purified by HPLC on silica gel with CH₂Cl₂-acetone (15:1) to give a dihydro derivative (16 mg). This compound was identical to dihydrovermistatin (**1**) based on the spectroscopic data including the optical rotation and CD curve.

ACKNOWLEDGEMENTS

We are grateful to Dr. H. Kasai, and Miss N. Kobayashi of Hoshi University for NMR and MS spectral measurements. This study was supported in part by Hoshi University Science/Technology Frontier Research Base from the Ministry of Education, Culture, Sports, Science and Technology, Japan and by a Cooperative Research Program of Research Center for Pathogenic Fungi and Microbial Toxicoses, Chiba University (2004-22).

REFERENCES

1. S. Komai, T. Hosoe, K. Nozawa, K. Okada, G. M. de Campos Takaki, K. Fukushima, M. Miyaji, Y. Horie, and K. Kawai, *Mycotoxins*, 2003, **53**, 11.
2. S. Komai, T. Hosoe, T. Itabashi, K. Nozawa, K. Okada, G. M. de Campos Takaki, T. Yaguchi, K. Fukushima, M. Miyaji, Y. Horie, and K. Kawai, *Chem. Pharm. Bull.*, 2005, in press.
3. T. Sassa, G. Niwa, H. Unno, and Y. Miura, *Tetrahedron Lett.*, 1974, 3941.
4. J. Fuska, D. Uhrin, B. Proksa, Z. Voticky, and J. Ruppeltdt, *J. Antibiotics*, 1986, **39**, 1605.
5. L Merlini, G. Nasini, and A. Selva, *Tetrahedron*, 1970, **26**, 2739.
6. I. Ohtani, T. Kusumi, Y. Kashman, and H. Kakisawa, *J. Am. Chem. Soc.*, 1991, **113**, 4092.
7. J. Fuska, A. Fuskova, and P. Nemec, *J. Antibiotics*, 1979, **32**, 667.
8. J. I. Pitt, "The Genus *Penicillium*", Academic Press, Inc., London, 1979.

9. R. K. Upadhyay, G. A. Strobel, S. J. Coval, and J. Clardy, *Experientia*, 1990, **46**, 982.
10. M. Arai, H. Tomoda, T. Okuda, H. Wang, N. Tabata, R. Masuma, Y. Yamaguchi, and S. Omura, *J. Antibiotics*, 2002, **55**, 172.