

HETEROCYCLES, Vol. 71, No. 7, 2007, pp. 1509 - 1516. © The Japan Institute of Heterocyclic Chemistry
Received, 2nd January, 2007, Accepted, 10th May, 2007, Published online, 11th May, 2007. COM-07-10988

MICROWAVE MEDIATED SYNTHESIS OF 3,4,4a,11a-TETRAHYDRO-5-PHENYL/5-(4'-SUBSTITUTEDPHENYL)-2H-INDENO[1,2-*b*]PYRANO-[3,2-*e*]PYRAN-6(5H)-ONES

Som N. Dhawan, Mamta Sharma, Sunita Bajaj, Kusum Sharma, and Somesh Sharma*

Department of Chemistry, Kurukshetra University, Kurukshetra –136119 Haryana, India. Email: sharmassomesh@yahoo.com

Abstract — Synthesis of 3,4,4a,11a-tetrahydro-5-phenyl/5-(4'-substituted phenyl)-2H-indeno[1,2-*b*]pyrano[3,2-*e*]pyran-6(5H)-ones have been reported under microwave conditions. The substrates for this being 2-benzylideneindane-1,3-dione (**1**) and 3,4-dihydro-2H-pyran (**2**); the product formation is governed by inverse Diels-Alder reaction conditions.


INTRODUCTION

Both inter- and intramolecular¹⁻⁷ Diels-Alder reactions have been extensively used in the synthesis of multicyclic ring systems primarily due to their ability to generate simultaneously upto four chiral centres in a highly stereoselective and largely predictable manner. In these reactions, α,β -unsaturated carbonyl compounds have been used both as dienes and as dienophiles depending upon the magnitude of electron density in the other component. In conventional Diels-Alder reactions, α,β -unsaturated compounds act as the dienophile while in inverse electron demand Diels-Alder reactions⁸⁻⁹ these act as dienes. In view of our continuous interest in the synthesis of novel multicyclic heterocyclic ring systems, we describe in this paper, the synthesis and spectral studies of hitherto unknown 2H-indeno[1,2-*b*]pyrano[3,2-*e*]pyran-6(5H)-one ring system using inverse electron demand Diels-Alder reactions between 2-benzylideneindane-1,3-dione (**1**) and 3,4-dihydro-2H-pyran (**2**) under microwave conditions.

RESULTS AND DISCUSSION

The reaction was carried out between 2-benzylideneindane-1,3-dione (**1**) and 3,4-dihydro-2H-pyran (**2**) in toluene for 10-20 min at 80 °C under microwave conditions. The chromatographic separation of the resulting reaction mixture afforded two isomeric cycloadducts (**3**) and (**4**) in good yields (Scheme 1). The

reaction was also carried out in the conventional way in acetonitrile, but the yield of the desired products was poor (Table 1). The structures of these cycloadducts have been ascertained through their IR, ^1H NMR and mass spectra.



Scheme 1

Table 1. Comparison of Inverse Diels-Alder reaction results under thermal and microwave condition.


R	Time (min)/MW	Yield (%)		Time (h)/Δ	Yield *(%)	
	Solvent (toluene/MeCN)	3	4	Solvent (toluene/ MeCN)	3	4
a = H	20/180)	45	47	48/5	17.6	14.6
b = CH ₃	45 / 180	0	0	72/10	0	0
c = OCH ₃	60 /180	0	0	72/10	0	0
d = F	15 /120	45	48	24/3	15.2	5.6
e = CO ₂ CH ₃	10 /150	48	43	24/3.5	14.9	6.2
f = NO ₂	10 /180	50	41	36/4.0	16.7	8.2

The regiochemistry and stereochemistry (*cis*) of cycloadducts could be explained on the basis of Woodward-Hoffmann rules,¹⁰ further supported by ^1H -NMR chemical shifts of the protons and their

coupling constants. In cycloadduct (**3f**), the protons $C_{4a}\text{-H}$, $C_{11a}\text{-H}$ and $C_5\text{-H}$ appeared in the regions δ 2.22-2.30, δ 5.75-5.77 ($J=2.4\text{Hz}$) and δ 4.28-4.30 ($J=6.0\text{Hz}$) respectively. In cycloadduct (**4f**), $C_{4a}\text{-H}$ and $C_{11a}\text{-H}$ appeared approximately in the same region except $C_5\text{-H}$ that appeared at δ 3.75 as a singlet (little broadened at the base presumably due to its low coupling constant). The dihedral angle between the planes defined by the atoms $4a\text{-H}$, $C_{4a}\text{-H}$ and C_5 ; and C_{4a} , C_5 and 5-H in cycloadduct **3f** is 42.8° as calculated using Tripos Alchemy III computer programme; and thus $J_{4a,5}$ is expected to have a value between 4-6 Hz while the dihedral angle between the same planes in cycloadduct **4f** is 84.6° and thus $J_{4a,5}$ is expected to have a value between 0-1Hz.

These assignments for chemical shifts of $C_5\text{-H}$, $C_{4a}\text{-H}$ and $C_{11a}\text{-H}$ and coupling constants ($J_{4a,5}$) are in excellent agreement with those reported for cycloadducts^{11,12} and further corroborated by HSQC (Figure 2) and the decoupling experiments of cycloadduct (**4f**). Upon double irradiation of $C_{4a}\text{-H}$ in cycloadduct (**3f**, $R=\text{NO}_2$), the doublet in the region δ 4.28 – 4.30 due to $C_5\text{-H}$ ($J=6.0\text{Hz}$) and doublet at δ 5.75-5.77 due to $C_{11a}\text{-H}$ ($J=2.4\text{ Hz}$) collapsed to sharp singlets. Similarly, when $C_{4a}\text{-H}$ in cycloadduct (**4f**, $R=\text{NO}_2$) was double irradiated, not only was the broadened singlet at δ 3.75 due to $C_5\text{-H}$ was reduced to a sharp singlet, but even the doublet in the region δ 5.45-5.46 ($J=2.6\text{ Hz}$) due to $C_{11a}\text{-H}$, also changed to a sharp singlet. These two isomeric cycloadducts differ in stereochemistry at $C_5\text{-phenyl}$ / substituted phenyl group. The stereochemical relationship between these three protons, viz., $C_{4a}\text{-H}$, $C_5\text{-H}$ and $C_{11a}\text{-H}$ in cycloadducts **3a** (*cis*) and **4a** (*trans*) have been confirmed through their molecular models (Figure 1) generated in their respective low energy orientation using Tripos Alchemy III computer programme.

Figure 1. Molecular models of cycloadducts of **3a** and **4a**



Current Data Parameters

NAME	12
EXPNO	1
PROCNO	1
F2 - Acquisition Parameters	
Date_	20070707
Time	17.35
INSTRUM	zgpg30
PROBHD	5 mm QNP1H/13
PULPROG	zgpg30
VS	132.6
RGZVARY	QCZ3
NUC1	13C
NUC2	1H
PCPD	3.216480 Hz
PCPD2	3.216480 Hz
PCPD3	3.216480 Hz
PCPD4	3.216480 Hz
PCPD5	3.216480 Hz
PCPD6	3.216480 Hz
PCPD7	3.216480 Hz
PCPD8	3.216480 Hz
PCPD9	3.216480 Hz
PCPD10	3.216480 Hz
PCPD11	3.216480 Hz
PCPD12	3.216480 Hz
PCPD13	3.216480 Hz
PCPD14	3.216480 Hz
PCPD15	3.216480 Hz
PCPD16	3.216480 Hz
PCPD17	3.216480 Hz
PCPD18	3.216480 Hz
PCPD19	3.216480 Hz
PCPD20	3.216480 Hz
PCPD21	3.216480 Hz
PCPD22	3.216480 Hz
PCPD23	3.216480 Hz
PCPD24	3.216480 Hz
PCPD25	3.216480 Hz
PCPD26	3.216480 Hz
PCPD27	3.216480 Hz
PCPD28	3.216480 Hz
PCPD29	3.216480 Hz
PCPD30	3.216480 Hz
PCPD31	3.216480 Hz
PCPD32	3.216480 Hz
PCPD33	3.216480 Hz
PCPD34	3.216480 Hz
PCPD35	3.216480 Hz
PCPD36	3.216480 Hz
PCPD37	3.216480 Hz
PCPD38	3.216480 Hz
PCPD39	3.216480 Hz
PCPD40	3.216480 Hz
PCPD41	3.216480 Hz
PCPD42	3.216480 Hz
PCPD43	3.216480 Hz
PCPD44	3.216480 Hz
PCPD45	3.216480 Hz
PCPD46	3.216480 Hz
PCPD47	3.216480 Hz
PCPD48	3.216480 Hz
PCPD49	3.216480 Hz
PCPD50	3.216480 Hz
PCPD51	3.216480 Hz
PCPD52	3.216480 Hz
PCPD53	3.216480 Hz
PCPD54	3.216480 Hz
PCPD55	3.216480 Hz
PCPD56	3.216480 Hz
PCPD57	3.216480 Hz
PCPD58	3.216480 Hz
PCPD59	3.216480 Hz
PCPD60	3.216480 Hz
PCPD61	3.216480 Hz
PCPD62	3.216480 Hz
PCPD63	3.216480 Hz
PCPD64	3.216480 Hz
PCPD65	3.216480 Hz
PCPD66	3.216480 Hz
PCPD67	3.216480 Hz
PCPD68	3.216480 Hz
PCPD69	3.216480 Hz
PCPD70	3.216480 Hz
PCPD71	3.216480 Hz
PCPD72	3.216480 Hz
PCPD73	3.216480 Hz
PCPD74	3.216480 Hz
PCPD75	3.216480 Hz
PCPD76	3.216480 Hz
PCPD77	3.216480 Hz
PCPD78	3.216480 Hz
PCPD79	3.216480 Hz
PCPD80	3.216480 Hz
PCPD81	3.216480 Hz
PCPD82	3.216480 Hz
PCPD83	3.216480 Hz
PCPD84	3.216480 Hz
PCPD85	3.216480 Hz
PCPD86	3.216480 Hz
PCPD87	3.216480 Hz
PCPD88	3.216480 Hz
PCPD89	3.216480 Hz
PCPD90	3.216480 Hz
PCPD91	3.216480 Hz
PCPD92	3.216480 Hz
PCPD93	3.216480 Hz
PCPD94	3.216480 Hz
PCPD95	3.216480 Hz
PCPD96	3.216480 Hz
PCPD97	3.216480 Hz
PCPD98	3.216480 Hz
PCPD99	3.216480 Hz
PCPD100	3.216480 Hz

***** CHANNEL F1 *****

NUC1	13C
PCPD	3.216480 Hz
PCPD2	3.216480 Hz
PCPD3	3.216480 Hz
PCPD4	3.216480 Hz
PCPD5	3.216480 Hz
PCPD6	3.216480 Hz
PCPD7	3.216480 Hz
PCPD8	3.216480 Hz
PCPD9	3.216480 Hz
PCPD10	3.216480 Hz
PCPD11	3.216480 Hz
PCPD12	3.216480 Hz
PCPD13	3.216480 Hz
PCPD14	3.216480 Hz
PCPD15	3.216480 Hz
PCPD16	3.216480 Hz
PCPD17	3.216480 Hz
PCPD18	3.216480 Hz
PCPD19	3.216480 Hz
PCPD20	3.216480 Hz
PCPD21	3.216480 Hz
PCPD22	3.216480 Hz
PCPD23	3.216480 Hz
PCPD24	3.216480 Hz
PCPD25	3.216480 Hz
PCPD26	3.216480 Hz
PCPD27	3.216480 Hz
PCPD28	3.216480 Hz
PCPD29	3.216480 Hz

Further, the effects of electron donating and withdrawing substituents have been explored on product formation. In case of electron donating groups, no formation of cycloadduct has been observed as it increases the energy gap between LUMO (diene) and HOMO (dienophile). The reaction was also tried in MeCN to obtain the comparative results with the thermal condition.

From the above discussion, we conclude that incorporation of electron-withdrawing groups at *p*-position of benzylidene moiety of 2-benzylideneindane-1,3-diones favours the cycloaddition between 1,3-diones and 3,4-dihydro-2*H*-pyran while incorporation of electron donating groups such as methyl and methoxy inhibit the cycloaddition reaction. Further, the formation of cycloadducts is controlled by the approach of 3,4-dihydro-2*H*-pyran from the less hindered site.

EXPERIMENTAL

Melting points were determined in open capillaries using a sulfuric acid bath and are uncorrected. IR spectra were recorded in Nujol mulls on a Perkin-Elmer 842 spectrophotometer and ^1H -NMR spectra were recorded on 90 MHz Perkin-Elmer R-32 instrument and ^{13}C -NMR were recorded on Bruker AV-400 (Top spin) in CDCl_3 using Me_4Si as internal standard. Low resolution mass spectra were recorded at 70eV using a VG-70S instrument. Elemental analysis was performed on Heraeus CHN-Rapid analyzer. Microwave reactions were carried out in Biotage initiator with power range of 0-400 W from magnetron at 2.45GHz.

General procedure for cycloaddition between 2-benzylideneindane-1,3-diones **1** and 3,4-dihydro-2H-pyran **2**.

Microwave Method: A solution of 2-benzylideneindane-1,3-dione (**1**, 1 g, 4.3 mmol) and 3,4-dihydro-2H-pyran (**2**, 0.36 g, 4.4 mmol) in toluene (2 mL) was subjected to microwave irradiations for 10-20min in 2-5 mL vial in Biotage initiator with 'high' absorption level, fixed hold time 'on' and pre-stirring for 30 sec adjusted in the instrument. The reaction mass was concentrated under vacuum and the crude product purified by column chromatography (silica gel, 100-200 mesh; eluent hexane: benzene).

Conventional Method: A solution of 2-benzylideneindane-1,3-dione (**1**, 1 g, 4.3 mmol) and 3, 4-dihydro-2H-pyran (**2**, 0.34 g, 4 mmol) in anhydrous acetonitrile (80 mL) was refluxed for 4.5 h. The progress of the reaction monitored by TLC. The reaction mass was concentrated under vacuum and the crude product purified by column chromatography (silica gel, 100-200 mesh; eluent hexane: benzene) to obtain (**3a**) as yellow solid, which upon crystallization from benzene-pet. Ether mixture afforded bright yellow crystals. Subsequent elution's afforded another adduct (**4a**); crystallized from benzene-pet. Ether mixture to yield yellow crystals.

3a: (4aR*,5R*,11aS*) 3,4,4a,11a-Tetrahydro-5-phenyl-2H-indeno[1,2-b]pyrano[3,2-e]pyran-6(5H)-one. mp 165-166 °C; IR (KBr) 1695, 1630 cm^{-1} ; ^1H -NMR (90MHz, CDCl_3): δ 1.33-1.48 (m, 2H, $\text{C}_4\text{-H}$), 1.54-1.75 (m, 2H, $\text{C}_3\text{-H}$), 2.17-2.25 (m, 1H, $\text{C}_{4a}\text{-H}$), 3.82-3.86 (m, 1H, $\text{C}_2\text{-H}$), 3.99-4.08 (m, 1H, $\text{C}_2\text{-H}$), 4.17-4.19 (d, $J = 6.0$ Hz, 1H, $\text{C}_5\text{-H}$), 5.72-5.75 (d, $J=2.6\text{Hz}$, 1H, $\text{C}_{11a}\text{-H}$), 7.08-7.40 (m, 9H, $\text{C}_2'\text{-H}$, $\text{C}_3'\text{-H}$, $\text{C}_4'\text{-H}$, $\text{C}_5'\text{-H}$, $\text{C}_6'\text{-H}$, $\text{C}_7\text{-H}$, $\text{C}_8\text{-H}$, $\text{C}_9\text{-H}$, and $\text{C}_{10}\text{-H}$); ^{13}C -NMR (100MHz, CDCl_3): δ 189.8 (C-6), 171.7 (C-10b), 136.9 (C-1'), 134.9 (C-6a), 131.1 (C-5a), 132.4 (C-7), 130.6 (C-9), 128.5 (C-2' & C-6'), 125.8 (C-3' & 5'), 123.0 (C-4'), 120.4 (C-8), 116.4 (C-10), 100.2 (C-5), 98.2 (C-11a), 62.1 (C-2), 40.8 (C-5a), 39.7 (C-4a), 29.6 (C-4), 24.5 (C-3); MS (EI) m/z (int %) 318 (M^+ , 38.7), 235 (63.4), 234 (21), 233 (47.0) and 84 (100). *Anal.* Calcd for $\text{C}_{21}\text{H}_{18}\text{O}_3$: C, 79.22; H, 5.70. Found: C, 79.26; H, 5.73.

4a: (4aR*,5S*,11aS*) 3,4,4a,11a-Tetrahydro-5-phenyl-2H-indeno[1,2-b]pyrano[3,2-e]pyran-6(5H)-

one. mp 168-169 °C; IR (KBr) 1695, 1630 cm^{-1} ; ^1H -NMR (90MHz, CDCl_3): δ 1.61-1.73 (m, 2H, $\text{C}_4\text{-H}$), 1.78-1.89 (m, 2H, $\text{C}_3\text{-H}$), 2.14-2.20 (m, 1H, $\text{C}_{4a}\text{-H}$), 3.66 (br s, 1H, $\text{C}_5\text{-H}$), 3.78-3.84 (m, 1H, $\text{C}_2\text{-H}$), 3.93-4.02 (m, 1H, $\text{C}_2\text{-H}$), 5.47-5.50 (d, $J=2.6\text{Hz}$, 1H, $\text{C}_{11a}\text{-H}$), 7.18-7.51 (m, 9H, $\text{C}_2'\text{-H}$, $\text{C}_3'\text{-H}$, $\text{C}_4'\text{-H}$, $\text{C}_5'\text{-H}$, $\text{C}_6'\text{-H}$, $\text{C}_7\text{-H}$, $\text{C}_8\text{-H}$, $\text{C}_9\text{-H}$, and $\text{C}_{10}\text{-H}$); C^{13} -NMR (100MHz, CDCl_3): δ 189.8 (C-6), 171.7 (C-10b), 136.9 (C-1'), 134.9 (C-6a), 131.1 (C-5a), 132.4 (C-7), 130.6 (C-9), 128.5 (C-2' & C-6'), 125.8 (C-3' & 5'), 123.0 (C-4'), 120.4 (C-8), 116.4 (C-10), 100.2 (C-5), 98.2 (C-11a), 62.1 (C-2), 40.8 (C-5a), 39.7 (C-4a), 29.6 (C-4), 24.5 (C-3); MS (EI) m/z (int %) 318 (M^+ , 37.2), 235 (64.1), 234 (18.9), 233 (45.1) and 84 (100). *Anal.* Calcd for $\text{C}_{21}\text{H}_{18}\text{O}_3$: C, 79.22; H, 5.70. Found: C, 79.25; H, 5.73.

3d: (4aR*,5R*,11aS*) 3,4,4a,11a-Tetrahydro-5-(4'-fluorophenyl)-2H-indeno[1,2-b]pyrano[3,2-e]-pyran-6(5H)-one. mp 182-183 °C; IR (KBr) 1695, 1620 cm^{-1} ; ^1H -NMR (90MHz, CDCl_3): δ 1.58-1.70 (m, 2H, $\text{C}_4\text{-H}$), 1.89-2.18 (m, 2H, $\text{C}_3\text{-H}$), 2.19-2.25 (m, 1H, $\text{C}_{4a}\text{-H}$), 3.71-3.86 (m, 1H, $\text{C}_2\text{-H}$), 3.98-4.07 (m, 1H, $\text{C}_2\text{-H}$), 4.15-4.17 (d, $J=6.0\text{Hz}$, 1H, $\text{C}_5\text{-H}$), 5.71-5.74 (d, $J=2.1\text{Hz}$, 1H, $\text{C}_{11a}\text{-H}$), 6.95-7.38 (m, 8H, $\text{C}_2'\text{-H}$, $\text{C}_3'\text{-H}$, $\text{C}_5'\text{-H}$, $\text{C}_6'\text{-H}$, $\text{C}_7\text{-H}$, $\text{C}_8\text{-H}$, $\text{C}_9\text{-H}$ and $\text{C}_{10}\text{-H}$); C^{13} -NMR (100MHz, CDCl_3): δ 188.8 (C-6), 172.7 (C-10b), 135.9 (C-1'), 132.9 (C-6a), 131.0 (C-10a), 131.4 (C-7), 130.6 (C-9), 128.5 (C-2' & C-6'), 126.8 (C-3' & 5'), 124.0 (C-4'), 120.4 (C-8), 116.4 (C-10), 102.2 (C-5) 99.2 (C-11a), 60.1 (C-2), 40.8 (C-5a), 39.7 (C-4a), 29.6 (C-4), 24.6 (C-3); MS (EI) m/z (int %) 336 (M^+ , 22.8), 253 (34.7), 252 (16.8), 251 (30.9) and 84 (100). *Anal.* Calcd for $\text{C}_{21}\text{H}_{17}\text{O}_3\text{F}$: C, 74.99; H, 5.09; F, 5.65. Found: C, 74.99; H, 5.11; F, 5.68.

4d: (4aR*,5S*,11aS*) 3,4,4a,11a-Tetrahydro-5-(4'-fluorophenyl)-2H-indeno[1,2-b]pyrano[3,2-e]-pyran-6(5H)-one. mp 188-189 °C; IR (KBr) 1695, 1620 cm^{-1} ; ^1H -NMR (90MHz, CDCl_3): δ 1.61-1.76 (m, 2H, $\text{C}_4\text{-H}$), 1.81-1.90 (m, 2H, $\text{C}_3\text{-H}$), 2.09-2.16 (m, 1H, $\text{C}_{4a}\text{-H}$), 3.64 (br s, 1H, $\text{C}_5\text{-H}$), 3.79-3.84 (m, 1H, $\text{C}_2\text{-H}$), 3.94-4.02 (m, 1H, $\text{C}_2\text{-H}$), 5.46-5.47 (d, $J=2.4\text{Hz}$, 1H, $\text{C}_{11a}\text{-H}$), 6.98-7.43 (m, 8H, $\text{C}_2'\text{-H}$, $\text{C}_3'\text{-H}$, $\text{C}_5'\text{-H}$, $\text{C}_6'\text{-H}$, $\text{C}_7\text{-H}$, $\text{C}_8\text{-H}$, $\text{C}_9\text{-H}$ and $\text{C}_{10}\text{-H}$); C^{13} -NMR (100MHz, CDCl_3): δ 189.4 (C-6), 172.7 (C-10b), 156.1 (C-4'), 141.9 (C-1'), 137.9 (C-6a), 133.1 (C-10a), 132.4 (C-7), 130.6 (C-9), 127.5 (C-2' & C-6'), 122.8 (C-3' & 5'), 121.5 (C-8), 118.5 (C-10), 105.7 (C-5), 98.2 (C-11a), 62.1 (C-2), 40.8 (C-5a), 39.7 (C-4a), 29.6 (C-4), 24.5 (C-3); MS (EI) m/z (int %) 336 (M^+ , 20.2), 253 (32.0), 252 (9.1), 251 (9.1) and 84 (100). *Anal.* Calcd for $\text{C}_{21}\text{H}_{17}\text{O}_3\text{F}$: C, 74.99; H, 5.09; F, 5.65. Found: C, 74.98; H, 5.11; F, 5.68.

3e: (4aR*,5R*,11aS*) 3,4,4a,11a-Tetrahydro-5-(4'-carbomethoxy)phenyl-2H-indeno[1,2-b]pyrano[3,2-e]pyran-6(5H)-one. mp 170-171 °C; IR (KBr) 1720, 1690 cm^{-1} ; ^1H -NMR (90MHz, CDCl_3): δ 1.51-1.63 (m, 2H, $\text{C}_4\text{-H}$), 1.67-1.86 (m, 2H, $\text{C}_3\text{-H}$), 2.18-2.25 (m, 1H, $\text{C}_{4a}\text{-H}$), 3.65-3.73 (m, 1H, $\text{C}_2\text{-H}$), 3.89 (s, 3H, $-\text{CO}_2\text{CH}_3$), 3.91-3.96 (m, 1H, $\text{C}_2\text{-H}$), 4.21-4.23 (d, $J=6.0\text{Hz}$, 1H, $\text{C}_5\text{-H}$), 5.71-5.73 (d, $J=1.4\text{Hz}$, 1H, $\text{C}_{11a}\text{-H}$), 7.21-7.39 (m, 6H, $\text{C}_2'\text{-H}$, $\text{C}_6'\text{-H}$, $\text{C}_7\text{-H}$, $\text{C}_8\text{-H}$, $\text{C}_9\text{-H}$ and $\text{C}_{10}\text{-H}$), 7.94-7.96 (d, $J=$

8.4Hz, 2H, C_{3'}-H, C_{5'}-H); C¹³-NMR (100MHz, CDCl₃): δ 191.2 (C-6), 172.5 (C-10b), 167.1 (CO), 148.0 (C-4'), 146.8 (C-1'), 136.8 (C-6a), 134.0 (C-10a), 132.4 (C-7), 130.6 (C-9), 126.5 (C-2' & C-6'), 121.8 (C-3' & 5'), 121.5 (C-8), 118.5 (C-10), 104.8 (C-5), 98.2 (C-11a), 62.1 (C-2), 51.1 (CH₃), 40.8 (C-5a), 39.7 (C-4a), 29.6 (C-4), 24.5 (C-3) MS (EI) m/z (int %) 376 (M⁺, 20.1), 293 (26.4), 292 (3.5), 233 (21.0), 84 (100). *Anal.* Calcd for C₂₃H₂₀O₅: C, 73.39; H, 5.36. Found: C, 73.41; H, 5.38.

4e: (4aR*,5S*,11aS*) 3,4,4a,11a-Tetrahydro-5-(4'-carbomethoxyphenyl)-2H-indeno[1,2-b]pyrano[3,2-e]pyran-6(5H)-one. mp 178-179 °C; IR (KBr) 1720, 1690 cm⁻¹; ¹H-NMR (90MHz, CDCl₃): δ 1.63-1.73 (m, 2H, C₄-H), 1.79-1.91 (m, 2H, C₃-H), 2.13-2.19 (m, 1H, C_{4a}-H), 3.70 (br s, 1H, C₅-H), 3.79-3.85 (m, 1H, C₂-H), 3.89 (s, 3H, -CO₂CH₃), 3.94-4.02 (m, 1H, C₂-H), 5.46-5.47 (d, J=2.4Hz, 1H, C_{11a}-H), 7.26-7.44 (m, 6H, C_{2'}-H, C_{6'}-H, C₇-H, C₈-H, C₉-H and C₁₀-H), 7.95-7.98 (d, J=8.4Hz, 2H, C_{3'}-H and C_{5'}-H); C¹³-NMR (100MHz, CDCl₃): δ 191.4 (C-6), 172.7 (C-10b), 167.0 (CO), 148.0 (C-4'), 146.9 (C-1'), 136.8 (C-6a), 134.0 (C-10a), 132.4 (C-7), 130.6 (C-9), 126.5 (C-2' & C-6'), 121.8 (C-3' & 5'), 121.5 (C-8), 118.5 (C-10), 105.8 (C-5), 98.2 (C-11a), 62.1 (C-2), 51.1 (CH₃), 40.8 (C-5a), 39.7 (C-4a), 29.6 (C-4), 24.5 (C-3); MS (EI) m/z (int %) 376 (M⁺, 24.3), 293 (27.0), 292 (3.8), 233 (23.3) and 84 (100). *Anal.* Calcd for C₂₃H₂₀O₅: C, 73.39; H, 5.36. Found: C, 73.39; H, 5.39.

3f: (4aR*,5R*,11aS*) 3,4,4a,11a-Tetrahydro-5-(4'-nitrophenyl)-2H-indeno[1,2-b]pyrano[3,2-e]pyran-6(5H)-one. mp 190-191°C; IR (KBr) 1700, 1620, 1510, 1340 cm⁻¹; ¹H-NMR (90MHz, CDCl₃): δ 1.60-1.64 (m, 2H, C₄-H), 1.68-1.78 (m, 2H, C₃-H), 2.22-2.30 (m, 1H, C_{4a}-H), 3.85-3.89 (m, 1H, C₂-H), 4.00-4.06 (m, 1H, C₂-H), 4.28-4.30 (d, J=6.0Hz, 1H, C₅-H), 5.75-5.77 (d, J=1.4Hz, 1H, C_{11a}-H), 7.29-7.42 (m, 6H, C_{2'}-H, C_{6'}-H, C₇-H, C₈-H, C₉-H and C₁₀-H), 8.16-8.20 (d, 2H, J=8.7Hz, C_{3'}-H, C_{5'}-H); C¹³-NMR (100MHz, CDCl₃): δ 192.4 (C-6), 173.7 (C-10b), 150.0 (C-4'), 146.9 (C-1'), 136.9 (C-6a), 133.1 (C-5a), 132.4 (C-7), 130.6 (C-9), 128.5 (C-2' & C-6'), 123.8 (C-3' & 5'), 121.4 (C-8), 118.4 (C-10), 104.1 (C-5), 98.2 (C-11a), 62.1 (C-2), 40.8 (C-5a), 39.7 (C-4a), 29.6 (C-4), 24.5 (C-3); MS (EI) m/z (int %) 363 (M⁺, 31.6), 333 (18.9), 249 (72.3), 248 (37.4) and 84 (100). *Anal.* Calcd for C₂₁H₁₇NO₅: C, 69.41; H, 4.72; N, 3.85. Found: C, 69.42; H, 4.74; N, 3.87.

4f: (4aR*,5S*,11aS*) 3,4,4a,11a-Tetrahydro-5-(4'-nitrophenyl)-2H-indeno[1,2-b]pyrano[3,2-e]pyran-6(5H)-one. mp 198-199 °C; IR (KBr) 1700, 1620, 1510, 1340 cm⁻¹; ¹H-NMR (90MHz, CDCl₃): δ 1.68-1.85 (m, 2H, C₄-H), 1.86-1.90 (m, 2H, C₃-H), 2.14-2.19 (m, 1H, C_{4a}-H), 3.75 (br s, 1H, C₅-H), 3.82-3.87 (m, 1H, C₂-H), 3.96-4.04 (m, 1H, C₂-H), 5.45-5.46 (d, J=2.5Hz, 1H, C_{11a}-H), 7.35-7.45 (m, 6H, C_{2'}-H, C_{6'}-H, C₇-H, C₈-H, C₉-H, and C₁₀-H), 8.16-8.20 (d, J=8.7Hz, 2H, C_{3'}-H, C_{5'}-H); C¹³-NMR (100MHz, CDCl₃): δ 192.4 (C-6), 173.7 (C-10b), 150.0 (C-4'), 146.9 (C-1'), 136.9 (C-6a), 133.0 (C-10a), 132.4 (C-7), 130.6 (C-9), 128.5 (C-2' & C-6'), 123.8 (C-3' & 5'), 121.5 (C-8), 118.5 (C-10), 105.7 (C-5),

98.2 (C-11a), 62.1 (C-2), 40.8 (C-5a), 39.7 (C-4a), 29.6 (C-4), 24.5 (C-3); MS (EI) m/z (int %) 363 (M^+ , 28.8), 333 (13.1), 249 (71.9), 248 (41.7) and 84 (100). *Anal.* Calcd for $C_{21}H_{17}NO_5$: C, 69.41; H, 4.72; N, 3.85. Found: C, 69.43; H, 4.75; N, 3.86.

ACKNOWLEDGEMENTS

Mamta Sharma thanks Kurukshetra University, Kurukshetra while Sunita Bajaj, Kusum Sharma and Somesh Sharma, are grateful to CSIR, New Delhi for generous financial grants.

REFERENCES

1. L. W. Butz and A. W. Rytina, *Org. Reactions*, 1949, **5**, 136.
2. J. Sauer, *Angew. Chem., Int. Ed. Engl.*, 1967, **6**, 16.
3. D. Craig, *Chem. Rev.*, 1987, **16**, 187.
4. L. F. Tientze, *J. Heterocycl. Chem.*, 1990, **27**, 47.
5. G. Brieger, *J. Am. Chem. Soc.*, 1963, **85**, 3783.
6. L. H. Klemm and K. W. Gopinath, *Tetrahedron Lett.*, 1963, **26**, 1243.
7. H. O. House and T. H. Cronin, *J. Org. Chem.*, 1965, **30**, 1061.
8. R. A. Carboni and R. V. Lindsey, *J. Am. Chem. Soc.*, 1959, **81**, 4342.
9. J. Sauer and D. Lang, *Angew. Chem.*, 1964, **76**, 603.
10. R. B. Woodward and R. Hoffmann, 'The Conservation of Orbital Symmetry,' Verlag Chemie, GmBH, Academic Press Inc., 1971, 145-151.
11. G. Desimoni, G. Cellevine, G. Minoli, and G. Tacconni, *Tetrahedron*, 1972, **28**, 4003.
12. G. Desimoni, M. J. Cook, and G. Tacconi, *Annali Chim. Rome*, 1970, **60**, 208.