

HETEROCYCLES, Vol. 73, 2007, pp. 857 - 862. © The Japan Institute of Heterocyclic Chemistry
Received, 18th June, 2007, Accepted, 23rd July, 2007, Published online, 25th July, 2007. COM-07-S(U)18

SYNTHESIS OF OPTICALLY PURE 4-HYDROXYMETHYL-3-PHENOXY-2-AZETIDINONE FROM D-GLUCAL

Pilar Areces,^{a*} Esther Carrasco,^a Maria Monterde,^b Mark E. Light,^c and Joaquín Plumet^{b*}

a) Department of Organic Chemistry, Faculty of Chemistry, University of Extremadura, 06071-Badajoz, Spain. b) Department of Organic Chemistry, Faculty of Chemistry, University Complutense, 28040-Madrid, Spain. c) School of Chemistry, University of Southampton, Highfield, Southampton SO17 1BJ, UK

e-mail corresponding authors: pareces@unex.es; plumety@quim.ucm.es

Abstract –A convenient approach to enantiomerically pure 4-hydroxymethyl-3-phenoxy-2-azetidinone has been carried out using the easily available aldehyde 5 as chiral starting material.

The use of carbohydrate derivatives as chiral auxiliaries in β -lactam synthesis using the Staudinger reaction is well documented in the literature.¹ Regarding the asymmetric induction obtained from imines derived from sugar aldehydes, compounds such as **1**,² **2**² or **3**³ (Figure 1) have been used although with different levels of diastereoselectivity depending on the chiral auxiliary employed.⁴





Figure 1. Imines derived from sugar aldehydes


A major problem associated with this methodology is the elimination of the chiral auxiliary because, in some cases, the experimental conditions necessary to achieve this process are incompatible with the β -lactam moiety. In this way, the introduction of an appropriate, synthetically versatile functionality between the imino group and the sugar moiety in the imino derivative partner may be a convenient

solution to this problem. In this case, the chiral auxiliary, not directly attached to the β -lactam ring, could be removed or transformed under smooth conditions, which would be compatible with the β -lactam ring. Within this general approach a double bond constitutes an appealing possibility. Considering the distance between the chiral auxiliary and the imino group, low levels of diastereoselectivity should be hoped and the method will be useful providing a convenient separation of the diastereomeric mixture. (Scheme 1)




Scheme 1

We have tested this approach in the following way (Scheme 2): aldehyde **5**, easily accessible from tri-O-acetyl-D-glucal (**4**) in two steps and 66% overall yield,⁵ was transformed into imine **6** under standard conditions ($\text{C}_6\text{H}_5\text{CH}_2\text{NH}_2$, molecular sieves, Et_2O , 97% isolated yield). Reaction of **6** with phenoxyacetyl chloride **7** in the presence of Et_3N gave a diastereomeric mixture (1.3:1) of β -lactams **8a** and **8b** in 70% isolated yield after purification by column chromatography of the reaction crude (silica gel, ethyl acetate-hexane 1:3). The *cis* relationship at the β -lactam nucleus was deduced from the $J_{\text{H}3/\text{H}4}$ value (4.8 Hz). Diastereomerically pure β -lactams were isolated by HPLC (ethyl acetate-hexane 1:1 as eluent, flow 5 ml/min, $\lambda_{\text{opt}} = 280 \text{ nm}$).



Scheme 2

Having achieved the steps a) and b) of our methodology (Scheme 1) we decided to check the procedure fusing the steps c) and d) in only one. In this way, ozonolysis of the β -lactam **8a** followed by treatment with NaBH_4 allowed us for the synthesis of enantiomerically pure 4-hydroxymethyl β -lactam **9** in 66% isolated yield (Scheme 3).



Scheme 3

The relative configuration of **9** (and hence **8**) was determined by X-ray crystallography as 3*S*, 4*R* (Figure 2).




Figure 2

In summary, in this communication we have described a convenient procedure for the synthesis of optically pure β -lactams having a tethered chiral auxiliary able to be appropriately functionalized.

EXPERIMENTAL

General Methods

Chemicals and solvents were obtained from commercial sources and distilled or dried according to standard methods. Progress of the reaction was monitored by thin layer chromatography, which was performed on Aldrich precoated plated (silica gel, POLIGRAM[®]SIL G/UV₂₅₄ Machery-Nagel) with thickness of 0.22 mm, by staining with Hanessian's stain. Chromatographic purification was performed on silica gel columns (Kieselgel 60, 230-400 mesh, Merck) with an indicated eluent. Melting points were determined on a Barnstead Electrothermal 9100 apparatus and are uncorrected. Optical rotations were recorded on a Perkin-Elmer 241 MC polarimeter. IR spectra were obtained on a Thermo IR-300 spectrometer (Electron Corporation), as a thin film on NaCl plates. ¹H NMR and ¹³C NMR spectra were recorded on a Bruker AC 200 spectrometer at room temperature and 400 MHz and 100 MHz respectively, in CDCl₃ as solvent, with TMS as an internal standard. Splitting patterns are indicated as s, singlet; d, doublet; t, triplet; q, quartet; m, multiplet. High resolution mass spectra were recorded using a Micromass AutoSpec under CI (at 3000 dalton resolution) or Bruker Microtof ESI-TOF under ESI-TOF. Separation of isomers was performed by an HPLC Agilent 1100 Series apparatus (Agilent Technologies)

with a semipreparative Zorbax RX-Sil USHL001118 column (9.4 x 250 mm, 5-micron). Crystal data were collected on a diffractometer having a *Nonius Kappa CCD* area detector.

(3*S*,4*R*)-4-[(1'*E*)-3',4',5'-Tri-*O*-acetyl-1',2'-dideoxy-D-*erythro*-penten-1-yl]-*N*-benzyl-3-phenoxy-2-azetidinone (8a) and (3*R*,4*S*)-4-[(1'*E*)-3',4',5'-tri-*O*-acetyl-1',2'-dideoxy-D-*erythro*-penten-1-yl]-*N*-benzyl-3-phenoxy-2-azetidinone (8b).

To a solution of the imine **6** (1.52 mmol) in CH₂Cl₂ (6 mL) was added triethylamine (4.56 mmol) under N₂ at 0°C, and a solution of phenoxyacetyl chloride **7** (2.28 mmol) in CH₂Cl₂ (4 mL) was introduced dropwise. Then, the reaction was warmed to rt and stirred for 6.5 h. After the reaction was completed, the reaction mixture was poured into water (10 mL), stirred for 15 min, and then extracted with CH₂Cl₂ (3 x 10 mL). The combined organic layers were washed with 5% NaHCO₃ (3 x 10 mL), dried (MgSO₄) and concentrated *in vacuo*. The crude product was purified by column chromatography (silica gel, EtOAc/hexane 1:3) to afford a mixture (1.3:1) of the two diastereomers (colourless oil, 70%), which were separated by HPLC (hexane/EtOAc 1:1).

Isomer 8a: $[\alpha]_D^{21} +18^\circ$ (c 0.25, CH₂Cl₂); IR: $\nu_{\max}$ (film): 1747 s (N-C=O), 1747 s (C=O, ester), 1227 s (C-O-C), 1049 (=C-O) cm⁻¹. HRMS (CI, M+H) calcd for C₂₇H₃₀NO₈ 496.1971, found 496.1985; m/z (%) 496 (M+1, 10), 436 (M+1-AcOH, 63), 376 (M+1-2AcOH, 72), 302 (53), 172 (65), 107 (60), 91 (CH₂Ph, 100); ¹H-NMR: (400 MHz, CDCl₃) δ (ppm) 7.38-7.32 (m, 3H, Ph), 7.27-7.23 (m, 4H, Ph), 6.97 (t, *J* = 7.2 Hz, 1H, Ph), 6.91 (d, *J* = 8.0 Hz, 2H, Ph), 6.99 (t, *J* = 2 Hz, 1H, Ph), 6.90 (2H, Ph), 5.73-5.72 (m, H-1', H-2'), 5.45-5.42 (m, H-3'), 5.29 (d, *J*_{3,4} = 4.8 Hz, H-3), 5.06 (quintet, *J*_{4',3'} = 3.8 Hz, *J*_{4',5'b} = 3.6 Hz, *J*_{4',5'a} = 7.6 Hz, H-4'), 4.67 (d, *J*_{1'',a,1''b} = 14.8 Hz, H-1''_a), 4.29-4.26 (m, H-4), 4.10 (d, *J*_{1'',a,1''b} = 14.8 Hz, H-1''_b), 3.98 (dd, *J*_{4',5'a} = 8.0 Hz, *J*_{5'a,5'b} = 12.4 Hz, H-5'_a), 3.87 (dd, *J*_{4',5'b} = 3.6 Hz, *J*_{5'a,5'b} = 12.4 Hz, H-5'_b), 2.03 (s, 6H, CH₃OAc), 1.96 (s, 3H, CH₃OAc); ¹³C-NMR: (100 MHz, CDCl₃) δ (ppm) 170.4, 170.1, 169.2 (CO, 3AcO), 164.8 (CO, lactam), 156.9 and 134.8 (2C-1''') 130.8 (C-1'), 129.5, 128.9 and 128.1 (C-2'''/C-6''', 2C-3'''/C-5''') 128.6 and 128.5 (2C-4'''), 122.1 (C-2'), 115.3 (C-2'''/C-6''', PhO), 81.5 (C-3), 71.2 (C-4'), 71.6 (C-3'), 61.3 (C-5'), 59.3 (C-4), 44.6 (C-1'') 20.8, 20.7, 20.6 (CH₃).

Isomer 8b: $[\alpha]_D^{23} +26.5^\circ$ (c 0.66, CH₂Cl₂); IR: $\nu_{\max}$ (film): 1747 s (N-C=O), 1747 s (C=O, ester), 1227 s (C-O-C), 1047 (=C-O) cm⁻¹; HRMS (CI, M+H) calcd for C₂₇H₃₀NO₈ 496.1971, found 496.1982; m/z (%) 496 (M+1, 15), 436 (M+1-AcOH, 100), 376 (M+1-2AcOH, 85); ¹H-NMR: (400 MHz, CDCl₃) δ (ppm) 7.37-7.31 (m, 3H, Ph), 7.26-7.22 (m, 4H, Ph), 6.97 (t, *J* = 7.6 Hz, 1H, Ph), 6.90 (d, *J* = 3.2 Hz, 2H, Ph), 5.80 (dd, *J*_{1',2'} = 16.0 Hz, *J*_{1',4'} = 8.8 Hz, H-1'), 5.68 (dd, *J*_{2',3'} = 6.0 Hz, *J*_{1',2'} = 15.2 Hz, H-2'), 5.38-5.35 (m, *J*_{4',3'} = 4.4 Hz, *J*_{2',3'} = 6.0 Hz, H-3'), 5.28 (d, *J*_{3,4} = 4.4 Hz, H-3), 5.08 (q, *J*_{4',3'} = 4.8 Hz, *J*_{4',5'a} = 5.0 Hz, *J*_{4',5'b} = 5.2 Hz, H-4'), 4.74 (d, *J*_{1'',a,1''b} = 14.4 Hz, H-1''_a), 4.24 (dd, *J*_{3,4} = 4.4 Hz, *J*_{4,1'} = 8.8 Hz, H-4), 4.06 (d, *J*_{1'',a,1''b} = 14.8 Hz, H-1''_b), 3.97 (d, *J*_{4',5'} = 5.2 Hz, 2H, H-5'), 2.03 (s, 3H, CH₃OAc), 2.01 (s, 3H, CH₃OAc), 1.99 (s, 3H, CH₃OAc); ¹³C-NMR: (100 MHz, CDCl₃) δ (ppm) 170.4, 170.0, 169.2 (CO,

3AcO), 164.8, (CO, lactam), 131.3 (C-1'), 156.9 and 134.8 (2C-1'''), 131.3 (C-1'), 129.6 and 128.0 (2C-4''') 128.9, 128.8 and 128.5 (C-2'''/C-6''', 2C-3'''/C-5'''), 122.3 (C-2') 115.3 (C-2'''/C-6''', PhO), 81.5 (C-3), 71.3 (C-3'), 71.3 (C-4'), 61.4 (C-5'), 58.9 (C-4), 44.4 (C-1''), 20.8, 20.1 (3CH₃).

(3S, 4R)-4-hydroxymethyl-N-benzyl-3-phenoxy-2-azetidinone (9).

Through a stirred solution of β -lactam **8a** (0.169 mmol) in a mixture of CH₂Cl₂ (3.5 mL) and MeOH (4 mL) at -78 °C, O₂ was bubbled for 2 min and then ozone for 35 min. After that, the solution was purged, first with O₂ (10 min) and afterwards with argon, and treated portionwise with NaBH₄ (1.18 mmol). The reaction mixture was stirred for 10 min and then 2 h at rt. The mixture was diluted with NH₄Cl solution (3.2 ml), and extracted with CH₂Cl₂. The organic phase was then washed with water and dried (MgSO₄). After evaporation, reaction crude was purified by column chromatography (silica gel, EtOAc/hexane 1:2) to give the enantiomerically pure alcohol **9** (31.6 mg, 66%), as a white solid which was recrystallized from hexane/CH₂Cl₂. mp 69-72°C; [α]_D¹⁹ +11.9°, (c 1.3, CH₂Cl₂); IR: $\nu_{\max}$ (film): 3441 s (OH), 2927 s (CH₂), 1746 s (N-C=O), 1236 s (C-O-C), 1048 m (=C-O) cm⁻¹; HRMS (ESI-TOF, M+H) calcd for C₁₇H₁₈NO₃ 284.1272, found 284.1281; m/z (%) 284 (M+1, 100), 253 (M+1-CH₂OH, 82), 150 (96); ¹H-NMR: (400 MHz, CDCl₃) δ (ppm) 7.40-7.29 (m, 5H, PhO), 7.11-7.05 (m, 5H, Ph), 5.3 (d, $J_{3,4}$ = 3.5 Hz, H-3), 4.74 (d, $J_{1''a,1''b}$ = 11.9 Hz, H-1''_a), 4.34 (t, $J_{1''b,1''a}$ = 11.9 Hz, H-1''_b), 3.9-3.8 (m, 3H, H-1'_a, H-1'_b, H-4) 1.95 (OH) cm⁻¹. ¹³C-NMR: (100 MHz, CDCl₃) δ (ppm) 166.0 (CO, lactam), 157.6, 137.9, 135.6, 130.1, 129.4, 128.8, 128.4, 123.1 (C-2'''/C-6''', 2C-3'''/C-5''', 2C-4'''), 116.1 (C-2'''/C-6''', PhO), 81.2 (C-3), 60.6 (C-1'), 58.3 (C-4), 45.4 (C-1''). Crystal structure analysis: C₁₇H₁₇NO₃, M_r = 283.32 g mol⁻¹, monoclinic, space group *Pbca*, a = 23.4831 (9), b = 5.7989 (2), c = 23.9343 (9) Å, β = 118.879 (2) °, V = 2853.96 (18) Å³, Z = 8, ρ = 1.319 g cm⁻³, μ = 0.091 mm⁻¹, F(000) = 1200, crystal size: 0.16 x 0.07 x 0.06 mm³. Crystal data were collected on a diffractometer having a *Nonius KappaCCD* area detector. A total of 14110 reflections (3.65 < θ < 27.47°) were collected of which 3265 were unique R(int) = 0.0931. Crystallographic data (excluding structure factors) for compound **9** have been deposited with the Cambridge Crystallographic Data Centre as supplementary publication number CCDC-614962.

ACKNOWLEDGEMENTS

This work was supported by the Ministerio de Educación y Ciencia, Spain, Grants nos. BQU 2003-04967-C01 and CTQ2005-07676. E.C. thanks Ministerio de Educación y Ciencia, Spain, for a predoctoral fellowship, and M.M. thanks Comunidad Autónoma de Madrid for a postdoctoral fellowship.

REFERENCES (AND NOTES)

1. For a comprehensive review on the asymmetric synthesis of β -lactams through the Staudinger

- reaction, see: C. Palomo, J. M. Aizpurua, I. Ganboa, and M. Oiarbide, *Curr. Med. Chem.*, 2004, **11**, 1837.
2. D. R. Wagle, C. Garai, M. G. Monteleone, and A. K. Bose, *Tetrahedron Lett.*, 1988, **29**, 1649.
 3. M. Arun, S. N. Jushi, V. G. Puranik, B. M. Bhawal, and A. R. A. S. Deshmukh, *Tetrahedron*, 2003, **59**, 2309.
 4. Compare: a) C. Palomo, M. Oiarbide, and A. Landa, *J. Org. Chem.*, 2000, **65**, 41. b) C. Palomo, M. Oiarbide, A. Esnal, A. Landa, J. J. Miranda, and A. Linden, *J. Org. Chem.*, 1998, **63**, 5838.
 5. F. González, S. Lesage, and A. S. Perlin, *Carbohydr. Res.*, 1975, **42**, 267.