

NEW ACRIDONE ALKALOID AND COUMARIN FROM CITRUS PLANTS¹

Motoharu Ju-ichi*, Hiromi Kaga, Miki Muraguchi, and Mami Inoue
Faculty of Pharmaceutical Sciences, Mukogawa Women's University,
Nishinomiya, Hyogo 663, Japan

Ichiro Kajiura

National Institute of Agrobiological Resources, Tsukuba,
Ibaragi 305, Japan

Mitsuo Omura

Okitsu Branch, Fruit Tree Research Station, Ministry of Agriculture
Forestry & Fisheries, Okitsu, Shimizu, Shizuoka 424-02, Japan

Hiroshi Furukawa

Faculty of Pharmacy, Meijo University, Tempaku, Nagoya 468, Japan


Abstract ——— A new acridone alkaloid citramine (1) and a new coumarin osthenon (2) were isolated from the root of the hybrid seedlings of Ogonkan x Hyuganatsu and their structures were elucidated on the basis of spectroscopic data.

Our extensive investigations of the constituents of the genus Citrus have resulted in the isolation of many kind of acridone alkaloids and coumarins.^{1,2} We now wish to report the isolation and structure elucidation of a new acridone alkaloid, citramine (1) and a new coumarin, osthenon (2) from the roots of several hybrid seedlings³ resulting from a cross of Ogonkan⁴ (Citrus spp.) x Hyuganatsu (C. tamurana Tan.) and also from other Citrus plants.


Citramine (1), light yellow prisms, mp 277-279°C, had the molecular formula $C_{16}H_{15}NO_6$. The uv [$\lambda_{\max}^{EtOH}$ nm (log ϵ): 222 (4.25), 274 (4.76), 335 (4.12), 384 (3.84)] and ir spectra (3400, 1620, 1605, 1580 cm^{-1}) showed the characteristic absorptions of the 1-hydroxy-9-acridone system.⁵ The existence of an N-methyl and two methoxy groups was clear from the ^{13}C -nmr signals (δ 39.51, 59.80, 60.68) and 1H -nmr signals (δ 4.04, 3.87, 3.78). The presence of three phenolic hydroxy groups was apparent from the ir (3400 cm^{-1}) and 1H -nmr spectra (δ 15.03, 9.04, 8.91). In the aromatic proton region

of the ^1H -nmr spectrum, ortho-coupled doublets at δ 8.05 and 6.95 (each 1H, J = 8.79 Hz) and one proton singlet at δ 6.49 were observed. The lower field signal at δ 8.05 was characteristic of H-8 and higher field signal at δ 6.49 was assignable to H-4 in the 9-acridone skeleton,^{6,7} thus suggesting the location of these three aromatic protons at C_4 , C_7 , and C_8 . In NOE experiments, irradiation at the frequency corresponding to the N-methyl proton at δ 4.04 gave enhancements of 19.5 % of the signal at δ 6.49 (H-4). On the other hand, irradiation of the methoxy proton signals at δ 3.87 and 3.78 showed no enhancements at any aromatic proton signals. The above results indicated the location of the two methoxy groups at C_2 and C_5 , respectively. Consequently, the structure of citramine was established as formula 1 and this alkaloid was also isolated from *C. natsudaoidai* Hayata.⁸ Many types of acridone alkaloids have been known so far,⁹ but the 1,2,3,5,6-penta-oxygenated type compound is rare and citramine is the fourth example from natural sources.¹⁰⁻¹²

Osthenon (2) was obtained as slightly yellow prisms, mp 141-142°C. The molecular formula $C_{14}H_{12}O_4$ was determined by mass spectrometry and elemental analysis. The uv spectrum ($\lambda_{\text{max}}^{\text{EtOH}}$ 216 and 306 nm) showed the characteristic absorption of 7-oxygenated coumarins.¹³ The ^1H -nmr spectrum showed the presence of one methoxy (δ 4.00, 3H, s) and one acetyl (δ 2.43, 3H, s) groups. In the aromatic proton region, two pairs of signals at δ 7.65, 6.30 (each 1H, d, J = 9.52 Hz) and 7.46, 6.91 (each 1H, d, J = 8.79 Hz) were observed and these signals were diagnostic to C_4 -H, C_3 -H, C_5 -H and C_6 -H of the coumarin nucleus.¹⁴ Two olefinic protons at δ 7.98 and 7.34 (each 1H, d, J = 16.6 Hz) were assumed to trans-oriented α,β -unsaturated carbonyl moiety. These spectral data suggested the presence of trans-but-3-en-1-enyl side chain at C_8 and this speculation was confirmed by ^{13}C -nmr spectrum [δ 27.68 (q), 131.22 (d), 132.56



(1)



(2)

(d) and 199.40 (s)]. From the above spectral data, the structure of osthenon was confirmed as formula 2. Osthenon was also isolated from *C. natsudaoidai*, *C. tachibana*, *C. rugulapsa*, *C. tamurana* x *C. kinokuni*, and *C. hassaku* x *C. grandis*.¹

EXPERIMENTAL

Extraction and Isolation : The dried roots (800 g) of hybrid seedlings resulting from a cross of Ogonkan x Hyuganatsu were extracted with acetone. Evaporation of the solvent yielded 120.8g of residue. The residue was chromatographed over silica gel and eluted with hexane, benzene, 50% benzene-AcOEt, AcOEt, CH_2Cl_2 , acetone and MeOH. Each eluate was further separated with repeated PTLC using the solvent systems of acetone- CHCl_3 (1:19), iso-propyl ether and acetone-hexane (1:1) to afford citramine (1) (6 mg) and osthenon (2) (86.1 mg) as well as known compounds. Detailed procedure of separation and structures of known compounds will be reported elsewhere.

Citramine (1): light yellow prisms, mp 277-279°C, high ms: m/z 317.0895 (M^+ , found), 317.0899 (calcd. for $\text{C}_{16}\text{H}_{15}\text{NO}_6$); ir $\nu_{\text{max}}^{\text{KBr}}$: 3400, 1620, 1605, 1580 cm^{-1} ; uv $\lambda_{\text{max}}^{\text{EtOH}}$ nm (log ϵ): 222 (4.25), 274 (4.76), 335 (4.12), 384 (3.84); ^1H -nmr (acetone- d_6 , δ): 15.03 (1H, s), 9.04 (1H, s), 8.91 (1H, s), 8.05 (1H, d, $J = 8.79$ Hz), 6.95 (1H, d, $J = 8.79$ Hz), 6.49 (1H, s), 4.04, 3.87, 3.78 (each 3H, s); ^{13}C -nmr ($\text{DMSO}-\text{d}_6$, δ): 39.51 (q), 59.80 (q), 60.68 (q), 91.86 (d), 103.69 (s), 112.68 (d), 115.28 (s), 121.91 (d), 129.03 (s), 135.31 (s), 138.58 (s), 142.55 (s), 155.28 (s), 156.10 (s), 157.44 (s), 179.37 (s); ei-ms m/z: 317 (M^+), 302 (base peak), 287, 274, 272, 263, 205, 189.

Osthenon (2): slightly yellow prisms, mp 141-142°C, high ms: m/z 244.0763 (M^+ , found), 244.0753 (calcd. for $\text{C}_{14}\text{H}_{12}\text{O}_4$); ir $\nu_{\text{max}}^{\text{CHCl}_3}$: 1730, 1595 cm^{-1} ; uv $\lambda_{\text{max}}^{\text{EtOH}}$ nm (log ϵ): 216 (sh, 4.35), 306 (4.46); ^1H -nmr (CDCl_3 , δ): 7.98 (1H, d, $J = 16.6$ Hz), 7.65 (1H, d, $J = 9.52$ Hz), 7.46 (1H, d, $J = 8.79$ Hz), 7.34 (1H, d, $J = 16.6$ Hz), 6.91 (1H, d, $J = 8.79$ Hz), 6.30 (1H, d, $J = 9.52$ Hz), 4.00, 2.43 (each 3H, s); ^{13}C -nmr (CDCl_3 , δ): 27.68 (q), 56.35 (q), 107.72 (d), 111.77 (s), 112.97 (s), 113.56 (d), 130.02 (d), 131.22 (d), 132.56 (d), 143.46 (d), 153.88 (s), 159.92 (s), 161.71 (s), 199.40 (s); ei-ms m/z: 244 (M^+), 229, 213 (base peak), 201, 186, 173, 158.

ACKNOWLEDGEMENT

The authors express their deep gratitude to Dr. H. Tanino for differential NOE (400 MHz) experiments and Misses K. Suwa and S. Horiyama for their measurements of ms and nmr spectra.

REFERENCES AND NOTES

- 1) Part X on "Constituents of Domestic Citrus Plants". Part IX. C. Ito, M. Matsuoka, T. Mizuno, K. Sato, Y. Kimura, M. Inoue, I. Kajiura, M. Omura, M. Ju-ichi, and H. Furukawa, Chem. Pharm. Bull., submitted.
- 2) M. Ju-ichi, M. Yoneda, K. Sakiyama, M. Inoue, and H. Furukawa, Heterocycles, 1987, 26, 2077.
- 3) The plant was cultivated and collected in Fruit Tree Research Station, Okitsu Branch, Shizuoka.
- 4) Ogonkan is a mandarin-like Citrus cultivated rarely in Japan. M. Iwamasa, "Kankitsu no Hinshu (Citrus cultivars)", Shizuoka Citrus Grower's Coop. Assoc., Shimizu, Japan, 1976, p 255.
- 5) J. Reisch, K. Szendri, E. Minker, and I. Novak, Pharmazie, 1972, 27, 208.
- 6) A. W. Sangster and K. L. Stuart, Chem. Rev., 1965, 65, 69; R. D. Brown and F. N. Lahey, Aust. J. Sci. Res., 1950, A3, 593.
- 7) J. S. Shah and B. K. Sabata, Ind. J. Chem., 1982, 21B, 16.
- 8) Isolation and characterization will be reported elsewhere.
- 9) M. F. Grundon, Nat. Prod. Rep., 1987, 4, 232 and references cited therein.
- 10) J. Vaquette, M. O. Cleriot, M. R. Paris, J. L. Pousset, A. Cave, and R. R. Paris, Plant. Med. Phytothera., 1974, 8, 57.
- 11) I. H. Bowen and Y. N. Patal, Phytochemistry, 1986, 25, 429.
- 12) D.-K. Qin, Acta Pharm. Sinica, 1986, 21, 683.
- 13) G. B. Guise, E. Ritchie, R. G. Senior, and W. C. Taylor, Aust. J. Chem., 1967, 20, 2429.
- 14) W. Steck and M. Mazurek, Lloydia, 1972, 35, 418.

Received, 6th May, 1988