

SYNTHESIS OF A NEW SKELETON, 2,6-EPITHIO-3-BENZAZOCINE

Mikio Hori*, Tadashi Kataoka, Hiroshi Shimizu, Eiji Imai, Noriyuki Iwata,
and Norihiro Kawamura

Gifu Pharmaceutical University, 6-1, Mitahora-higashi 5-chome, Gifu 502,
Japan


Masayasu Kurono

Nagoya Laboratory, Sanwa Kagaku Kenkyusho, Ltd., 1212, Gejo-cho, Kasugai,
Aichi 486, Japan

Abstract — Treatment of isothiochroman 2-oxide derivative (9) with acetic anhydride followed by heating in Dowtherm A (a mixture of biphenyl and diphenyl ether) afforded a novel 2,6-epithio-3-benzazocine derivative (2).

In connection with our study on sulfur-containing analgesic compounds, thieno[3,4-b]morphinans,¹ 8-mercapto-3-benzazocines,² and [1]benzothiopyrano[3,4-b]pyrroles³ have been synthesized in our laboratory.

It is already known that 3-benzazocines (3) had not clinically shown analgesic activity.⁴ 2,6-Epithio-3-benzazocines (2) were designed on the assumption that the epithiobenzazocines (2) would be metabolized with a radical cleavage of the C-S bond to give non-analgesic 3-benzazocine derivatives.⁵ Therefore, the epithiobenzazocines (2) would be a candidate for the non-narcotic analgesics. In this communication, the synthesis of 3-ethoxycarbonyl-1,1,6-trimethyl-2,6-epithio-3-benzazocine is described in order to learn the chemical properties of the 2,6-epithio-3-benzazocine skeleton in which the carbon atom of 11-position of benzomorphans (1) is displaced by a sulfur atom.



1-(2-Ethoxycarbonylaminoethyl)-1,4,4-trimethylisothiochroman 2-oxide (9), a key intermediate for the synthesis of 2,6-epithio-3-benzazocine skeleton, was synthesized from 4,4-dimethylisothiochroman (4) in several steps as shown in Chart 1.




Chart 1

4,4-Dimethylisothiochroman (4)⁶ was chlorinated with sulfuryl chloride and treated with mercuric cyanide to give 1-cyano-4,4-dimethylisothiochroman (5). The cyanide (5) was methylated with sodium hydride and methyl iodide, and then the cyano group was hydrolyzed with KOH to lead to isothiochroman-1-carboxylic acid (6). The acid (6) was submitted to reduction with LiAlH_4 and the resulting alcohol was chlorinated with $\text{CCl}_4\text{-PPh}_3$. Treatment of the chloride with KCN gave the cyanide (7). Reduction of the cyanide (7) with LiAlH_4 and then treatment with ethyl chloro-carbonate gave 1-(2-ethoxycarbonylaminoethyl)isothiochroman (8).

In order to cyclize the compound (8) under the Pummerer reaction conditions,⁷ the compound (8) was oxidized with m-CPBA to give the sulfoxide (9), which was a diastereoisomeric mixture in the ratio of 1:1.





Chart 2

The sulfoxide (9) was refluxed in acetic anhydride for 24 hr to give the desired sulfide (10)⁸ (18%), the 3-acetoxy product (11) (34.1%), and 3-acetoxy-N-acetyl compound (12) (45.0%). When the reaction was followed by TLC, it was observed that the compound (11) appeared first and then was converted into 10 and 12. On refluxing 9 in acetic anhydride for 1 hr 11 was obtained in 86.6% yield. Therefore, the cyclization conditions of 11 to 10 were investigated and we found the optimal conditions that the sulfide (11) was heated in Dowtherm A at 200 °C for 2.5 hr to give 10 in 71.3% yield. As a result, 2,6-epithio-3-benzazocine (10) has been eventually synthesized from 9 via 11 in 61.7% yield.

Since the sulfide (11) was a diastereomeric mixture in the ratio of 1:1 and the yield was over 50%, the cyclization reaction in Dowtherm A is considered to proceed definitely via the S_N2 mechanistic path in view of the recent work of Uchida and Oae.⁹



The structure of 10 was determined by the ¹H-nmr spectral data showing two singlets at 5.08 and 5.27 ppm, which were attributable to 2-H. This observation could be explained in terms of tautomerism of the urethane moiety.

Pharmacological evaluations of 2,6-epithio-3-benzazocine derivatives are now in progress.

REFERENCES AND NOTES

1. M. Ban, Y. Baba, K. Kimura, Y. Kondo, K. Suzuki, and M. Hori, *Chem. Pharm. Bull.*, 1976, 24, 1679.
2. M. Hori, M. Ban, E. Imai, N. Iwata, Y. Baba, H. Fujimura, M. Nozaki, and M. Niwa, *Heterocycles*, 1983, 20, 2359; M. Hori, M. Ban, E. Imai, Y. Suzuki, Y. Baba, T. Morita, H. Fujimura, M. Nozaki, and M. Niwa, *J. Med. Chem.*, 1985, 28, 1656.
3. M. Hori, T. Kataoka, H. Shimizu, E. Imai, N. Iwata, N. Kawamura, and M. Kurono, *Heterocycles*, 1985, 23, 1381.
4. Y. Sawa, T. Kato, A. Morimoto, T. Masuda, M. Hori, and H. Fujimura, *Yakugaku Zasshi*, 1975, 95, 261.
5. S. Oae, K. Asada, and T. Yoshimura, *Tetrahedron Lett.*, 1983, 24, 1265; S. Oae, A. Mikami, T. Matsuura, K. Ogawa-Asada, Y. Watanabe, K. Fujimori, and T. Iyanagi, *Biochem. Res. Commun.*, 1985, 131, 567; T. Numata, Y. Watanabe, and S. Oae, *Tetrahedron Lett.*, 1979, 1411; Y.

- Watanabe, T. Numata, T. Iyanagi, and S. Oae, Bull. Chem. Soc. Jpn., 1981, 54, 1163; Y. Watanabe, S. Oae, and T. Iyanagi, ibid., 1982, 55, 188.
6. A. M. El-Khawasa, M. F. El-Zohry, M. T. Ismail, A. A. Abdel-Wahab, and A. A. Khalaf, Phosphorus and Sulfur, 1987, 29, 265.
7. S. Oae and T. Numata, Tetrahedron, 1974, 30, 2641; I. Nagakura, H. Oka, and Y. Nitta, Heterocycles, 1975, 3, 453; see the whole views on the Pummerer reaction in S. Oae and T. Numata, "The Pummerer Reactions and Related Reactions" Chapter 2 of "Isotopes in Organic Chemistry" Vol. 5 ed. by E. Buncl and C. C. Lee, Elsevier Ltd. (1980) and S. Oae, T. Numata, and T. Yoshimura, "Heterosulfonium Salts" Chapter 15 of "The Chemistry of the Sulfonium Group" ed. by J. M. Stirling and S. Patai, John Wiley & Sons Ltd. (1981).
8. 10: Ir (KBr) cm^{-1} : 1700 (C=O). $^1\text{H-Nmr (CDCl}_3\text{)}$ δ : 1.27 and 1.32 (3H, t, $J=7.0$ Hz, CH_2CH_3), 1.44 and 1.50 (3H, s, 1-H), 1.70 (3H, s, 6- CH_3), 3.20 (3H, m, 4-H and 5-H), 3.90 - 4.50 (1H, m, 4-H), 4.21 and 4.24 (2H, q, $J=7.0$ Hz, CH_2CH_3), 5.08 and 5.27 (1H, s, 2-H), 7.20 - 7.50 (4H, m, ArH). MS m/e : 305 (M^+), 158 (base).
9. Y. Uchida and S. Oae, Gazz. Chim. Ital., 1987, 117, 649.

Received, 25th May, 1988