

SYNTHESIS OF 5-ALKYLPYRROLIZIDIN-3-ONES FROM THE
 LUKES-ŠORM LACTAM BY GRIGNARD REACTION AND RECYCLIZATION
 OF INTERMEDIATE 5-BROMOALKYLPYRROLIDIN-2-ONES

Kimio Takahashi and Arnold Brossi*

Section on Medicinal Chemistry, Laboratory of
 Analytical Chemistry, NIDDK, National Institutes of
 Health, Bethesda, Maryland 20892, U.S.A.


Abstract - Synthesis of 5-alkylpyrrolizidin-3-ones 7 and 8 from pyrrolizidin-3,5-dione 1 is described. Grignard reaction of 1 gave 5-(3'-oxobutyl)pyrrolidin-2-one 2. Reduction of 2 with sodium borohydride, followed by mesylation, bromination, and cyclization afforded a diastereomeric mixture 7a and 8a, which were separated by chromatography. The same procedure with 3, also obtained from 1 by a Grignard reaction, afforded 7b and 8b.

Lukes-Šorm's¹ dilactam 1 is a valuable compound in medicine and in chemistry. Under the generic name rolziracetam, 1 has recently been reported to have amnesia-reversal activity and was subject of a multicenter, phase 2 clinical study in memory-impaired elderly patients, as well as in individuals with primary degenerative dementia (PDD)². We reported³ a synthesis of cis-5-*n*-propylpyrrolizidine (9), whose analogs are found among ant toxins⁴⁻⁶. Recently, we reported a conversion of 1 into trans-2,5-dialkylpyrrolidine⁶, also found among natural ant toxins, and an improved synthesis of 1 and ring opening reaction of 1 with Grignard reagents and other nucleophiles⁷. We now report a simple conversion of 1 into cis-5-alkylpyrrolizidin-3-ones (7a,b) and trans-isomers 8a,b.

Reaction of 1 with methylmagnesium bromide and *n*-propylmagnesium chloride gave ketolactams 2 and 3, respectively, in good yields⁷. Reduction of ketone 2 with sodium borohydride on silica gel gave the alcohol 4a as a mixture of stereo-

isomers. Because hydroxylactam **4a** is soluble in water, supported borohydride gave better result. Mesylation of the alcohol **4a** and treatment of the mesylate **5a** with lithium bromide in acetone gave the bromide **6a** in 64% yield. The same procedure with **3** yielded **6b**. Reaction of the bromide **6a** with sodium hydride in tetrahydrofuran gave a mixture of *cis*-5-methylpyrrolizidin-3-one (**7a**) and *trans*-isomer **8a**, in 37% and 35% yield, respectively. Although the synthesis described

SCHEME



here is not stereoselective, the isomers are effectively separated by flash chromatography affording the *cis*-isomer as the less polar material. Bromide **6b** when similarly treated with sodium hydride afforded *cis*-5-*n*-propylpyrrolizidin-3-one (**7b**) and *trans*-isomer **8b**, in 34% and 32% yields, respectively, which were identical with the samples prepared by the reported procedure². The *cis*-isomer **7b**, whose stereochemistry was confirmed by X-ray analysis³ of the thioketal of *cis*-5-acetonylpyrrolizidin-3-one, shows the nmr signals of C₅- and C₆-protons at 3.55 and 3.88 ppm, respectively. On the other hand, the nmr signals of the protons at C₅- and C₆-positions of *trans*-isomer **8b** were observed at 3.55-4.10 ppm as a multiplet². The nmr comparison of **7a** and **8a** allowed to assign to **7a** the *cis*- and to **8a** the *trans*-configuration.

Ketones **2** and **3** obtained from dilactam **1** by Grignard reaction were used for synthesis of 5-alkylpyrrolidin-2-ones and derived pyrrolidine ant alkaloids. It is now shown that these ketolactams are also versatile intermediates for synthesis of 5-alkylpyrrolizidin-3-ones and derived pyrrolizidine ant alkaloids^{4,5}.

EXPERIMENTAL

Ir spectra were determined by using a Beckman 4230 instrument in chloroform solution. ^1H -Nmr spectra were determined in deuterochloroform by using a Varian HR-220 spectrometer or a Jeol FX-100 spectrometer. Ms were obtained with a V. G. Micromass 7070F spectrometer with a Perkin-Elmer Sigma 3 gas chromatograph equipped with 2% OV-1 column, or a Hitachi-Perkin-Elmer RMU-6E spectrometer (70 eV). Chemical ionization (CI) Ms were obtained by using a Finnigan 1015D spectrometer. Silica gel (230-400 mesh) for flash chromatography was from EM Laboratories.

5-(3'-Hydroxy-n-butyl)pyrrolidin-2-one (4a). A mixture of the ketone **2** (155 mg, 1.0 mmol), NaBH_4 on silica gel (Aldrich, 8%, 160 mg) in CH_2Cl_2 (2.0 ml) was stirred at 25°C for 10 min. The mixture was diluted with 5% MeOH in CH_2Cl_2 , filtered through a silica gel column to give **4a** (101 mg, 64%) as an oil: ir 3440 and 1695 cm^{-1} ; nmr 7.25 (1H, br s, NH), 3.77 and 3.64 (each 1H, each br s), 2.72 (1H, br s), 2.39-2.05 (4H, m), 1.81-1.36 (6H, m), and 1.16 (3H, d, $J = 6\text{ Hz}$, C-Me); CI-Ms (NH_3 , m/z) 158 ($M^+ + 1$) and 174 ($M^+ + 17$).

5-(3'-Hydroxy-n-hexyl)pyrrolidin-2-one (4b). The ketone **3** (1.50 g, 8.2 mmol) was treated with NaBH_4 on silica gel (Aldrich, 8%, 1.60 g) in CH_2Cl_2 (50 ml) as above to give **4b** (1.32 g, 97%) as an oil: Ir 3440 and 1680 cm^{-1} ; nmr 7.25 (1H, br s, NH), 3.80-3.32 (2H, m), 2.95-2.65 (1H, m), 2.45-1.10 (12H, m), and 0.87 (3H, br t, $J = 6\text{ Hz}$, C-Me); ms (m/z) 185 (M^+).

5-(3'-Bromo-n-butyl)pyrrolidin-2-one (6a). MsCl (573 mg, 5.0 mmol) was added to a stirred solution of the alcohol **4a** (985 mg, 5.0 mmol) and Et_3N (505 mg, 5.0 mmol) in CH_2Cl_2 (30 ml) at 0°C . The mixture was stirred at 0°C for 1 h, washed with brine, dried (MgSO_4), and evaporated to give **5a** as an oil: Nmr 8.00 (1H, br s, NH), 4.95-4.65 (1H, m), 3.95-3.70 (1H, m), 3.00 (3H, s, Ms), 2.70-1.50 (8H, m), and 1.42 (3H, d, $J = 6\text{ Hz}$, C-Me), which was used in the next step without further purification. A mixture of the mesylate **5a**, LiBr (1.00 g, 11.5 mmol), and acetone (50 ml) was stirred at 25°C for 15h. The mixture was diluted with CH_2Cl_2 , washed with brine, dried (MgSO_4), and evaporated to give **6a** (705 mg, 64%) as an oil: Ir 3440 and 1680 cm^{-1} ; nmr 7.65 (1H, br s, NH), 4.30-3.80 and 3.80-3.35 (each 1H, each m), 2.45-1.40 (8H, m), and 1.76 (3H, d, $J = 6\text{ Hz}$, C-Me); ms (m/z) 219 and 221 (M^+).

5-(3'-Bromo-n-hexyl)pyrrolidin-2-one (6b). The alcohol **4b** (1.85 g, 10 mmol) was treated with MsCl (1.14 g, 10 mmol) and Et_3N (1.01 g, 10 mmol) in CH_2Cl_2 (50

ml) as above to give 5-(3'-mesyloxy-n-hexyl)pyrrolidin-2-one (**5b**): Nmr 7.65 (1H, br s, NH), 4.90-4.35 (1H, m), 3.85-3.35 (1H, m), 3.00 (3H, s, Ms), 2.70-1.10 (12H, m), and 0.92 (3H, br t, $J = 6$ Hz, C-Me). Reaction of the mesylate **5b** and LiBr (3.0 g, 34.5 mmol) in acetone (50 ml) carried out as above yielded **6b** (1.56 g, 63%): Ir 3440 and 1690 cm^{-1} ; nmr 7.30 (1H, br s, NH), 4.15-3.30 (2H, m), 2.55-1.10 (12H, m), and 0.89 (3H, br t, $J = 6$ Hz, C-Me); ms (m/z) 247 and 249 (M^+).

cis-5-Methylpyrrolizidin-3-one (7a) and trans-5-Methylpyrrolizidin-3-one (8a). A mixture of the bromide **6a** (220 mg, 1.0 mmol) and NaH (50% dispersion in mineral oil, 48 mg, 1.0 mmol) in THF (2.0 ml) was stirred at 25°C for 15 h. The mixture was diluted with CH_2Cl_2 , washed with brine, dried (MgSO_4), and evaporated to give an oil, which was chromatographed on silica gel with $\text{CH}_2\text{Cl}_2/\text{MeOH} = 95/5$ as the eluent yielding **7a** (53 mg, 37%) as an oil: Ir 1680 cm^{-1} ; nmr 3.95 (1H, m, $\text{C}_\alpha\text{-H}$), 3.55 (1H, m, $\text{C}_\beta\text{-H}$), 3.10-1.00 (8H, m), and 1.05 (3H, d, $J = 6$ Hz, C-Me); ms (m/z) 139 (M^+), and oily **8a** as the more polar compound (49 mg, 35%): Ir 1680 cm^{-1} ; nmr 4.16-3.55 (2H, m, $\text{C}_\alpha\text{-H}$ and $\text{C}_\beta\text{-H}$), 3.00-1.00 (8H, m), and 1.05 (3H, d, $J = 6$ Hz, C-Me); ms (m/z) 139 (M^+).

cis-5-n-Propylpyrrolizidin-3-one (7b) and trans-5-n-Propylpyrrolizidin-3-one (8b). A mixture of the bromide **6b** (1.00 g, 4.0 mmol), NaH (50% dispersion in mineral oil, 192 mg, 4.0 mmol) in THF (30 ml) was stirred at 25°C for 15 h. Work-up as mentioned above yielded **7b** (228 mg, 34%) and **8b** as the more polar compound (216 mg, 32%), which were both identical with authentic samples⁹ by TLC, ir, and nmr comparison.

ACKNOWLEDGEMENT

We would like to thank Dr. P. A. Wehrli of the Research Division of Hoffmann-La Roche, Nutley, N.J., for having initiated several reductive aminations of γ -oxopimeric acid for the preparation of lactam **1**.

REFERENCES AND NOTES

The compounds described in this paper are racemic, and only one series of enantiomers is depicted for convenience.

1. R. Lukes and F. Šorm, Collect. Czech. Chem. Commun., 1947, **12**, 278.
2. D. E. Butler, J. D. Leonard, B. W. Caprathe, Y. J. L'Italien, M. R. Pavia, F. M. Hershenson, P. H. Poschel, and J. G. Marriot, J. Med. Chem., 1987, **30**, 498.
3. P. Buchs, A. Brossi, and J. Flippen-Anderson, J. Org. Chem., 1982, **47**, 719.
4. T. H. Jones, M. S. Blum, H. M. Fales, and C. R. Thompson, J. Org. Chem., 1980, **45**, 4778.
5. A. Numata and T. Ibuka, "Alkaloids from Ants and Other Insects", in "The Alkaloids", (ed. A. Brossi), Vol. **31**, Academic Press, 1987, pp 193-315.
6. W. Gessner, K. Takahashi, A. Brossi, M. A. Kaliner, and M. Kowalski, Helv. Chim. Acta, 1987, **70**, 2003.
7. R. Alonso, W. Gessner, K. Takahashi, and A. Brossi, Synth. Commun., 1988, **18**, 37.
8. G. Hall, J. K. Sugden, and M. B. Waghela, Synthesis, 1987, 10.
9. M. Ikeda, T. Sato, and H. Ishibashi, Heterocycles, 1988, **27**, 1465.

Received, 4th July, 1988