

SYNTHESIS OF VINCA ALKALOIDS AND RELATED COMPOUNDS. XLIII¹.

UNEXPECTED REARRANGEMENT OF 3-ACYLINDOLENINES

Zsuzsanna Kardos-Balogh, Ferenc Sóti, Mária Incze,



Mária Kajtár-Peredy, Lajos Radics, and Csaba Szántay*

Central Research Institute for Chemistry of the Hungarian

Academy of Sciences, H-1525 Budapest, POB 17, Hungary


Abstract - The stable 3-acylindolenine derivative (1) was transformed to new heterocyclic compounds (2, 3) through unusual rearrangements.

Recently, we reported the synthesis of the 3-acylindolenine derivative 1 via intramolecular acylation². Our subsequent studies on the chemical behaviour of this molecule disclosed a remarkable reluctance of the oxo group towards usual carbonyl reactions. Thus, treatment of 1 with NaBH₃CN in CF₃COOH has left the keto function intact and led, rather, to the reduction of the C-N double bond giving product 2 (in addition to minor amounts of 3).


12 R=H3 R=CH₂CF₃

With the conjugated double bond thus eliminated, much of the regular chemical reactivity of oxo group could be restored: reaction of 2 with hydroxylamine gave

the oxime 4 while the reduction of 2 with LiAlH_4 afforded the epimeric alcohols 5 and 6.




4




	R ₁	R ₂
<u>5</u>	HO	H
<u>6</u>	H	OH


Surprisingly, however, acetylation of 2 with $\text{AcOH}/\text{Ac}_2\text{O}$ mixture gave 7, a rearranged triacetyl derivative, in high yield rather than the expected N-acetyl-2. In its turn, when hydrolyzed with sodium hydroxide in ethanol under reflux, 7 is converted into 8 (40.5 %) and 9 (47.5 %). A closer study of this reaction disclosed that products 8 and 9 are in equilibrium and can be mutually interconverted. Refluxing of either 8 or 9 in ethanol affords the same product mixture in which the ratio of 8:9 is approx. 2.2:1.



7



8




9


The constitution and stereochemistry of the new products as portrayed in 7-9 were inferred from high field ^1H (400 MHz) and ^{13}C (100.6 MHz) nmr spectra. The assignment of the resonances in terms of proton and carbon chemical shifts, δ_{H} , δ_{C} , ^{13}C -multiplicities and interproton couplings, J_{HH} , was performed by means of standard one- and two-dimensional (1D, 2D) FT nmr techniques and the fully assigned spectral parameters are collected in the Experimental. First, proton-proton chemical shift correlation (COSY) experiment³ was run to establish ^1H - ^1H connectivities within the molecular framework and, then, these pieces of informa-

tion, combined with results of carbon-proton chemical shift correlation experiment (mediated by one-bond C-H couplings)³ served as the starting point for the determination of the carbon-carbon connectivities. Carbon-carbon sequences involving quaternary ^{13}C atoms were inferred from a series of carbon-proton chemical shift correlation experiments in which the relevant time periods were systematically varied such as to obtain observable proton-to-carbon magnetization transfer for the expected range (1.5 to 7 Hz) of multiple bond $^n\text{J}_{\text{CH}}$ ($n=2,3,4$) couplings³. Examination of the collected spectral parameters shows that these are best accommodated within the proposed formulas 7, 8 and 9. The most relevant stereochemical feature of 7 and 8 follows from the value of $^3\text{J}_{11\text{a},11\text{b}}$ (6.7 Hz) which suggests that these protons assume a *gauche* steric disposition. The stereochemistry thus inferred has received further support by the occurrence of sizable four-bond ^1H - ^1H couplings between $\text{H}_{12\text{eq}}$ and $\text{H}_{11\text{b}}$ (1.7 Hz), as well as between $\text{H}_{12\text{ax}}$ and $\text{H}_{2\text{ax}}$ (2.3 Hz). Formation of 9 gives rise to major variations in the stereochemistry of the molecule. This is clearly reflected by the 9.9 Hz assumed by the vicinal coupling $^3\text{J}_{12\text{a},12\text{b}}$ in 9, a value characteristic of protons in *trans* relative orientation. Corroborative evidence for the resulting stereochemistry as shown in 9 was readily available from selective ^1H - $\{^1\text{H}\}$ NOE experiments. Preirradiation of the resonance due to H-12a resulted in signal enhancements of resonances due to H-5_a, H-13B, H-2_a, H-4_a and H-14A, while performing the same experiment with the preirradiation frequency set at the resonance of H-12b gave rise to enhanced signal intensities for protons N-COCH₃ and CH_AH_BCH₃. It can be seen that the spatial proximities reflected by the observed NOE effects are best accounted for by the stereochemistry portrayed in 9. Thus two unexpected rearrangements (2 $\rightarrow$ 7 and 8 $\rightarrow$ 9) were observed. In the first case (2 $\rightarrow$ 7) the rearrangement presumably starts by the attack of an acetate anion at C₆, the latter site being in α -position to the protonated nitrogen. In principle, the attack could also occur at the proton of the C₇-H bond causing a Hoffmann type elimination⁴, or at several other electrophilic centers of the molecule. Position C₆ is apparently favoured owing to the through-space neighbouring group effect of the sterically close keto-function⁵. The ring opening is followed by the formation of the aminocarbino function and subsequent O-acetylation. Again, the last step is rather surprising since under the given conditions pseudobasic aminocarbinols usually afford the corresponding iminium salt⁶. In our case, however, formation of the bridge-head double bond is unfavoured

as Bredt's rule points out.



Compound 7 can in principle also be formed through an elimination-addition mechanism.



The second rearrangement, 8 $\rightleftharpoons$ 9, can be rationalized by assuming the formation of 10, an intermediate ketone, which is in equilibrium with both 8 and 9. The ready conversion of 8 $\rightleftharpoons$ 10 is the consequence of the destabilization of the amino-carbinol function by deacetylation, while transition 10 $\rightleftharpoons$ 9 is governed by the facile transacetylation of the two, sterically proximate amino groups.

EXPERIMENTAL

Infrared spectra were recorded on a Nicolet 7199 Fourier transform spectrometer and the frequencies (cm^{-1}) of significant peaks are reported. All nmr spectra were run on deuteriochloroform solutions at ambient temperature using a Varian Associates model XL-100 for low-field and model XL-400 instrument for high field conventional and 2D experiments. Selective ^1H - ^1H NOE measurements were performed in the difference

mode. Mutual $^1\text{H} - ^1\text{H}$ couplings are given only once, at their first occurrence in the Experimental. Mass spectra were recorded on an AEI MS-902 mass spectrometer (70 eV, ion source temp. 200 °C, direct inlet). The purification of the compounds was carried out by column chromatography on silica gel (Merck Kieselgel 60, 0.063 - 0.2 mm).

(-)-(1S:7aR:12aS:12bS)-1-Ethyl-15-oxo-1,2,3,4,6,7,7a,12,12a,12b-decahydro-1,7a-propanoindolo[2,3-a]quinolizin 2

(-)-(1S:7aR:12aS:12bS)-1-Ethyl-15-oxo-12-(2,2,2-trifluoro-ethyl)-1,2,3,4,6,7,7a,12,12a,12b-decahydro-1,7a-propanoindolo[2,3-a]quinolizin 3.

A solution of 1 (10.0 g, 0.0324 mol) in CF_3COOH (100 ml) was cooled to -10 °C, and NaBH_3CN (5.0 g, 0.0798 mol) was added in small portions in argon atmosphere. The reaction mixture was stirred at -10 ° for 1 h and at room temperature overnight, then poured onto ice (500 g). The organic layer was separated and the aqueous phase was extracted with CHCl_3 (2 x 300 ml). The combined extracts were neutralized by 5 % NaHCO_3 , then dried over MgSO_4 and evaporated in *vacuo*, the residue was dissolved in EtOH to afford 2 (5.45 g, 54 %); mp 237-238 °C (ethanol); $[\alpha]_{\text{D}}^{25} -557.4^\circ$ ($c = 1.0$, CHCl_3); ms (m/z , %) 310 (M^+ , 8.5), 254 (11), 180 (89), 144 (11), 143 (14), 124 (100); ir (KBr), (ν , cm^{-1}) 3350 (NH, indoline), 2805, 2748 (Bohlmann bands), 1686 (CO), 1603, 762 (aromatics); ^1H -nmr (δ , ppm) 0.92 (3H, t, $J = 7.5$ Hz, CH_2CH_3), 1.93 (2H, q, CH_2CH_3), 2.55 (1H, d, $J_{12a,12b} = 8.0$ Hz, C12b-H), 1.1-2.9 (12H, m, $\text{C2-H}_2 + \text{C3-H}_2 + \text{C4-H}_2 + \text{C7-H}_2 + \text{C13-H}_2 + \text{C14-H}_2$), 3.0-3.5 (2H, m, C6-H₂), 3.93 (1H, br d, NH), 4.16 (1H, dd, $J_{12a,\text{NH}} = 4.0$ Hz, C12a-H), 6.6 - 7.2 (4H, m, aromatics); ^{13}C -nmr (δ , ppm) 7.9 (CH_2CH_3), 21.2 (C3), 29.3 (CH_2CH_3), 29.7 (C13), 31.8 (C7), 34.9 (C2), 37.8 (C14), 40.0 (C1), 47.7 (C6), 56.1 (C4), 57.4 (C7a), 65.5 (C12a), 69.0 (C12b), 110.9 (C11), 119.6 (C9), 124.2 (C8), 128.3 (C10), 129.6 (C7b), 152.3 (C11a), 208.8 (C15); Calc. for $\text{C}_{20}\text{H}_{26}\text{N}_2\text{O}$ (310.43) C, 77.38; H, 8.44; N, 9.03. Found C, 77.42; H, 8.40; N, 9.05. The mother liquor was concentrated and subjected to column chromatography ($\text{EtOAc-Et}_2\text{NH}$ 9:1 v/v, R_f 0.4) to give 3 (2.3 g, 18 %); mp 138-140 °C (ethanol); $[\alpha]_{\text{D}}^{25} -443.4^\circ$ ($c = 1.0$, CHCl_3); ms (m/z , %) 392 (M^+ , 7), 336 (10), 226 (7), 225 (7), 180 (100), 124 (71); ir (KBr), (ν , cm^{-1}) 2799, 2741 (Bohlmann bands), 1700 (CO, amide), 1602 (aromatics), 1262, 1147, 1130 (CF_3), 752 (aromatics); ^1H -nmr (δ , ppm) 0.87 (3H, t, $J = 7.4$ Hz, CH_2CH_3), 2.74 (1H, d,

$J_{12a,12b} = 6.7$ Hz, C12b-H), 1.0-3.5 (16H, m, C2-H₂ + C3-H₂ + C4-H₂ + C6-H₂ + C7-H₂ + C13-H₂ + C14-H₂ + CH₂CH₃), 3.88 (1H, dq, $J_{gem} = 16.2$ Hz, $J_{H,F} = 9.2$ Hz, NCH_AH_BCF₃), 4.05 (1H, dq, $J_{H,F} = 10.0$ Hz, NCH_AH_BCF₃), 4.06 (1H, d, C12a-H), 6.65-7.25 (4H, m, aromatics); ¹³C-nmr (δ, ppm) 7.5 (CH₂CH₃), 20.5 (C3), 26.8 (CH₂CH₃), 28.9 (C13), 32.9 (C7), 35.3 (C2), 37.1 (C14), 40.6 (C1), 47.7 (C6), 49.8 ($^2J_{C,F} = 30.2$ Hz, NCH₂CF₃), 56.3 (C4), 57.5 (C7a), 69.2 (C12b), 69.4 (C12a), 110.5 ($^5J_{C,F} = 1.9$ Hz, C11), 121.3 (C9), 124.9 (C8), 125.5 ($^1J_{C,F} = 287$ Hz, NCH₂CF₃), 128.7 (C10), 129.4 (C7b), 151.6 (C11a), 209.1 (C15); Calc. for C₂₂H₂₇F₃N₂O (392.46) C, 67.32; H, 6.94; F, 14.52; N, 7.14. Found C, 67.41; H, 6.92; F, 14.70; N, 7.15.

(-)-(1S:7aR:12aS:12bS)-1-Ethyl-15-(hydroxyimino)-1,2,3,4,6,7,7a,12,12a,12b-decahydro-1,7a-propanoindolo[2,3-a]quinolizine 4

A solution of **2** (8.0 g, 0.0258 mol) and NH₂OH.HCl (9.0 g, 0.129 mol) in anhydrous pyridine (70 ml) was refluxed for 24 h then the solvent was removed in *vacuo*. The residue was dissolved in a mixture of CHCl₃ (200 ml) and water (100 ml), then treated with saturated Na₂CO₃ solution. The organic layer was separated, the aqueous phase was extracted with CHCl₃ (2x100 ml). The combined extracts were dried over MgSO₄ and evaporated in *vacuo*. The residue was purified by column chromatography (toluene-Et₂NH 10:1 v/v) to give **4** (3.2 g, 38 %). Light yellow oil; $[\alpha]_D^{25} -382.3^\circ$ (c = 1.0, CHCl₃); ms (m/z, %) 325 (M⁺, 22), 308 (22), 307 (30), 306 (22), 267 (100), 195 (69); ir (CHCl₃) (ν, cm⁻¹) 3587, 3260 (OH), 3389 (NH), 1607 (aromatics), 935 (NO), 751 (aromatics); ¹H-nmr (δ, ppm) 0.89 (3H, t, J = 7.5 Hz, CH₂CH₃), 1.0-3.4 (17H, m, C2-H₂ + C3-H₂ + C4-H₂ + C6-H₂ + C7-H₂ + C13-H₂ + C14-H₂ + C12b-H), 3.76 (1H, br s, NH), 4.05 (1H, br d, J = 7.2 Hz, C12a-H), 5.5 (1H, br, N-OH), 6.6-7.2 (4H, m, aromatics); ¹³C-nmr (δ, ppm) 7.8 (CH₂CH₃), 20.0 (C3), 20.2 (C14), 30.1^x (CH₂CH₃), 30.2^x (C13), 34.1 (C7), 35.7 (C2), 39.4 (C1), 47.8 (C6), 50.5 (C7a), 55.9 (C4), 67.5 (C12b), 68.4 (C12a), 110.7 (C11), 120.4 (C9), 124.3 (C8), 127.6 (C10), 134.0 (C7b), 150.7 (C11a), 165.6 (C15); Calc. for C₂₀H₂₇N₃O (325.45) C, 73.81; H, 8.36; N, 12.91. Found C, 73.64; H, 8.27; N, 12.95.

(-)-(1S:7aR:12aS:12bS)-1-Ethyl-15 α -hydroxy-1,2,3,4,6,7,7a,12,12a,12b-decahydro-1,7a-propanoindolo[2,3-a]quinolizine 5

(-)-(1S:7aR:12aS:12bS)-1-Ethyl-15 β -hydroxy-1,2,3,4,6,7,7a,12,12a,12b-decahydro-1,7a-propanoindolo[2,3-a]quinolizine 6.

LiAlH_4 (1.0 g, 0.0264 mol) was suspended in THF (100 ml) in argon atmosphere. A solution of **2** (1.0 g, 0.0032 mol) in THF (25 ml) was added dropwise at reflux temperature within 30 min. The reaction mixture was refluxed for one more hour, then cooled and the excess LiAlH_4 was decomposed with water (1 ml), 15 % NaOH (1 ml) and water (3 ml). The precipitated solids were filtrated and washed with CHCl_3 (50 ml). The combined filtrates and washings were dried over MgSO_4 and evaporated in *vacuo*. The residue was separated by column chromatography (cyclohexane-EtOAc 1:1 v/v). The fraction with higher retention factor (R_f 0.38) was evaporated and crystallized from EtOH to give **5** (0.54 g, 53.6 %); mp 123-126 °C (ethanol); $[\alpha]_D^{25}$ -165.1 ° (c = 1.0, CHCl_3); ms (m/z , %), 312 (M^+ , 32), 311 (5), 255 (12), 182 (100); ir (CHCl_3), (ν , cm^{-1}) 3570 (OH), 3382 (NH), 1024 (C-OH); ^1H -nmr (δ , ppm) 0.87 (3H, t, J = 7.5 Hz, CH_2CH_3), 1.78 (2H, q, CH_2CH_3), 2.49 (1H, d, $J_{12a,12b}$ = 8.0 Hz, C12b-H), 1.0-2.9 (14H, m, C2-H₂ + C3-H₂ + C4-H₂ + C6-H₂ + C7-H₂ + C13-H₂ + C14-H₂), 3.5 (2H, br, OH + NH), 4.02 (1H, d, C12a-H), 4.45 (1H, dd, J = 9.6 Hz, 1 Hz, C15-H), 6.6-7.2 (4H, m, aromatics); ^{13}C -nmr (δ , ppm) 8.2 (CH_2CH_3), 22.2 (C3), 30.6 (C13), 31.6 (CH_2CH_3), 32.0 (C7), 34.5 (C14), 35.3 (C2), 39.5 (C1), 48.3 (C6), 50.4 (C7a), 56.3 (C4), 66.6 (C12a), 68.9 (C12b), 79.7 (C15), 110.5 (C11), 119.1 (C9), 126.2 (C8), 127.9 (C10), 132.1 (C7b), 152.6 (C11a); Calc. for $\text{C}_{20}\text{H}_{28}\text{N}_2\text{O}$ (312.44) C, 76.88; H, 9.03; N, 8.97. Found C, 77.24; H, 8.76; N, 8.96.

The fraction with lower retention factor (R_f 0.22) was evaporated and crystallized from EtOH to give **6** (0.28 g, 27.8 %); mp 206-209 °C (ethanol); $[\alpha]_D^{25}$ -138.1 ° (c = 1.0, CHCl_3); ms (m/z , %) 312 (M^+ , 17), 311 (15), 182 (100); ir (CHCl_3), (ν , cm^{-1}) 3392 (NH), 3170 (OH), 1090 (C-OH); ^1H -nmr (δ , ppm) 0.89 (3H, t, J = 7.4 Hz, CH_2CH_3), 1.80 and 1.90 (2H, dq, CH_2CH_3), 2.51 (1H, d, $J_{12a,12b}$ = 8.0 Hz, C12b-H), 1.0-3.2 (14H, m, C2-H₂ + C3-H₂ + C4-H₂ + C6-H₂ + C7-H₂ + C13-H₂ + C14-H₂), 3.5 (1H, br, NH), 3.55 (1H, br d, J = 6.8 Hz, C15-H), 3.92 (1H, d, C12a-H), 6.25 (1H, br, C15-OH), 6.0-7.15 (4H, m, aromatics); ^{13}C -nmr (δ , ppm) 8.2 (CH_2CH_3), 22.3 (C3), 26.4 (C13), 27.4 (C7), 31.2 (CH_2CH_3), 31.3 (C14), 35.0 (C2), 40.7 (C1), 47.9 (C6), 51.4 (C7a), 55.6 (C4), 66.4 (C12a), 69.6 (C12b), 74.8 (C15), 110.4 (C11), 120.0

(C9), 123.5 (C8), 127.3 (C10), 137.2 (C7b), 150.7 (C11a); Calc. for $C_{20}H_{28}N_2O$ (312.44) C, 76.88; H, 9.03; N, 8.97. Found C, 76.77; H, 8.87; N, 8.91.

(-)-6 α -Acetoxy-6 $\alpha\alpha$ -(2-acetoxyethyl)-11-acetyl-1,2,3,4,6,6a,11a,11b α -octahydro-1 β ,6 β -ethanoindolizino[1,2-b]indole (7)

A solution of **2** (1.0 g, 3.22 mmol) in Ac_2O (10 ml) and $AcOH$ (1 ml) was refluxed for 3 h, then poured onto ice (50 g). The aqueous mixture was extracted with $CHCl_3$ (3 x 50 ml), the combined extracts were neutralized by 5 % $NaHCO_3$, then washed with water, dried over $MgSO_4$ evaporated in *vacuo*. The residue was crystallized from $EtOH$ to give **7** (1.3 g, 88.8 %); mp 223-225 °C (ethanol); $[\alpha]_D^{25} -18^\circ$ (c=1.0, $CHCl_3$); ms (m/z, %) 454 (M^+ , 19), 411 (18), 209 (28), 167 (100), 152 (20), 139 (11), 138 (21); ir (KBr), (ν , cm^{-1}) 1738 (two CO, esters), 1671 (CO, amide), 1596 (aromatics), 1248 (two COC, esters), 763 (aromatics); 1H -nmr (δ , ppm) 0.46 (1H, dddd, $J_{gem} = 15.0$ Hz, $J_{12A,13A} = 11.7$ Hz, $J_{12A,13B} = 8.2$ Hz, $J_{12A,2\alpha} = 2.3$ Hz, C12- H_A), 0.68 (3H, t, $J = 7.5$ Hz, CH_2CH_3), 1.17 (1H, dq, $J_{gem} = 13.5$ Hz, $CH_AH_BCH_3$), 1.22 (1H, dddd, $J_{12B,13A} = 8.5$ Hz, $J_{12B,13B} \sim 1$ Hz, $J_{12B,11b} = 1.7$ Hz, C12- H_B), 1.30 (1H, dq, $CH_AH_BCH_3$), 1.37 (1H, dddd, $J_{gem} = 13$ Hz, $J_{2\alpha,3\alpha} = 5.8$ Hz, $J_{2\alpha,3\beta} = 13$ Hz, C2- H_α), 1.45 (1H, m, C3- H_A), 1.73 (1H, m, C2- H_β), 1.78 (1H, m, C13- H_A), 1.83 (1H, m, C3- H_B), 1.88 (3H, s, $OCOCH_3$), 1.98 (1H, dt, $J_{gem} = 13.9$ Hz, $J_{vic} = 6.9$ Hz, C14- H_A), 2.12 (1H, dt, C14- H_B), 2.20 (3H, s, $OCOCH_3$), 2.39 (3H, s, $NCOCH_3$), 2.68 (1H, ddd, $J_{gem} = 14$ Hz, C13- H_B), 2.94 (1H, ddd, $J_{gem} = 15.0$ Hz, $J_{4\alpha,3\alpha} = 5.8$ Hz, $J_{4\alpha,3\beta} = 13.2$ Hz, C4- H_α), 3.36 (1H, ddd, $J_{vic} = 6.3$ and ~ 1 Hz, C4- H_β), 3.39 (1H, dd, $J_{11a,11b} = 6.7$ Hz, C11b-H), 3.81 (1H, dt, $J_{gem} = 11.6$ Hz, $J_{vic} = 6.9$ Hz, C15- H_A), 3.82 (1H, dt, C15- H_B), 4.73 (1H, d, C11a-H), 7.11 (1H, dd, $J_{7,8} = 7.7$ Hz, $J_{8,9} = 7.4$ Hz, C8-H), 7.28 (1H, ddd, $J_{9,10} = 8.0$ Hz, $J_{7,9} = 1.0$ Hz, C9-H), 7.92 (1H, dd, C7-H), 8.14 (1H, d, C10-H); ^{13}C -nmr (δ , ppm) 7.2 (CH_2CH_3), 20.8 ($OCOCH_3$), 22.0 (C3), 22.1 ($OCOCH_3$), 24.3 ($NCOCH_3$), 28.6 (C12), 29.8 (C13), 30.8 (CH_2CH_3), 32.2 (C1), 36.1 (C2), 36.8 (C14), 42.6 (C4), 59.8 (C6a), 60.8 (C15), 65.2 (C11b), 69.0 (C11a), 100.5 (C6), 116.8 (C10), 124.3 (C8), 126.9 (C7), 128.7 (C9), 131.9 (C6b), 144.6 (C10a), 168.8 ($OCOCH_3$), 169.4 ($NCOCH_3$), 170.6 ($OCOCH_3$); (Products **7** and **8** were shown by nmr spectra to occur as ca. 60:40 mixtures of rotational isomers due to the restricted rotation of the N-acetyl group. The data reported here refer to the major component in which the carbonyl oxygen is pointing toward the aromatic ring.) Calc. for $C_{26}H_{34}N_2O_5$ (454.55) C, 68.70; H, 7.54; N, 6.16. Found C, 68.55; H, 7.52;

N, 6.15.

(-)-6 α -Hydroxy-6 α -(2-hydroxyethyl)-11-acetyl-1,2,3,4,6,6a,11a α ,11b α -octahydro-1 β ,6 β -ethanoindolizino[1,2-b]indole 8

(-)-1-Acetyl-4a β -ethyl-7a α -(2-hydroxyethyl)-1,2,3,4,4a,5,6,7,7a,12,12a α ,12b β -dodecahydropyrido[3,2':6,7]cyclohept[1,2-b]indole 9

A solution of 7 (8.0 g, 0.0176 mol) and NaOH (16.0 g, 0.4 mol) in 96 % EtOH (700 ml) was refluxed for 2.5 h. The solvent was removed in *vacuo*, the residue was diluted with water (500 ml) and extracted with CHCl₃ (3x200 ml). The organic layer was dried over MgSO₄ and evaporated in *vacuo*. The residue was separated by column chromatography on Al₂O₃ (CH₂Cl₂:EtOH 10:1 v/v). The fraction with lower retention factor (R_f 0.66) was evaporated and crystallized from ethanol to give 8 (2.6 g, 40.5 %); mp 207-209 °C (ethanol); $[\alpha]_D^{25}$ -96.4 ° (c = 1.0, CHCl₃); ms (m/z , %) 370 (M^+ , 57), 167 (48), 162 (13), 161 (14), 152 (38), 139 (30), 138 (51), 130 (36), 111 (100); ir (CHCl₃), (ν , cm⁻¹) 3400 (OH), 3170 (OH), 1656 (CO, amide), 1394 (OH), 1189 (C-O), 1064 (C-O), 600 (OH); ¹H-nmr (δ , ppm) 0.57 (1H, m, C12-H_A), 0.72 (3H, t, J = 7.5 Hz, CH₂CH₃), 1.22 (1H, m, C12-H_B), 1.25 (1H, dq, CH_AH_BCH₃), 1.33 (1H, m, C2-H_A), 1.42 (1H, dq, CH_AH_BCH₃), 1.43 (1H, m, C3-H_A), 1.54 (1H, ddd, J_{gem} = 15 Hz, $J_{14A,15A}$ = 2.0 Hz, $J_{14A,15B}$ = 2.7 Hz, C14-H_A), 1.68 (1H, ddd, J_{gem} = 14.5 Hz, J_{vic} = 8.2 and ~ 1 Hz, C13-H_A), 1.76 (1H, ddd, J_{gem} = 13 Hz, $J_{2\beta 3\alpha}$ = 1 Hz, $J_{2\beta 3\beta}$ = 5.5 Hz, C2-H_B), 1.81 (1H, m, C13-H_B), 1.87 (1H, m, C3-H_B), 2.38 (3H, s, NCOCH₃), 2.53 (1H, ddd, $J_{14B,15A}$ = 5.0 Hz, $J_{14B,15B}$ = 11.6 Hz, C14-H_B), 2.82 (1H, ddd, J_{gem} = 14.8 Hz, $J_{3\alpha,4\alpha}$ = 5.2 Hz, $J_{3\beta,4\alpha}$ = 12.8 Hz, C4-H_A), 3.38 (1H, dd, $J_{11a,11b}$ = 7.0 Hz, $J_{11b,12B}$ = 1.4 Hz, C11b-H), 3.39 (1H, ddd, $J_{3\alpha,4\beta}$ = 1 Hz, $J_{3\beta,4\beta}$ = 6.2 Hz, C4-H_B), 3.68 (1H, ddd, J_{gem} = 12.4 Hz, C15-H_A), 3.84 (1H, ddd, C15-H_B), 4.80 (1H, d, C11a-H), 4.55 and 5.3 (1H + 1H, br lines, 2xOH), 7.07 (1H, dd, $J_{7,8}$ = 7.5 Hz, $J_{8,9}$ = 7.0 Hz, C8-H), 7.20 (1H, dd, $J_{7,9}$ = 1 Hz, C7-H), 7.25 (1H, ddd, $J_{9,10}$ = 8.0 Hz, C9-H), 8.12 (1H, d, C10-H); ¹³C-nmr (δ , ppm) 7.30 (CH₂CH₃), 21.5 (C3), 24.4 (NCOCH₃), 27.9 (C12), 31.3 (C13 + CH₂CH₃), 32.2 (C1), 35.6 (C2), 38.5 (C14), 41.1 (C4), 57.7 (C15), 59.0 (C6a), 65.2 (C11b), 65.7 (C11a), 91.2 (C6), 117.3 (C10), 124.0 (C8), 124.1 (C7), 128.2 (C9), 135.3 (C6b), 143.2 (C10a), 169.7 (NCOCH₃); Calc. for C₂₂H₃₀N₂O₃ (370.48) C, 71.32; H, 8.16; N, 7.56. Found C, 71.42; H, 8.23; N, 7.49.

The fraction with higher retention factor (R_f 0.79) was evaporated and crystallized

from EtOH to give 9 (3.1 g, 47.5 %); mp 204-206 °C (ethanol); $[\alpha]_D^{25}$ -193 ° (c = 1.0, CHCl₃); ms (m/z, %) 370 (M⁺, 12), 161 (100), 153 (27), 130 (66); ir (CHCl₃), (ν, cm⁻¹) 3630 (OH), 3538 (OH, bridged) 3390 (NH, indoline), 1702 (CO), 1643 (CO, amide), 1607, 753 (aromatics); ¹H-nmr (δ, ppm) 0.78 (3H, t, J = 7.5 Hz, CH₂CH₃), 1.24 (1H, dq, J_{gem} = 14.3 Hz, CH_AH_BCH₃), 1.48 (3H, s, NCOCH₃), 1.45-1.55 (2H, m, C3-H₂), 1.51 (1H, m, C4-H_α), 1.66 (1H, dqd, J_{HB,5α} = 1.3 Hz, CH_AH_BCH₃), 1.72 (1H, m, C4-H_β), 1.76 (1H, ddd, J_{gem} = 15.7 Hz, J_{5β,6α} = 4.2 Hz, J_{5β,6β} = 4.2 Hz, C5-H_β), 1.95 (1H, dddd, J_{5α,6α} = 4.3 Hz, J_{5α,6β} = 14.1 Hz, C5-H_α), 2.05 (1H, ddd, J_{gem} = 13.9 Hz, J_{13A,14A} = 4.7 Hz, J_{13A,14B} = 6.0 Hz, C13-H_A), 2.28 (1H, ddd, J_{gem} = 13.2 Hz, J_{2α,3α} = 4.5 Hz, J_{2α,3β} = 10.2 Hz, C2-H_α), 2.35 (1H, ddd, J_{13B,14A} = 7.6 Hz, J_{13B,14B} = 5.1 Hz, C13-H_B), 2.60 (1H, ddd, J_{gem} = 18.0 Hz, C6-H_β), 2.83 (1H, br, OH), 2.86 (1H, ddd, C6-H_α), 3.21 (1H, d, J_{12a,12b} = 9.9 Hz, C12b-H), 3.46 (1H, dddd, J_{gem} = 11.6 Hz, J_{14A,OH} = 2.5 Hz, C14-H_A), 3.59 (1H, dddd, J_{14B,OH} = 6.0 Hz, C14-H_B), 3.84 (1H, s, NH), 4.34 (1H, d, C12a-H), 4.73 (1H, br dd, J_{vic} = 2.6 Hz, C2-H_β), 6.62 (1H, ddd, J_{10,11} = 7.9 Hz, J_{9,11} = 1.0 Hz, J_{8,11} = 0.6 Hz, C11-H), 6.82 (1H, ddd, J_{8,9} = 7.6 Hz, J_{9,10} = 7.4 Hz, C9-H), 7.09 (1H, ddd, J_{8,10} = 1.3 Hz, C10-H), 7.40 (1H, ddd, C8-H); ¹³C-nmr (δ, ppm) 6.9 (CH₂CH₃), 20.2 (C3), 21.6 (NCOCH₃), 25.0 (CH₂CH₃), 27.8 (C5), 29.4 (C4), 36.9 (C6), 38.3 (C2), 38.7 (C4a), 40.5 (C13), 59.5 (C14), 60.9 (C12a), 62.3 (C12b), 65.0 (C7a), 109.5 (C11), 119.5 (C9), 126.5 (C7b), 128.7 (C8), 129.0 (C10), 148.6 (C11a), 172.0 (NCOCH₃), 210.0 (C7); Calc. for C₂₂H₃₀N₂O₃ (370.48) C, 71.32; H, 8.16; N, 7.56. Found C, 71.49; H 8.07; N, 7.61.

Refluxing of either pure 8 or pure 9 in ethanol for 12 hours, after evaporation and chromatographic separation (on Al₂O₃ column, CH₂Cl₂:EtOH 10:1 v/v) affords the same product mixture in which the ratio of 8:9 is approx. 2.2:1.

ACKNOWLEDGEMENTS

The authors are grateful to Drs J. Tamás and G. Keresztury (Central Research Institute for Chemistry) for the recording and interpretation of the ms and ir spectra, respectively. Financial support from the Hungarian Academy of Sciences and Gedeon Richter Pharmaceutical Works (Budapest) are greatly acknowledged.

REFERENCES

1. For part XLII. see J. Sápi, L. Szabó, E. Baitz-Gács, Gy. Kalaus, and Cs. Szántay, Tetrahedron, in press.
2. Zs. Kardos-Balogh, F. Sóti, M. Incze, M. Kajtár-Peredy, and Cs. Szántay, Tetrahedron Letters, 1985, 26, 5593.
3. A. Bax, Two Dimensional Nuclear Magnetic Resonance in Liquids, Delft University Press, 1982.
4. M. Incze, F. Sóti, Zs. Kardos-Balogh, and Cs. Szántay, Heterocycles, 1985, 23, 2843.
5. T. Irie, and H. Tanida, J. Org. Chem., 1980, 45, 1795.
6. D. Beke, Advances in Heterocyclic Chemistry: Heterocyclic Pseudobases, Vol. 1, ed. by A.R. Katritzky, Academic Press, Inc., London, 1963, pp 167.

Received, 29th July, 1988