

DEGRADATION AND MANIPULATIONS OF THE IMMUNOSUPPRESSANT FK506: PREPARATION OF POTENTIAL SYNTHETIC INTERMEDIATES

Robert S. Coleman¹ and Samuel J. Danishefsky*
 Department of Chemistry
 Yale University
 New Haven, Connecticut 06511 USA

Abstract - Degradation of the immunosuppressant FK506 is described.


The fermentation product FK506 (**1**), isolated from *Streptomyces tsukubaensis* no. 9993 by a group of Fujisawa scientists, has stimulated a great deal of research.²⁻⁴ Not the least source of interest in FK506 arises from its potent immunosuppressive powers. From a chemical viewpoint, the novel 23-membered macrolide structure with its varied menu of appendages and stereogenic centers provides a great deal of challenge to any program in total synthesis. Of course, the ultimate goal in studying FK506 is to identify with precision the minimum structural domains that are required for its immunosuppressive activity. Such information might provide a better insight into the mode of action of this molecule, and, by inference, begin to clarify the steric and electronic character of its receptor(s).

Our laboratory is investigating several facets of the FK506 problem. An effort in total synthesis is being conducted in parallel with a degradative program. In this paper we describe some of our initial accomplishments in the latter area.

We first addressed one of the sources of the chemical instability of FK506, *i.e.*, the β -hydroxyketone sector (carbons 22-24).⁵ Reduction of **1** with Li(*s*-Bu)₃BH (THF, -78°C, 1.5 h) occurred selectively at C₂₂ to afford a single alcohol (**2**, Scheme 1, ~90% yield). There was little or no interference from competing reduction of either the C₉ or the C₁₀ masked ketone. A rigorous assignment of the stereochemistry at C₂₂ of this product is not offered.⁶ However, examination of molecular models based on the molecular geometry provided in the previously published crystal structure² suggested that attack of hydride would occur from the β -face of the molecule (as drawn), to provide a product with the 22*S* stereochemistry. Treatment of **2** with lead tetraacetate (1.5-2.0 equiv) in methanol (0-25°C) afforded the amide diester **3** (52%, 33% recovered **2**). The

We dedicate this paper to Professor Derek H. R. Barton on the occasion of his 70th birthday. We are thankful for his continuing leadership in organic chemistry.


pipecolinate residue of **3** could be removed by treatment with lithium hydroxide (10 equiv, THF-H₂O, 25°C) followed by acidification (10% HCl, pH 1, 25°C, 12 h), thereby providing the lactone tetraol **5**⁷ (65%).




Scheme 1

It was of interest to determine the stereochemical outcome when reduction of the C₂₂ ketone was carried out on a *seco* variation of FK506. Treatment of **1** with lead tetraacetate in methanol afforded **4**, (81%), which upon reduction with Li(*s*-Bu)₃BH (THF, -78°C) gave rise to **3** as the only product (Scheme 1). The fact that the two routes provided the same reduction product suggests, but does not prove, that the local environment in the area of the C₂₂ ketone is similar in **1** and **4**.

The oxalylpipecolinate residue could be excised and retrieved from the molecule. This degradative sequence (Scheme 2) began with treatment of **2** with lithium hydroxide (1.1 equiv, THF-H₂O, 25°C) followed by acidification (Dowex 15W, H⁺ form). Esterification of the resulting carboxylic acid with diazomethane (ether, 0°C) led, in low yield (25%), to a substance formulated as **6**, the methyl ester of the 22-dihydro-*seco*-acid corresponding to FK506. Treatment of **6** with lead tetraacetate in methanol afforded **5** and **7**.⁸



We further investigated selective differentiation of the C₂₂ oxygen by exploiting its proximal allyl group (Scheme 3). Toward this end, compound 2 was treated with *N*-iodosuccinimide in CH₂Cl₂ (1.5-2.0 equiv, 25°C, 2-4 days). A very slow reaction was noted. We were unable to drive the process to completion under more forcing conditions. Under the conditions described above, 37% of a single iodoether 8 was obtained (36% recovered 2; stereochemistry at C₃₇ not rigorously assigned). When this compound was subjected to the sequence: (i) Pb(OAc)₄, MeOH, 52%; (ii) LiOH, THF-MeOH-H₂O (3:2:1), 25°C, 89%; (iii) acetone, CuSO₄, CSA, 76%; the lactone acetonide 10⁹ was produced. Ozonolysis of 10 (-78°C, CH₂Cl₂; Me₂S, 25°C) afforded 11¹⁰ and 12.¹¹ The results of additional degradative and synthetic studies will be disclosed in due course.



ACKNOWLEDGMENTS We wish to thank the National Institutes of Health (PHS grant AI 16943) for financial support of this work. Nmr spectra were obtained through the auspices of the Northeast Regional NSF/NMR facility at Yale University, which was supported by NSF Chemistry Division Grant CHE 7916210. We extend our sincere appreciation to Dr. Roland Wenger of Sandoz, Ltd. (Basel, Switzerland) for the generous gift of FK506.

REFERENCES AND NOTES

1. National Institutes of Health (National Cancer Institute) Postdoctoral Fellow, 1988-1990 (CA 08388).
2. H. Tanaka, A. Kuroda, H. Marusawa, H. Hatanaka, T. Kino, T. Goto, M. Hashimoto, and T. Taga, *J. Am. Chem. Soc.*, **1987**, 109, 5031.
3. T. Kino, H. Hatanaka, M. Hashimoto, M. Nishiyama, T. Goto, M. Okuhara, M. Kohsaka, H. Aoki, and H. Imanaka, *J. Antibiot.*, **1987**, 60, 1249. T. Kino, H. Hatanaka, S. Miyata, N. Inamura, M. Nishiyama, T. Yajima, T. Goto, M. Okuhara, M. Kohsaka, H. Aoki, and T. Ochiai, *J. Antibiot.*, **1987**, 60, 1256.
4. D. Askin, R. P. Volante, R. A. Reamer, K. M. Ryan, and I. Shinkai, *Tetrahedron Lett.*, **1988**, 29, 277. S. Mills, R. Desmond, R. A. Reamer, R. P. Volante, and I. Shinkai, *Tetrahedron Lett.*, **1988**, 29, 281.
5. Treatment of FK506 with 1 N NaOH is reported to cause cleavage of the C₂₃-C₂₄ and C₂₅-C₂₆ carbon-carbon bonds via a retroaldol process: see reference 2. We were unable to selectively hydrolyze the macrolactone ester linkage of FK506 using stoichiometric LiOH (THF-MeOH-H₂O, 0 °C).
6. Analysis and assignment of the ¹H nmr spectra of FK506 and related cyclic intermediates are severely complicated by the presence of rotamers involving the pipecolinic acid residue: see also reference 2.
7. Compound **5** was characterized: ¹H Nmr (CDCl₃, 500 MHz) δ 5.78 (dd, 1H, J = 9.2, 1.3 Hz), 5.09 (d, 1H, J = 9.9 Hz), 5.04 (dd, 1H, J = 17.0, 2.0 Hz), 4.97 (dd, 1H, J = 10.1, 2.1 Hz), 4.39 (m, 2H), 4.08 (dd, 1H, J = 7.9, 1.6 Hz), 3.86 (br m, 1H), 3.70 (ddd, 1H, J = 11.1, 7.8, 4.5 Hz), 3.50 (ddd, 1H, J = 7.2, 5.5, 1.6 Hz), 3.42 (s, 3H), 3.41 (s, 3H), 3.39 (s, 3H), 3.02 (ddd, 1H, J = 11.2, 8.7, 4.1 Hz), 2.70 (br s, 1H), 2.48 (m, 1H), 2.41 (m, 1H), 2.31 (m, 3H), 2.13 (m, 2H), 2.01 (m, 2H), 1.94 (m, 1H), 1.84 (m, 1H), 1.75-0.95 (m, 14H), 1.61 (d, 3H, J = 1.2 Hz), 1.55 (d, 3H, J = 0.3 Hz), 1.29 (d, 3H, J = 7.0 Hz), 1.89 (m, 3H), 0.60 (d, 3H, J = 7.1 Hz).

8. Compound **7**, which consisted of major and minor rotameric forms (*ca.* 2:1), was identical (by ^1H nmr and SiO_2 TLC comparison) with synthetic material prepared from L-pipecolinic acid. Compound **7** was characterized: ^1H Nmr (CDCl_3 , 490 MHz) δ 5.25 (d, 1H, $J = 5.7$ Hz, major), 4.54 (d, 1H, $J = 5.2$ Hz, minor), 4.46 (br d, 1H, $J = 12.7$ Hz, minor), 3.89 (s, 3H, major), 3.85 (s, 3H, minor), 3.78 (s, 3H, minor), 3.77 (s, 3H, major), 3.58 (br d, 1H, $J = 11.6$ Hz, major), 3.34 (td, 1H, $J = 13.2, 3.1$ Hz, major), 2.91 (td, 2H, $J = 13.0, 3.5$ Hz, minor), 2.31 (br d, 1H, $J = 14.1$ Hz, major), 2.28 (br d, 1H, $J = 13.7$ Hz, minor), 1.8-1.3 (m, 5H); EIMS, m/e (relative intensity) 229 (M^+ , 16), 170 (base), 142 (60), 110 (7), 82 (13), 70 (23).
9. Compound **10** was characterized: ^1H Nmr (CDCl_3 , 490 MHz) δ 5.27 (d, 1H, $J = 9.0$ Hz), 5.09 (d, 1H, $J = 10.0$ Hz), 4.25-4.15 (m, 4H), 4.07 (dd, 1H, $J = 7.9, 1.7$ Hz), 3.70 (ddd, 1H, $J = 11.1, 7.9, 4.5$ Hz), 3.5 (ddd, 1H, $J = 8.0, 6.1, 1.7$ Hz), 3.43 (m, 1H), 3.41 (s, 3H), 3.40 (s, 3H), 3.39 (s, 3H), 3.27 (dd, 1H, $J = 9.8, 5.1$ Hz), 3.22 (dd, 1H, $J = 9.8, 6.7$ Hz), 3.07 (m, 1H), 3.01 (ddd, 1H, $J = 11.3, 8.8, 4.2$ Hz), 2.66 (br s, 1H), 2.49 (m, 1H), 2.31 (m, 2H), 2.12 (dd, 1H, $J = 12.7, 3.8$ Hz), 2.01 (dt, 2H, $J = 12.7, 4.3$ Hz), 1.94 (dd, 2H, $J = 7.0, 5.3$ Hz), 1.80-0.85 (m, 12H), 1.60 (d, 3H, $J = 1.2$ Hz), 1.54 (d, 3H, $J = 1.0$ Hz), 1.46 (s, 3H), 1.41 (s, 3H), 1.29 (d, 3H, $J = 7.0$ Hz), 0.84 (d, 3H, $J = 6.2$ Hz), 0.65 (d, 3H, $J = 6.8$ Hz).
10. Compound **11** was characterized: ^1H Nmr (CDCl_3 , 490 MHz) δ 9.81 (d, 1H, $J = 2.4$ Hz), 4.46 (ddd, 1H, $J = 10.3, 5.8, 4.5$ Hz), 4.32 (d, 1H, $J = 2.7$ Hz), 4.19 (m, 2H), 3.27 (d, 2H, $J = 5.6$ Hz), 3.13 (m, 1H), 2.54 (ddd, 1H, $J = 13.3, 7.2, 2.8$ Hz), 2.17 (s, 3H), 1.94-1.81 (m, 3H), 1.46 (s, 3H), 1.47 (s, 3H), 0.80 (d, 3H, $J = 6.8$ Hz).
11. Compound **12** was characterized: ^1H Nmr (CDCl_3 , 490 MHz) δ 4.11 (dd, 1H, $J = 8.0, 1.7$ Hz), 3.68 (ddd, 1H, $J = 11.1, 8.0, 4.5$ Hz), 3.46 (ddd, 1H, $J = 7.8, 6.1, 1.7$ Hz), 3.41 (s, 3H), 3.40 (s, 3H), 2.50 (m, 2H), 2.32 (m, 2H), 2.14 (s, 3H), 2.11 (m, 1H), 1.7-1.5 (m, 3H), 1.30 (d, 3H, $J = 7.0$ Hz), 0.95 (d, 3H, $J = 6.6$ Hz).

Received, 29th August, 1988