

REINVESTIGATION OF CYCLOADDITION OF NITROSOBENZENE TOWARDS  
1,3-DIPHENYLISOBENZOFURANS




Cheng-Tung Lin<sup>\*a</sup>, Li-Wei Din<sup>a,1</sup>, and Sue-Lein Wang<sup>b</sup>




a. Department of Chemistry, Tunghai University,  
Taichung, Taiwan 40098 R.O.C.

b. Department of Chemistry, National Tsing Hua University,  
Hsinchu, Taiwan 30042 R.O.C.


**Abstract** — The title reaction gives rearrangement product of *N*-phenyl- $\alpha$ -phenyl- $\alpha$ -(*o*-benzoylphenyl)nitrones (8) rather than the earlier reported Diels-Alder cycloadduct 2. The molecular structure of 8a has been confirmed by a single crystal X-ray diffraction study.

Nitrosobenzene is one of the carbon-nitroso compounds known to be a reactive heterodienophile in the Diels-Alder reaction.<sup>2</sup> Extensive studies of cycloaddition reveal that nitrosobenzene reacts regioselectively towards unsymmetrical dienes.<sup>3</sup> Earlier, Mustafa reported that nitrosobenzene cyclized with 1,3-diphenylisobenzofuran (1a) to afford the corresponding Diels-Alder cycloadduct 2a.<sup>4</sup> Several years ago, Kirby and his co-workers found that nitrosyl cyanate reacted with 1,3-diphenylisobenzofuran in a different manner. The major products were 1,2-dibenzoylbenzene and *N*-cyanoimine 3 in a ratio of 1 : 1.<sup>5</sup> The outcome of the reaction was rationalized by oxygen atom transfer from the transient intermediate, *N*-cyanonitrene 4 or *N*-cyanoxaziridine 5, to the reactant, 1,3-diphenylisobenzofuran and/or by the recombination of unreacted reactant with cyanonitrene, a fragment cleaved from the unstable Diels-Alder cycloadduct 6.





In connection with our studies on the Diels-Alder reaction of trichloronitrosomethane,<sup>6</sup> we undertook the reinvestigation of the reaction of nitrosobenzene with 1,3-diphenylisobenzofurans. In this paper, we wish to describe the results that are different with the ones reported earlier by Mustafa<sup>4</sup> and Kirby<sup>5</sup>.



a R = H  
b R = CH<sub>3</sub>

Treatment of nitrosobenzene with 1a in ethyl alcohol as reported<sup>4</sup> afforded a stable light yellow crystal in the yield of 64%. Its melting point, microanalysis and molecular ion of the mass spectrum were found to be identical with the data reported for cycloadduct 2.<sup>4</sup> However, the ir spectrum shows a strong absorption band at 1665 cm<sup>-1</sup> which can only be accounted by a conjugated C=O stretching. The presence of carbonyl group is further established by the signal at 194.0 ppm in its <sup>13</sup>C-nmr spectrum. Apparently, the structure of product was assigned incorrectly by earlier investigator.<sup>4</sup> Due to the absence of critical evidence for distinguishing between two possible structures 7 and 8a, an X-ray structure analysis was taken. The crystal structure<sup>7</sup> reveals a notable C-N-O angle of 124.1° and an N-C(14) length of 1.311 Å (Tables 1 and 2) which should exclude the consideration of N-phenyloxaziridine 7 and unambiguously established the structure of product as nitronone 8a.<sup>8</sup> A view of molecule 8a is given in Figure 1.

Under the similar conditions, the reaction of nitrosobenzene with 5,6-dimethyl-1,3-diphenylisobenzofuran (1b) exhibited similar results to afford 8b.<sup>9</sup> Attempts of isolation of the preliminary cycloadduct 2 by decreasing the reaction temperature to  $-78^{\circ}\text{C}$  were unsuccessful. Neither the reaction




Fig. 1 ORTEP diagram of the X-ray crystal structure of nitrone 8a

Table 1. Bond angles data for nitrone 8a

O(1)-N-C(14)	124.1(3)	O(1)-N-C(21)	114.4(3)
C(14)-N-C(21)	121.3(3)	C(2)-C(1)-C(6)	118.5(4)
C(2)-C(1)-C(7)	122.5(4)	C(6)-C(1)-C(7)	119.0(4)
C(1)-C(2)-C(3)	120.7(5)	C(2)-C(3)-C(4)	120.1(5)
C(3)-C(4)-C(5)	120.1(5)	C(4)-C(5)-C(6)	120.0(5)
C(1)-C(6)-C(5)	120.6(5)	O(2)-C(7)-C(1)	121.1(4)
O(2)-C(7)-C(8)	119.0(3)	C(1)-C(7)-C(8)	119.8(3)
C(7)-C(8)-C(9)	119.6(4)	C(7)-C(8)-C(13)	120.7(3)
C(9)-C(8)-C(13)	119.3(4)	C(8)-C(9)-C(10)	121.2(4)
C(9)-C(10)-C(11)	120.0(5)	C(10)-C(11)-C(12)	119.7(5)
C(11)-C(12)-C(13)	121.4(4)	C(8)-C(13)-C(12)	118.4(4)
C(8)-C(13)-C(14)	123.3(3)	C(12)-C(13)-C(14)	118.2(3)
N-C(14)-C(13)	119.6(3)	N-C(14)-C(15)	120.6(3)
C(13)-C(14)-C(15)	119.8(3)	C(14)-C(15)-C(16)	119.7(4)
C(14)-C(15)-C(20)	120.8(3)	C(16)-C(15)-C(20)	119.4(4)
C(15)-C(16)-C(17)	120.1(4)	C(16)-C(17)-C(18)	120.3(5)
C(17)-C(18)-C(19)	119.8(5)	C(18)-C(19)-C(20)	121.4(5)
C(15)-C(20)-C(19)	119.0(4)	N-C(21)-C(22)	118.7(3)
N-C(21)-C(26)	119.5(3)	C(22)-C(21)-C(26)	121.8(4)
C(21)-C(22)-C(23)	118.7(4)	C(22)-C(23)-C(24)	119.9(4)
C(23)-C(24)-C(25)	121.0(5)	C(24)-C(25)-C(26)	120.0(4)
C(21)-C(26)-C(25)	118.3(4)		

temperature (-78 °C to 80 °C) nor solvents (methylene chloride, ether and benzene) have influence on the formation and the yield of nitrone 8.

Table 2. Bond lengths data (Å) for nitrone 8a

O(1)-N	1.299 (4)	O(2)-C(7)	1.218 (5)
N-C(14)	1.311 (5)	N-C(21)	1.452 (5)
C(1)-C(2)	1.389 (6)	C(1)-C(6)	1.380 (7)
C(1)-C(7)	1.404 (5)	C(2)-C(3)	1.382 (6)
C(3)-C(4)	1.371 (8)	C(4)-C(5)	1.370 (8)
C(5)-C(6)	1.390 (7)	C(7)-C(8)	1.497 (6)
C(8)-C(9)	1.383 (6)	C(8)-C(13)	1.402 (5)
C(9)-C(10)	1.382 (7)	C(10)-C(11)	1.370 (7)
C(11)-C(12)	1.380 (7)	C(12)-C(13)	1.395 (6)
C(13)-C(14)	1.481 (5)	C(14)-C(15)	1.484 (5)
C(15)-C(16)	1.378 (6)	C(15)-C(20)	1.381 (6)
C(16)-C(17)	1.385 (7)	C(17)-C(18)	1.359 (8)
C(18)-C(19)	1.356 (8)	C(19)-C(20)	1.386 (7)
C(21)-C(22)	1.377 (6)	C(21)-C(26)	1.368 (6)
C(22)-C(23)	1.381 (7)	C(23)-C(24)	1.363 (7)
C(24)-C(25)	1.363 (7)	C(25)-C(26)	1.388 (6)

4-Nitrosopyridine-N-oxide, known for its "ene" reaction with tetramethylethylene<sup>10</sup> has not hitherto been employed as a dienophile. When 1,3-diphenylisobenzofuran was treated with 4-nitrosopyridine-N-oxide at 0 °C, the desired product 9 was isolated in the yield of 38 %. The structure of nitrone 9 was determined on the basis of spectroscopic analysis and high resolution mass spectrum.<sup>11</sup>

Attention was then turned to the examination of oxygen atom transfer between nitrone 8 and isobenzofurans as suggested by the mechanism proposed by Kirby et al. for the reaction of nitrosyl cyanate with 1.<sup>5</sup> Thus, refluxing the mixture of 1,3-diphenylisobenzofuran and nitrone 8a in benzene or toluene under oxygen-free nitrogen atmosphere for 24 hours did not give any products. All the reactants were chromatographically recovered. The nitrones 8 and 9 are relatively stable in the air for weeks, however, hydrolysis of 8a in aqueous acid solution produced 1,2-dibenzoylbenzene in quantitatively yield.

#### ACKNOWLEDGEMENT

This work was supported by grant from the National Science Council of the republic of China (NSC-76-0208-M029-03).

#### REFERENCES AND NOTES

1. Parts of this work were taken from the Master of Science thesis (chinese) of

- L. W. Din, Tunghai University, 1988
2. a). S. M. Weinreb and R. R. Staib, Tetrahedron, 1982, 38, 3087; b) G. W. Kirby, Chem. Soc. Rev., 1977, 6, 1; c) J. Hamer and M. Ahmad, "Organic Chemistry--A series of Monographs", Vol. 8, Academic Press, New York, 1967, Chapter 12.
  3. a) G. Kresze and W. Kosbahn, Tetrahedron, 1971, 27, 1931 b) G. Kresze and O. Korpiun, Tetrahedron, 1966, 22, 2493
  4. A. Mustafa, J. Chem. Soc., 1949, 256
  5. P. Horsewood, G. W. Kirby, R. P. Sharma, and J. G. Sweeny, J. Chem. Soc., Perkin 1, 1981, 1802
  6. a) C. T. Lin and J. H. Chen, J. Chin. Chem. Soc., 1987, 34, 317; b) We have found that trichloronitrosomethane reacted with 1,3-diphenylisobenzofurans to produce 1,2-dibenzoylbenzenes in high yield. The corresponding Diels-Alder cycloadducts were unable to be isolated. See also ref.1.
  7. Crystallographic data for  $C_{26}H_{19}NO_2$  (8a): orthorhombic, space group Pbca (No.61),  $a = 11.264(4) \text{ \AA}$ ,  $b = 16.418(3) \text{ \AA}$ ,  $c = 21.771(6) \text{ \AA}$ ,  $V = 4026(2) \text{ \AA}^3$ ,  $Z = 8$ ,  $FW = 377.4$ ,  $D_c = 1.245 \text{ g/cm}^3$ ,  $\lambda = 0.71073 \text{ \AA}$ . Of the 3414 reflections measured at room temperature on a Nicolet R3m/V diffractometer, 3023 were unique and 1470 reflections were considered observed ( $I > 3 \sigma(I)$ ) after LP corrections. The structure was solved by direct methods and refined by full matrix least-squares refinement.  $R = 0.042$ ,  $R_w = 0.043$ ,  $GOF = 1.42$ . Highest peak on final difference map =  $0.37 \text{ e/\AA}^3$ . All calculations were performed on a Micro VAX II based Nicolet SHELXTL PLUS system.
  8. 8a (64 % yield): mp  $227^\circ \text{C}$  (lit.<sup>4</sup>  $230^\circ \text{C}$ ),  $^1\text{H-nmr}$  ( $\text{CDCl}_3$ )  $\delta$  8.01 (d, 1 H,  $J = 8.1 \text{ Hz}$ ), 7.93 (d, 1 H,  $J = 8.4 \text{ Hz}$ ), 7.41 (m, 6 H), 7.19 (m, 11 H);  $^{13}\text{C-nmr}$  ( $\text{CDCl}_3$ )  $\delta$  194.0 (s), 147.0 (s), 146.7 (s), 139.8 (s), 136.2 (s), 135.2 (s), 134.5 (s), 132.4 (d), 131.3 (d), 130.7 (d), 130.6 (d), 130.3 (d), 128.8 (d), 128.7 (d), 128.6 (d, two peaks), 128.4 (d), 128.1 (d, two peaks), 124.3 (d); ir (KBr) 3030,  $1665 \text{ cm}^{-1}$ ; ms (12 eV)  $m/z$  377 ( $\text{M}^+$ ), 361 ( $\text{M}^+ - 16$ ), 270 ( $\text{M}^+ - \text{C}_6\text{H}_5\text{NO}$ , base peak); Anal. Calcd for  $C_{26}H_{19}NO_2$ : C, 82.74; H, 5.07; N, 3.71; Found: C, 82.80; H, 5.02; N, 3.73
  9. 8b (89 %): mp  $189^\circ \text{C}$ ;  $^1\text{H-nmr}$  ( $\text{CDCl}_3$ )  $\delta$  7.98 (d, 1 H,  $J = 7.2 \text{ Hz}$ ), 7.90 (d, 1 H,  $J = 7.7 \text{ Hz}$ ), 7.46 (m, 3 H), 7.15 (m, 11 H), 6.95 (s, 1 H), 2.25 (s, 3 H,  $\text{CH}_3$ ), 2.20 (s, 3 H,  $\text{CH}_3$ );  $^{13}\text{C-nmr}$  ( $\text{CDCl}_3$ )  $\delta$  194.2 (s), 147.6 (s), 147.1 (s), 139.2 (s), 137.2 (s), 136.6 (s), 135.2 (s), 132.1 (d), 132.1 (s), 132.0

(d), 130.5 (d), 130.4 (d), 130.0 (d), 129.9 (s), 128.5 (d), 128.4 (d), 128.3 (d), 127.9 (d), 127.8 (d), 124.2 (d), 19.62 (q), 19.58 (q); ir (KBr) 3030, 1665  $\text{cm}^{-1}$ ; ms (12 ev)  $m/z$  405 ( $M^+$ ), 389 ( $M^+-16$ ), 298 ( $M^+-C_6H_5NO$ , base peak); Anal. Calcd for  $C_{28}H_{23}NO_2$ : C, 82.94; H, 5.72; N, 3.45; Found: C, 82.96; H, 5.76; N, 3.32

10. R. A. Abramovitch and E. M. Smith, *J. Heterocycl. Chem.*, 1975, 12, 969

11. 9 (38 %): mp 198 °C;  $^1\text{H}$ -nmr ( $\text{CDCl}_3$ )  $\delta$  8.05-7.85 (m, 4 H), 7.15-7.70 (m, 14 H);  $^{13}\text{C}$ -nmr ( $\text{CDCl}_3$ )  $\delta$  194.2 (s), 149.5 (s), 143.5 (s), 139.6 (d), 139.4 (s), 135.7 (s), 134.1 (s), 133.6 (s), 132.7 (d), 131.3 (d), 130.7 (d), 130.5 (d), 130.4 (d), 129.9 (d), 129.3 (d), 128.8 (d), 128.7 (d), 128.2 (d), 122.2 (d); ir (KBr) 3035, 1660  $\text{cm}^{-1}$ ; HRMS for  $C_{25}H_{18}N_2O_3$  Calcd : 394.1317; Found: 384.1321

Received, 31st October, 1988