

2-(HETEROATOM-SUBSTITUTED)METHYL PENEMS.
IV.¹ OXYGEN DERIVATIVES

Marco Alpegiani, Angelo Bedeschi, Franco Giudici, Ettore Perrone,
Giuseppina Visentin, Franco Zarini, and Giovanni Franceschi*

Farmitalia Carlo Erba, Research & Development, Infectious Diseases Dep.,
Via dei Gracchi 35, Milano, Italy

Abstract — The synthesis of "2-CH₂X" penems wherein X is an oxygen atom part of an acyloxy, N-substituted carbamoyloxy, alkoxy or aryl-oxy residue is described, with emphasis to procedures which exploit a common 2-hydroxymethylpenem precursor. Correlations are attempted between chemical structure of the X moiety and antibacterial activity of obtained compounds.










The subtle influence exerted by the C₂-functionalization of penems on their antibacterial properties has fostered chemists to synthesize hundreds of compounds differing for the sidechain at this position.² Structural analogies with carbapenems and cephalosporins, the carbocyclic and homo counterparts of penems, suggested an attentive investigation in the class of 2-alkylthio and 2-(heteroatom-substituted)methyl derivatives. Recent papers from our laboratories dealt with "2-CH₂X" penems where X is sulphur^{3,4} or nitrogen.¹ This work describes the 2-oxymethylpenems, allocated into three main subclasses, depending on the oxygen being part of an ester (2~25, Scheme I), a carbamate (26~41, Scheme II), or an ether (43~66, Scheme III). For brevity, only the most direct synthetic access will be reported, whenever possible exploiting the pivotal 2-hydroxymethyl intermediate,⁵ variously protected at the C₃-carboxyl and C₈-hydroxyl (1a, a', a'').

2-(Acyloxymethyl)penems. Compound 3c, bearing the acetoxymethyl sidechain featured by the natural cephalosporin C and by some commercial hemisynthetic cephalosporins (e.g., cefotaxime) was the first target in our programme. Its *in vitro* activity,⁶ one order of magnitude higher than that displayed by the reference compound (68, Table III), encouraged the synthesis of analogs (Scheme I). Most of the modifications here described were performed in the attempt to improve stability towards human serum esterases (through bulkier acyl residues⁷: 4c, 5c, 11c), penetration across the outer membrane of Gram-negative bacteria (introduction of charged residues: 13c, 23~25), activity against *Pseudomonas* spp. (basic groups: 23), and intrinsic antibacterial potency (screening of different sidechains, including the aminothiazolyl-2-methoxyiminoacetic⁸ group: 8c). The method of choice was the mild Mitsunobu-Volante procedure,⁹ involving treatment of a suitably protected carbinol (1a, a') and the appropriate carboxylic acid (1~1.2 mol equiv., THF, a few min) with a slight excess of preformed triphenylphosphine - diethyl azodicarboxylate complex (TPP-DEAD). Accordingly, the fully protected penems 2~5a, 7a, 10a, 16a' were isolated in yields ranging from 75% to 95% after flash chromatography. Other methods from 1a included direct acylation with acid chlorides (12a: allyloxyoxalyl chloride, NEt₃, CH₂Cl₂, 69%; 14a: ethyl chlorocarbonate, NEt₃, CH₂Cl₂, 85%) and ketene addition (6a: diketene, cat. amount of NEt₃, CH₂Cl₂, 1 h, 45%).



Scheme I

2-(Acyloxymethyl)penems

a	P= SiMe ₂ t-Bu	P'= allyl
b	P= H	P'= allyl
c	P= H	P'= Na
a'	P= SiMe ₂ t-Bu	P'= p-NB
b'	P= H	P'= p-NB



16 a', b'	W= NHCO ₂ p-NB	n= 0
17 a	W= OSiPh ₂ t-Bu	n= 0
18 a	W= OSiPh ₂ t-Bu	n= 1
19 a	W= OH	n= 0
20 a	W= OH	n= 1
21 a, b	W=	n= 0
22 a, b	W=	n= 1



Functional protecting groups in the acyl residue (10a, 12a, 16a') were selected in order to allow concomitant deblocking, with the exception of *N*-chloroacetyl in 7a, which required an extra deprotection step (thiourea 4 mol equiv., EtOH, 6 h; 85%). Partial oxime isomerization (*Z*:*E* = 3:1) accompanied dechloroacetylation; on prolonging the reaction time, the *E* (*anti*) isomer 9a was exclusively isolated. The quaternary ammonium derivatives 21a, 22a were obtained from the novel carbinol intermediates 19a and 20a (3 mol equiv. of tertiary amine, 1.5 equiv. of triflic anhydride, CH₂Cl₂, -50°C; quenching with 4% aq. HCl and EtOAc extraction), in turn prepared from 1a by Mitsunobu condensation with silylated glycolic¹⁰ and *p*-hydroxymethylbenzoic acid¹¹ (84% and 98%, respectively), followed by selected unmasking of the primary hydroxyl of obtained 17a, 18a (Bu₄NF·3H₂O, HOAc-THF, 1 h, 56% and 90%).

Removal of C₈-hydroxyl and C₃-carboxyl protecting groups was achieved by procedures by now customary in penem chemistry. Thus, prolonged (12–24 h) exposure to the fluoride reagent afforded 2–6b, 8b, 14b, 16b', 21b, 22b (65–85%) from the corresponding *tert*-butyldimethylsilyl ethers. The bis-silylated compound 10a was deprotected to 11b (60%). Compound 12a afforded a mixture of the expected product 12b (34%) and the deacylated penem 1b (25%), owing to competitive fluoride attack at the oxalic ester moiety. Palladium-mediated transallylation (Pd(PPh₃)₄ 0.2 mol equiv., PPh₃, CH₂Cl₂, 15 min) with sodium 2-ethylhexanoate (1.2 mol equiv.) afforded the sodium salts 2–6c, 8c, 11c, 14c and the disodium salt 13c, isolated as white lyophiles (60–92%) after reverse-phase chromatography (LiChroprep C-18, H₂O-MeCN). Substitution of acetic acid for sodium 2-ethylhexanoate in the deallylation of 21b, 22b gave zwitterions 24 and 25. Compound 23 was obtained by reductive debenzoylation of 16b' (Fe/NH₄Cl, aq. THF; 29%).

2-(Carbamoyloxymethyl)penems. The carbamoyloxymethylpenem 28c (code-named FCE 22101),⁵ structurally related to cephamycin C, cefoxitin and cefuroxime, was specifically designed for its predicted stability to human serum esterases.¹² Actually FCE 22101, *in vitro* roughly equivalent to the acetate 3c, proved superior when tested *in vivo*,¹³ even beyond our expectations. In order to investigate the effects of *N*-substitution on antimicrobial activity, several congeners were synthesized (Scheme II).


A first set of products (29–32) arose in connection with a study aimed at an alternative to trichloroacetyl isocyanate for the pilot-plant carbamoylation of 1a. Addition of chlorosulphonyl isocyanate (1 mol equiv., CaCO₃, CH₂Cl₂, -60°C) to 1a gave 27a (not isolable), which could be hydrolyzed *in situ* smoothly, with concomitant loss of SO₃, by quenching the reaction mixture with aqueous dioxane and warming up to room temperature under vigorous stirring (2 h). Conventional deblocking of obtained 28a afforded FCE 22101 in 80% overall yield. By varying the hydrolytic conditions of 27a, the sodium *N*-sulphonate 30a (4% aq. NaHCO₃; 78%), the ethyl *N*-sulphonate 31a (EtOH, 1 h; 72%), and the *N*-sulphonamide 32a (30% aq. NH₄OH; 45%) were isolated. The allophanate 29a'' was obtained by iterative carbamoylation of trichloroethyl bis-protected penem carbinol 1a''. After CSI addition and hydrolysis to 28a'', the second carbamoyl unit was introduced by reaction with trichloroacetyl isocyanate (CH₂Cl₂, 10 min, 0°C) followed by silica gel catalyzed methanolysis (4 h, 25°C; 83%).

Synthesis of the *N*-methyl, *N*-cyclohexyl, and *N*-phenyl carbamates 33–35a, requiring DMAP catalysis for isocyanate addition, has already been reported.¹⁴ It was interesting to ascertain whether the loss of activity against most *Enterobacteriaceae* observed with the last two compounds could be related to their high lipophilicity. Introduction of a charged substituent (pyridiniumomethyl; compound 39) was therefore sought.¹⁵ Curtius degradation of *p*-(*tert*-butyldiphenylsilyloxymethyl)benzoyl azide (refluxing benzene, 4 h) afforded a solution of the corresponding isocyanate, which was directly added to penem carbinol 1a (DMAP 0.1 mol equiv., refluxing

Scheme II

2-(Carbamoyloxymethyl)penems

a	P= SiMe ₂ t-Bu	P'= allyl
b	P= H	P'= allyl
c	P= H	P'= Na
a''	P= COOCH ₂ CCl ₃	P'= CH ₂ CCl ₃



CHCl_3 , 2 h) to produce 36a (75%). Selective desilylation and introduction of the pyridinium moiety (36a→37a→38a; 26%) strictly paralleled the previously described sequence leading to 22a. Tertiary carbamates were obtained by acylating 1a with chlorocarbonyl derivatives of secondary amines. Thus *N*-chlorocarbonylmorpholine reacted with 1a (*N,N*-diethylisopropylamine, CH_2Cl_2) within one night to give 40a (54%), while immediate acylation to 41a (51%) occurred with the more reactive 1-chlorocarbonyl-4-ethyl-2,3-dioxopiperazine.

Removal of protecting groups was carried out as usual. Both trichloroethyl esters in 29a' were cleaved with Zn/HOAc (THF, 4 h), releasing 29c in 25% unoptimized yield after NaHCO_3 treatment and reverse-phase chromatography. Sequential desilylation and deallylation were performed on 30~35a, 40a, 41a to prepare the sodium salts 30~35c, 40c, 41c (60~80% overall). Similarly, zwitterion 39 (66%) was obtained from 38a, save that the Pd-mediated deallylation was carried out in the presence of excess HOAc in place of sodium ethylhexanoate. Unexpected by-products from these reactions included 28b (from 31a), denouncing cleavage of the N-S bond by the fluoride reagent, and FCE 22101 (27c), which formed in increasing amounts while recording the nmr spectrum of disodium salt 30c ($\text{DMSO}-d_6$, 45 °C). Following this observation, FCE 22101 allyl ester (28b) was quantitatively obtained from the sodium *N*-sulphonate 30b by thermolysis in $\text{DMSO}-\text{MeCN}$ (1:9, 80 °C, 6 h).

2-(Alkoxyethyl)- and 2-(Aryloxyethyl)penems. Very recently, 3-alkoxyethyl-cephalosporins have received attention for their interesting pharmacokinetic properties, in particular improved oral bioavailability.¹⁶ A brief investigation in the corresponding class of penems was therefore undertaken (Scheme III). Instability of 2-halomethylpenems to solvolytic conditions advised for a different approach, involving the versatile silver azetidinyl mercaptide 42b.¹⁷ Acylation of the latter with the appropriate alkoxyacetyl chloride (CH_2Cl_2 , -10 °C, 20 min) followed by intramolecular Wittig reaction (refluxing toluene, 2 h) proceeded smoothly, affording penems 43~48b in 50~70% overall yields. The 2-hydroxymethylpenem intermediate 1a could displace in acceptable yields (45~55%) the most electrophilic halides (allyl and benzyl bromides) in the presence of silver triflate (2,6-lutidine, CH_2Cl_2 , 20 min), providing an alternative access to penem ethers 46a and 47a.


The presence of zinc at the active site of enzymes implicated in the degradation of β -lactam antibiotics (mammalian dehydropeptidases and class B bacterial β -lactamases) conferred potential interest to polyethers, such as MEM-ether 50a, THP-ether 51a, and orthoformate 52a. These products were uneventfully obtained from carbinol 1a under customary conditions (50a: MEM-chloride, Hünig base, CH_2Cl_2 , overnight, 76%; 51a: 3,4-dihydro-2*H*-pyran, PTSA catalysis, CH_2Cl_2 , 1 h quantitative; 52a: neat trimethyl orthoformate, 60 °C, 7 h, 70%).

Aryloxyethylpenems present a substitution pattern unusual for hemisynthetic cephalosporins. For the synthesis of a few representatives (53c, 57c, 58c, 60c, 65, 66) the Mitsunobu-Volante procedure was again selected as the basic strategy. Not unexpectedly, yields of condensation impressively varied according to the pK_A of the reagent. Thus, 1a reacted with unsubstituted phenol (THF, 1 h) to give a very modest yield of 53a, *m*-nitrophenol gave a mixture of penem 55a and 2-exomethylene-penam 67a¹⁸ (30:70; 36%), while the more acidic *o*- and *p*-isomers reacted within a few seconds at 0 °C to give the anticipated products 54a (83%) and 56a (74%). Consistently, access to the *p*-hydroxymethyl derivative 62a by this procedure was inconvenient, and in fact *p*-(*tert*-butyldiphenylsilyloxyethyl)phenol condensed with 1a in low and capricious yields (up to 23%). Condensations targetted at the quarternary ammonium compounds 65, 66, and at carbamate 60c, were best run with *p*-hydroxybenzaldehyde, here selected as an electron-poor equivalent of *p*-(hydroxymethyl)phenol. Use of this reagent led to the formyl derivative 59a in yields

Scheme III

2-(Alkoxymethyl and Aryloxymethyl)penems

a	P= SiMe ₂ t-Bu	P'= allyl
b	P= H	P'= allyl
c	P= H	P'= Na



exceeding 80%; subsequent reduction, either by excess sodium cyanoborohydride (HOAc, THF, r.t., 2 h, 88%) or stoichiometric K-selectride (THF, -60°C, 30 min, 91%), afforded **62a** in a reproducible sequence, setting the stage for carbamoylation to **60b** (Cl_3CCONCO , CH_2Cl_2 , -20°C, then MeOH/SiO_2 , 4 h; 55%) or for quaternarization to **63a**, **64a** (isolated crude; quantitative) under conditions identical to those described for the synthesis of **21a**, **22a**.

Desilylation of **46a**, **47a**, **50a**, **51a**, **53a**, **54a**, **56a**, **60a**, **63a**, **64a** ($\text{Bu}_4\text{NF} \cdot 3\text{H}_2\text{O}$, HOAc, THF, overnight) proceeded uneventfully, save that the mixed orthoformate **52a** released the hydroxymethylpenem **1b** instead of **52b** (quantitative conversion). The 30:70 mixture of *m*-nitrophenoxy endo/exo isomers afforded an unvaried proportion of desilylated compounds **55b**, **67b** (not processed further). The *o*- and *p*-nitro compounds **54b**, **56b** were subjected to dissolving metal reduction (Zn/HOAc , CH_2Cl_2 , 30 min) to obtain the aminophenoxy derivatives **57b** (68%) and **58b** (48%). Finally, Pd-catalyzed transallylation with sodium 2-ethylhexanoate or excess acetic acid gave penem monosodium salts **43~47c**, **50c**, **51c**, **53c**, **57c**, **58c**, **60c** (40~85%), the disodium salt **49c** (82%), and zwitterions **65**, **66** (30% overall from **62a**) required for microbiological evaluation.

Table I - Spectral data of key intermediates

Compd	ir (ν_{max} , cm^{-1}) ^(a)	¹ H nmr (δ , ppm) ^(b)
8a	1775, 1735, 1705	3.74(1H, dd, J=1.8 and 4.2Hz), 4.04(3H, s), 5.54(2H, ABq, J=15.4Hz), 5.74(1H, d, J=1.8Hz), 6.80(1H, s)
9a	1785, 1740-1695	3.72(1H, dd, J=1.8 and 4.2Hz), 4.12(3H, s), 5.48(2H, ABq, J=15.1Hz), 5.61(1H, d, J=1.8Hz), 7.48(1H, s)
15a	1785, 1750, 1710	3.71(1H, dd, J=1.9 and 4.5Hz), 3.82(3H, s), 5.10 and 5.56 (2H, two d, J=15.0Hz), 5.60(1H, d, J=1.9Hz) [200MHz]
19a	1790, 1750, 1705	2.7(1H, br, exch. D_2O), 3.71(1H, dd, J=1.6 and 4.4Hz), 4.20 (2H, s), 5.18 and 5.59(2H, two d, J=15.0Hz), 5.60(1H, d, 1.6Hz)
20a	1785, 1715	2.34(1H, br, exch. D_2O), 3.65(1H, dd, J=2 and 4.4 Hz), 4.70 (2H, s), 5.45(2H, ABq, J=15.5Hz), 5.57(1H, d, J=2Hz)
28a''	3480, 3320, 1770, 1730, 1700 (KBr)	3.93(1H, dd, J=2 and 8Hz), 4.73(2H, s), 4.81(2H, s), 5.26 (2H, ABq, J=15.5Hz), 5.62(1H, d, J=2Hz)
29a''	3500, 3405, 1805 1755, 1725	4.02(1H, dd, J<2 and 7Hz), 4.77(2H, s), 4.87(2H, s), 5.38 (2H, ABq, J=15Hz), 5.70(1H, ABq, J<2Hz)
52a	1775, 1700 (KBr)	3.35(6H, s), 3.67(1H, dd, J=1.6 and 4.8Hz), 4.67 and 4.89 (2H, two d, J=15.5Hz), 5.08(1H, s), 5.56(1H, d, J=1.6Hz)
54b	1790.1690 (KBr)	2.50(1H, br, exch. D_2O), 3.78(1H, dd, J=1.8 and 6.0Hz), 5.40(2H, ABq, J=16Hz), 5.65(1H, d, J=1.8Hz), 6.9-8.0(4H, m)
55b	1785, 1705	3.75(1H, dd, J=2 and 4.3Hz), 5.38(2H, ABq, J=15Hz), 5.63 (1H, d, J=2Hz), 7.3-7.9(4H, m)
56b	1790, 1695 (KBr)	3.76(1H, dd, J=1.8 and 6.1Hz), 5.40(2H, ABq, J=15.5Hz), 5.63(1H, d, J=1.8Hz), 6.95(2H, d, J=8.5Hz), 8.18(2H, d, J=8.5Hz)
59a	1790, 1695	3.72(1H, dd, J=1.6 and 4.4Hz), 5.20 and 5.51(2H, two d, J=15.1Hz), 5.61(1H, d, J=1.6Hz), 7.03(2H, d, J=8.8Hz), 7.84(2H, d, J=8.8Hz), 9.90(1H, s) [200MHz]
62a	1790, 1715	2.4(1H, br, exch. D_2O), 3.65(1H, dd, J=1.9 and 4.1Hz), 4.82 (2H, s), 5.47(2H, ABq, J=15.6Hz), 5.67(1H, d, J=1.9Hz), 7.49(2H, d, J=8Hz), 8.08(2H, d, J=8Hz)

(a) In CHCl_3 unless otherwise stated.

(b) In CDCl_3 at 60 MHz unless otherwise stated (salient data).

Table II - Spectral data of 2-(oxygen-substituted)methylpenem-3-carboxylic acids (sodium or internal salts)

Compd	ir (KBr) $\nu_{\max}$ (cm) ⁻¹	uv (H ₂ O) $\lambda_{\max}$ (nm)	¹ H nmr (D ₂ O) ¹ δ (ppm)
2c	1765, 1720, 1610, 1585	258(ϵ =5358), 308(ϵ =7587)	1.30(3H, d, J=6.7Hz), 3.95(1H, dd, J=1.4 and 6.0Hz), 4.26(1H, dq, J=6.0 and 6.7Hz), 5.18 and 5.56(2H, two d, J=13.9Hz), 5.69(1H, d, J=1.4Hz), 8.19(1H, s)
3c	1770, 1750, 1610	258(ϵ =4630), 308(ϵ =6870)	1.31(3H, d, J=6.5Hz), 2.14(3H, s), 3.93(1H, dd, J=1.4 and 5.8Hz), 4.26(1H, m), 5.10 and 5.46(2H, two d, J=14.4Hz), 5.68(1H, d, J=1.4Hz)
4c	1760, 1740, 1610, 1590	214(ϵ =2688), 259(ϵ =2197), 307(ϵ =3708)	1.19(3H, t, J=7.5Hz), 1.56(3H, d, J=6.3Hz), 1.89(2H, m), 2.66(2H, t, J=7.1Hz), 4.16(1H, dd, J=1.6 and 6.0Hz), 4.50(1H, m), 5.38 and 5.72(2H, two d, J=14.2Hz), 5.91(1H, d, J=1.6Hz) [taken at 45°C]
5c	1760, 1605, 1580	306(ϵ =6267)	1.29(3H, d, J=6.5Hz), 3.90(1H, br. d, J=5.9Hz), 4.23(1H, dq, J=5.9 and 6.5Hz), 5.62(1H, br. s), 5.32 and 5.66(2H, two d, J=13.9Hz), 7.53(2H, m), 7.68(1H, m), 8.02(2H, d, J=8.2Hz)
6c		260(ϵ =2630), 305(ϵ =4879)	1.29(3H, d, J=6.5Hz), 2.32(3H, s), 3.92(1H, dd, J=1.5 and 5.7Hz), 4.24(1H, m), 5.15 and 5.53(2H, each d, J=14.0Hz), 5.65(1H, d, J=1.5Hz)
8c	1760, 1740, 1605, 1580	304	1.32(3H, d, J=6.3Hz), 3.91(1H, dd, J=1.5 and 6.0Hz), 4.03(3H, s), 4.27(1H, m), 5.60(2H, ABq, J=14.0Hz), 5.66(1H, d, J=1.5Hz), 6.99(1H, s) [60MHz]
11c	1755(br), 1595	258(ϵ =3370),	1.30(3H, d, J=6.4Hz), 3.17(1H, dd, J=2.5 and 15.2Hz), 3.41(1H, dd, J=1.5 and 15.2Hz), 3.93(1H, dd, J=1.5 and 5.9Hz), 4.25(1H, dq, J=5.9 and 6.4Hz), 4.41(1H, dd, J=2.4 and 5.6Hz), 5.19 and 5.60(2H, two d, J=14.0Hz), 5.67(1H, d, J=1.5Hz)
13c	1750(br), 1660, 1610	258(ϵ =3121), 308(ϵ =4824)	
14c	1750(br), 1610-1580	255(ϵ =3058), 307(ϵ =3760)	1.35(3H, t, J=7.0Hz), 1.36(3H, d, J=6.4Hz), 3.95(1H, dd, J=1.7 and 5.8Hz), 4.27(1H, m), 4.29(2H, q, J=7.0Hz), 5.37(2H, ABq, J=14.0Hz), 5.68(1H, d, J=1.7Hz) [60MHz]
23	1750, 1620-1580	259(ϵ =2204), 308(ϵ =3020)	1.30(3H, d, J=6.5Hz), 3.94(1H, dd, J=1.6 and 5.8Hz), 3.98(2H, s), 4.25(1H, m), 5.19 and 5.61(2H, two d, J=13.8Hz), 5.69(1H, d, J=1.6Hz)
24	1775-1745, 1605, 1585	258(ϵ =3258), 308(ϵ =4919)	1.29(3H, d, J=6.4Hz), 2.24(4H, m), 3.27(3H, s), 3.6-3.9(4H, m), 3.93(1H, dd, J=1.4 and 5.9Hz), 4.24(1H, dq, J=5.9 and 6.4Hz), 4.45(2H, s), 5.22 and 5.63(2H, two d, J=14.0Hz), 5.68(1H, d, J=1.4Hz)
25	1765, 1720, 1605, 1575	236(ϵ =18923), 307(ϵ =5600)	1.27(3H, d, J=6.3Hz), 3.79(1H, dd, J=1.6 and 5.8Hz), 4.20(1H, m), 5.12 and 5.42(2H, two d, J=14.8Hz), 5.51(1H, d, J=1.6Hz), 5.92(2H, s), 7.50 and 7.94(4H, two d, J=8.3Hz), 8.15(2H, m), 8.63(1H, m), 8.99(2H, m)

Table II - Continued

28c	1755, 1730, 1600-1570	258(ϵ =4150), 306(ϵ =6030)	1.31(3H,d,J=6.5Hz), 3.91(1H,dd,J=1.5 and 6.0Hz), 4.25(1H,dq,J=6.0 and 6.5Hz), 5.02 and 5.36(2H,two d,J=14.5Hz), 5.66(1H,J=1.5Hz)
29c	1770,1710, 1660,1600	257(ϵ =3548), 306(ϵ =4246)	1.29(3H,d,J=6.5Hz), 3.93(1H,dd,J=1.6 and 6.0Hz), 4.24(1H,m), 5.14 and 5.55(2H,two d,J=14.2Hz), 5.67(1H,d,J=1.6Hz)
30c	1775(br), 1605	255(ϵ =2349), 307(ϵ =3130)	1.30(3H,d,J=6.4Hz), 3.96(1H,dd,J=1.4 and 5.9Hz), 4.25(1H,m), 5.12 and 5.52(2H,two d,J=14.5Hz), 5.67(1H,d,J=1.4Hz)
31c	1760, 1650-1580	258(ϵ =4434), 307(ϵ =6632)	1.29(3H,d,J=6.3Hz), 1.33(3H,t,J=7.0Hz), 3.91(1H,dd,J=1.6 and 6.0Hz), 4.18(2H,q,J=7.0Hz), 4.24(1H,m), 5.03 and 5.38(2H,two d,J=14.9Hz), 5.64(1H,d,J=1.6Hz)
32c	1750, 1620(br)	256(ϵ =2326), 304(ϵ =2804)	1.31(3H,d,J=6.4Hz), 3.93(1H,dd,J=1.6 and 5.8Hz), 4.25(1H,m), 5.03 and 5.37(2H,two d,J=14.9Hz), 5.66(1H,d,J=1.6Hz)
33c	1770,1720, 1590 [Nujol]	304(ϵ =5670) [EtOH]	1.30(3H,d,J=6.5Hz), 2.73(3H,s), 3.90(1H,dd,J=1.6 and 6.0Hz), 4.24(1H,dq,J=6.0 and 6.5 Hz), 5.07 and 5.36(2H,two d,J=14.6Hz), 5.65(1H,d,J=1.6Hz)
34c	1765,1705, 1590 [Nujol]	306(ϵ =7600) [EtOH]	1.31(3H,d,J=6.4Hz), 1.2-1.9(10H,m), 3.3-3.4(1H,m), 3.90(1H,dd,J=1.3 and 6.2Hz), 4.24(1H,dq,J=6.2 and 6.4Hz), 5.09 and 5.42(2H,two d,J=14.8Hz), 5.65(1H,d,J=1.3Hz)
35c	1760-1710, 1595	306(ϵ =5229) [EtOH]	1.29(3H,d,J=6.5Hz), 3.90(1H,dd,J=1.4 and 6.0Hz), 4.23(1H,dq,J=6.0 and 6.5Hz), 5.14 and 5.51(2H,two d,J=13.6Hz), 5.64(1H,d,J=1.4Hz), 7.2-7.4(5H,m)
39	1760,1720, 1590	307	1.24(3H,d,J=6.4Hz), 3.82(1H,dd,J=1.6 and 5.8Hz), 4.18(1H,dq,J=5.8 and 6.4Hz), 5.05 and 5.42(2H,two d,J=14.6Hz), 5.55(1H,d,J=1.6Hz), 5.75(2H,m), 7.43(4H,m), 8.05(2H,m), 8.53(1H,m), 8.89(2H,m)
40c	1755,1720, 1590	305(ϵ =5795)	1.31(3H,d,J=6.4Hz), 3.53(4H,m), 3.75(4H,m), 3.93(1H,dd,J=1.4 and 6.0Hz), 4.26(1H,dq,J=6.0 and 6.4Hz), 5.13 and 5.49(2H,two d,J=14.6Hz), 5.67(1H,d,J=1.4Hz)
41c	1755,1720, 1675,1600	224,306	1.20(3H,t,J=7.3Hz), 1.30(3H,d,J=6.4Hz), 3.51(2H,q,J=7.3Hz), 3.73(2H,m), 3.95(1H,dd,J=1.7 and 5.9Hz), 4.12(2H,m), 4.25(1H,m), 5.65(1H,d,J=1.7Hz), 5.26 and 5.66(2H,two d,J=14.2Hz)
43c	1755,1650, 1600,1580	258(ϵ =4044), 306(ϵ =6076)	1.30(3H,d,J=6.3Hz), 3.38(3H,s), 3.91(1H,dd,J=1.7 and 6.1Hz), 4.25(1H,dq,J=6.1 and 6.3 Hz), 4.50 and 4.80(2H,two d,J=14.0Hz), 5.66(1H,d,J=1.7Hz)
44c	1760,1605, 1580	306(ϵ =5613)	1.19(3H,t,J=7.1Hz), 1.30(3H,d,J=6.4Hz), 3.5-3.7(2H,m), 3.90(1H,dd,J=1.6 and 6.0Hz), 4.48(1H,dq,J=6.0 and 6.4Hz), 4.52 and 4.85(2H,two d,J=14.2Hz), 5.65(1H,d,J=1.6Hz)

Table II - Continued

45c	1760, 1600-1560	258(ϵ =4100), 306(ϵ =6050)	0.92(3H,t,J=7.5Hz), 1.32(3H,d,J=6.5Hz), 1.61(2H,m), 3.5-3.6(2H,m), 3.91(1H,dd,J=1.6 and 6.0Hz), 4.26(1H,dq,J=6.0 and 6.5Hz), 4.55 and 4.68(2H,two d, J=14.9Hz), 5.66(1H,d, J=1.6Hz)
46c	1760,1585	258(ϵ =3764), 307(ϵ =5637)	1.31(3H,d,J=6.5Hz), 3.92(1H,dd,J=1.6 and 5.9Hz), 4.0-4.2(2H,m), 4.26(1H,dq,J=5.9 and 6.5Hz), 4.55 and 4.90(2H,two d,J=14.1Hz), 5.2-5.4(2H,m), 5.67(1H,d,J=1.6Hz), 5.9-6.1(1H,m)
47c	1760, 1590-1570	258(ϵ =3870), 307(ϵ =5474)	1.34(3H,d,J=6.5Hz), 3.88(1H,dd,J=1.4 and 4.7Hz), 4.24(1H,dq,J=4.7 and 6.5Hz), 4.61(2H,m), 4.62 and 4.93(2H,two d,J=14.4Hz), 5.62(1H,d,J=1.4Hz), 7.44(5H,m)
49c	1765, 1590(br)	258(ϵ =3187), 307(ϵ =4864)	1.32(3H,d,J=6.5Hz), 3.94(1H,dd,J=1.6 and 6.0Hz), 3.93 and 4.01(2H,two d,J=15.5Hz), 4.26(1H,dq,J=6.0 and 6.5Hz), 4.59 and 4.89(2H,two d,J=14.4Hz), 5.67(1H,d,J=1.6Hz)
50c	1760, 1600-1580	259(ϵ =3350), 307(ϵ =5412)	1.21(3H,d,J=6.4Hz), 3.32(2H,s), 3.54(2H,m), 3.68(3H,m), 4.22(1H,m), 4.71(2H,m), 4.80 and 4.93(2H,two d,J=13.8Hz), 5.66(1H, br s)
51c	1760,1580	260(ϵ =2678) 306(ϵ =5025)	1.32(3H,d,J=6.4Hz), 1.4-1.9(6H,m), 3.60(1H,m), 3.90(2H,m), 4.25(1H,m), 4.58, 4.73, 4.94 and 5.05(2H,each d,J=14Hz), 4.78(1H,m), 5.67(1H,d,J=1.4Hz) [taken at 45°C]
53c	1760,1590	262(ϵ =4484) 308(ϵ =5513)	1.31(3H,d,J=6.5Hz), 3.89(1H,dd,J=1.6 and 6.0Hz), 4.25(1H,dq,J=6.0 and 6.5Hz), 5.20 and 5.53(2H,two d,J=14.6Hz), 5.62(1H,d,J=1.6Hz), 7.06-7.46(5H,m)
57c	1765,1600	306(ϵ =5273)	1.27(3H,d,J=6.4Hz), 3.85(1H,dd,J=1.2 and 6.0Hz), 4.21(1H,m), 5.15 and 5.8(2H,two d, J=14.5Hz), 5.57(1H,d,J=1.2Hz), 6.75-7.3(4H,m)
58c	1765, 1600-1580	305(ϵ =5466)	1.27(3H,d,J=6.4Hz), 3.86(1H,dd,J=1.6 and 6.0Hz), 4.21(1H,m), 5.07 and 5.41(2H,two d, J=14.5Hz), 5.58(1H,d,J=1.6Hz), 6.86(4H,m)
60c	1760,1705, 1610,1585	308(ϵ =5387)	1.26(3H,d,J=6.3Hz), 3.79(1H,dd,J=1.6 and 5.8Hz), 4.19(1H,dq,J=5.8 and 6.3Hz), 4.98(2H,s), 5.12 and 5.44(2H,two d,J=14.6Hz), 5.52(1H,d,J=1.6Hz), 6.97(2H,d,J=8.5Hz), 7.31(2H,d,J=8.5Hz)
65	1765,1600	228,260sh, 308(ϵ =4780)	1.27(3H,d,J=6.3Hz), 2.22(4H,m), 2.93(3H,s), 3.30-3.65(4H,m), 3.84(1H,d,J=6.0Hz), 4.23(1H,m), 4.45(2H,s), 5.19 and 5.51(2H,two d, J=14Hz), 5.58(1H,s), 6.95-7.55(4H,m)
66	1760,1605, 1580	258(ϵ =8935) 306(ϵ =5647)	1.22(3H,d,J=6.3Hz), 3.72(1H,dd,J=1.5 and 6.0Hz), 4.14(1H,dq,J=6.0 and 6.3Hz), 5.02 and 5.34(2H,two d,J=14.4Hz), 5.43(1H,d, J=1.5Hz), 5.70(2H,s), 6.95 and 7.39(4H,two d,J=8.6Hz), 8.01(2H,dd,J=5.6 and 7.0Hz), 8.50(1H,d,J=7.0Hz), 8.87(2H,d,J=5.6Hz)

(a) At 200 MHz unless otherwise stated.

Table III - *In vitro* antibacterial activity of penems












Compd		MIC(^a)					
		<i>S.a.</i>	<i>S.f.</i>	<i>E.c.</i>	<i>E.c.</i> (+)	<i>S.t.</i>	<i>C.f.</i> (+)
Acyloxymethyl compounds							
2c	OCHO	0.011	6.25	0.78	0.78	0.78	nd(^b)
3c	OCOMe	0.022	3.12	0.39	0.39	0.39	nd
4c	OCOPr- <i>n</i>	0.045	6.25	0.39	0.78	12.5	>50
5c		0.022	3.12	0.19	0.78	50	>50
6c	OCOCH ₂ COMe	<0.09	nd	0.78	1.56	nd	nd
8c		0.09	12.5	nd	nd	>25	>25
11c		0.19	3.12	0.78	0.78	3.12	6.25
13c	OCOCO ₂ Na	0.09	12.5	1.56	6.25	0.78	nd
14c	OCOEt	0.045	12.5	nd	nd	25	>25
23	OCOCH ₂ NH ₂	nd	12.5	nd	nd	3.12	nd
24		0.045	12.5	3.12	3.12	3.12	nd
25		0.011	0.78	0.39	1.56	0.39	6.25
Carbamoyloxymethyl compounds							
28c	CONH ₂	0.045	3.12	0.78	0.78	0.78	1.56
29c	CONHCONH ₂	0.011	6.25	0.78	3.12	1.56	nd
30c	CONHSO ₃ Na	0.19	25	1.56	25	0.78	nd
31c	CONHSO ₃ Et	0.78	50	6.25	25	3.12	>25
32c	CONHSO ₂ NH ₂	0.39	100	6.25	25	6.25	>25
33c	CONHMe	0.22	3.12	0.78	0.78	0.78	nd
34c		0.045	50	0.78	0.78	100	>25
35c		0.022	1.56	0.19	0.39	6.25	>50
39		0.045	0.39	0.78	1.56	0.78	3.12
40c		0.09	6.25	0.78	6.25	25	>25
41c		0.39	12.5	0.78	25	0.78	nd

Table III - Continued

Compd		MIC ^(a)					
		S.a.	S.f.	E.c.	E.c.(+)	S.t.	C.f.(+)
Hydroxy-, alkoxy-, aryloxymethyl compounds							
1c	OH	0.19	nd	3.12	6.25	3.12	≥25
43c	OMe	0.045	3.12	0.39	0.78	0.39	6.25
44c	OEt	0.09	12.5	0.78	0.78	1.56	25
45c	OPr-n	0.09	12.5	0.78	0.78	6.25	>50
46c	OCH ₂ CH=CH ₂	0.045	12.5	0.39	0.39	1.56	>50
47c		0.045	12.5	0.78	0.78	>50	>50
49c	OCH ₂ COONa	1.56	>50	1.56	1.56	1.56	6.25
50c	OCH ₂ OCH ₂ CH ₂ OMe	0.09	12.5	0.39	0.39	0.78	>50
51c		0.09	12.5	0.78	0.78	12.5	>50
53c		0.011	1.56	0.78	1.56	100	>25
57c		0.022	1.56	1.56	3.12	25	>25
58c		0.022	nd	1.56	3.12	25	>25
60c		0.011	1.56	0.39	0.78	25	>25
65		0.022	3.12	0.78	1.56	1.56	nd
66		0.011	1.56	0.78	0.78	0.78	1.56
Reference							
68		0.38	12.5	3.12	3.12	1.56	>25

(^a) MICs (mcg/ml) were determined by the standard two-fold agar dilution method in Bacto Antibiotic Medium 1 (Difco); inoculum size 10⁴ cfu. Organisms included in this table are: S.a., *Staphylococcus aureus* Smith; S.f., *Streptococcus faecium* ATCC 8043; E.c., *Escherichia coli* B; E.c.(+), *E. coli* B β-lactamase producer; S.t., *Salmonella typhi* ATCC 14028; C.f.(+), *Citrobacter freundii* ATCC 4051 cephalosporin-resistant. (^b) nd = Not determined.

Table III shows the *in vitro* activity of the title compounds against six representative bacterial strains, in comparison with the unsubstituted 2-methylpenem reference 68. In particular, *Staphylococcus aureus* Smith and *Salmonella typhi* ATCC 14028 were selected as common Gram-positive and Gram-negative organism, respectively; data on *Escherichia coli* B, characterized by a permeable outer membrane, reflect more closely the intrinsic activity of each compound. With the exception of *Pseudomonas aeruginosa*, which proved resistant to all of the tested compounds, and of some "difficult" opportunistic pathogens, here exemplified by *Citrobacter freundii* ATCC 4051, the vast majority of compounds showed good levels of activity, usually superior to that of the unsubstituted 2-methyl reference. It is hard to identify general features proper of this family of "2-CH₂X" penems, either in comparison with others^{1,3,4} characterized by a different X hetero atom, or according to a division into chemical subclasses. Within each of these (esters, carbamates, ethers), simple substituents proved often the best for imparting a wide spectrum of activity (3c, 28c, 43c). Homologation of the carbon chain of the acyl (4c, 5c), *N*-substituted carbamate (34c, 35c), or ether residue (44c, 45c, 47c) almost immediately led to a decrease in activity against Gram-negative rods, as a result of impaired outer membrane penetration. Introduction of negatively ionizable groups (13c, 30-32c, 49c) restored good penetration properties but depressed the intrinsic activity (compare MIC values on *E. coli* and *S. typhi*).

On the contrary, introduction of a quaternary ammonium overcame the permeability problem without affecting intrinsic activity on both Gram-positive and -negative bacteria (25, 39, 65, 66). The 2-aryloxymethyl compounds, reported by Ciba scientist as devoid of activity against enterobacteria,¹⁷ are particularly interesting in this respect. Compound 66 (FCE 24386) was selected for its good *in vitro* activity and unusually extended plasma half-life;¹⁵ unfortunately, it also displayed unacceptable levels of CNS toxicity (anoxic convulsions in mice), which prevented further development.

Other sidechain modifications met with little success. Introduction of the amino-thiazolyl moiety (8c), or basic groups (23, 57c, 58c), or potential metal ligands (49c, 50c, 51c) did not result in improved antibacterial potency and anti-pseudomonal activity, nor in increased stability towards renal degrading enzymes (dehydropeptidases, data not shown). Further works confirmed 28c (FCE 22101) and 43c (FCE 24964)¹⁹ as the most interesting compounds in this penem family. An oral pro-drug formulation of the latter, FCE 25199,²⁰ now undergoing toxicological studies in animals, in a short time will hopefully join the exiguous maniple of clinically evaluated penems.

ACKNOWLEDGEMENTS

Acknowledgements are due to Costantino Della Bruna, Giuseppe Meinardi, Daniela Jabes for antimicrobial evaluation, and to Daniela Borghi for nmr spectroscopy.

REFERENCES AND NOTES

1. Part III of this series: M. Alpegiani, A. Bedeschi, E. Perrone, F. Zarini, and G. Franceschi, *Heterocycles*, 1988, 27, 1329.
2. S.W. McCombie and A.K. Ganguly, *Medicinal Research Reviews*, 1988, 8, 393
3. M. Alpegiani, A. Bedeschi, E. Perrone, F. Zarini, and G. Franceschi, *Heterocycles*, 1985, 23, 2255.
4. M. Alpegiani, E. Perrone, and G. Franceschi, *Heterocycles*, 1988, 27, 49.
5. G. Franceschi, M. Foglio, M. Alpegiani, C. Battistini, A. Bedeschi, E. Perrone, F. Zarini, F. Arcamone, C. Della Bruna, and A. Sanfilippo, *J. Antibiotics*, 1983, 36, 938.

6. A. Sanfilippo, C. Della Bruna, D. Jabes, E. Morvillo, and G. Schioppacassi, *J. Antibiotics*, 1982, 35, 1248.
7. 3-Acetoxyethylcephalosporins are known to be metabolized by hydrolysis to the corresponding, less active 3-hydroxymethyl compounds. It has been shown that the more bulky is the 3-acyloxymethyl group, the more resistant is the cephalosporin molecule to enzymatic hydrolysis: T. Takaya, H. Takasugi, T. Murakawa, and H. Nakano, *J. Antibiotics*, 1981, 34, 1300. See also note 12.
8. The intrinsic potency of β -lactam antibiotics depends on their molecular recognition by the bacterial targets, the cell-wall synthesizing enzymes. The three-dimensional requirements for such recognition are not known. The 2-aminothiazol-4-ylacetamido sidechain with a syn-methoxyimino group at C₂ (ATMO) is the most potent activator of cepheids: R. Bucourt, R. Heymes, A. Lutz, L. Pénasse, and J. Perronnet, *Tetrahedron*, 1978, 34, 2233.
9. R.P. Volante, *Tetrahedron Lett.*, 1981, 22, 3199.
10. E. Fontana, M. Alpegiani, E. Perrone, and G.P. Vicario, *J. Label. Compounds Radiopharm.*, 1987, 24, 41.
11. F.H. Case, *J. Am. Chem. Soc.*, 1925, 47, 3303.
12. Carbamates are not easily hydrolyzed by esterases or other enzymes present in human serum. An indirect evidence is provided by the following result. The chemical half-lives of 2-acetoxyethylpenem 3c and 2-carbamoyloxymethylpenem 28c were determined by HPLC in 4:1 human serum /10 mM pH 7.4 phosphate buffer and in phosphate buffer alone (37°C, 0.33 mM initial concentration). While under the latter conditions the two compound showed almost identical stability, in the presence of serum depletion of 3c was much faster ($t_{1/2}$ = 2.6 h) than that of 28c ($t_{1/2}$ = 8.2 h). We thank Riccardo Corigli for this experiment.
13. G. Franceschi, E. Perrone, M. Alpegiani, A. Bedeschi, C. Battistini, F. Zarini, and C. Della Bruna, *J. Antimicrob. Chemother.*, 1989, 23 (Suppl. C), 1.
14. C. Battistini, S. Vioglio, C. Scarafile, and G. Franceschi, *Heterocycles*, 1985, 23, 1929.
15. E. Perrone, M. Alpegiani, A. Bedeschi, F. Giudici, F. Zarini, G. Franceschi, C. Della Bruna, D. Jabes, and G. Meinardi, *J. Antibiotics*, 1987, 40, 1636.
16. K. Fujimoto, S. Ishihara, H. Yanagisawa, J. Ide, E. Nakayama, H. Nakao, S.-I. Sugawara, and M. Iwata, *J. Antibiotics*, 1987, 40, 370.
17. H.G. Capraro, M. Lang, E. Hungerbühler, P. Schneider, W. Tosch, and O. Zak, Program and Abstracts of the 28th Intersci. Conf. on Antimicrob. Agents Chemother., Los Angeles, 1988, No. 224, p. 149.
18. Compounds 67a and 67b were obtained as single isomers. Relevant analytical data for the latter are as follows: ir, $\nu_{\max}$ (CHCl₃) 1780, 1740 cm⁻¹; nmr, δ (CDCl₃) 3.44(1H, dd, J=1.8 and 4.5Hz), 3.50(1H, br s, exch. D₂O), 5.41(1H, d, J=1.8Hz), 5.47(1H, d, J=1.2Hz), 6.94(1H, d, J=1.2Hz), 7.25-8.10(4H, m). The high deshielding experienced by the C₃ proton (δ = 5.47) is characteristic of the 3*S*-stereochemistry, as discussed in reference 3. Assignment of the *Z*-alkene geometry is based on analogy considerations: 2-heterocyclylthiomethylpenems isomerize exclusively or prevalently ($\geq$ 9:1) to *Z*-configured 2-exomethylene-penams. See again reference 3.
19. C. Della Bruna, D. Jabes, P. Castellani, R. Rossi, G. Meinardi, M. Alpegiani, F. Zarini, E. Perrone, and G. Franceschi, Program and Abstracts of the 28th Intersci. Conf. on Antimicrob. Agents Chemother., Los Angeles, 1988, No 225, p. 149.
20. C. Della Bruna, P. Castellani, D. Jabes, G. Meinardi, R. Rossi, G. Franceschi, E. Perrone, and R. Roncucci, Program and Abstracts of the 29th Intersci. Conf. on Antimicrob. Agents Chemother., Houston, 1989, No 87, p. 115.

Received, 16th October, 1989