

HIGH PRESSURE DIELS-ALDER REACTION OF 1-METHYL-2(1*H*)-PYRIDONES HAVING A PHENYL GROUP WITH *N*-PHENYLMALEIMIDE

Hiroshi Tomisawa*, Hiroto Nakano, and Hiroshi Hongo

Tohoku College of Pharmacy

4-4-1 Komatsushima, Aoba-ku, Sendai 981, Japan

Abstract — Diels-Alder reaction of 1-methyl-2(1*H*)-pyridones (**Ia-d**) having a phenyl group at 3,4,5, and 6 position with *N*-phenylmaleimide (**II**) under 10 kbar at 110 °C for 72 h gave a mixture of the *endo* and *exo* adducts (**IIIa-d** and **IVa-d**) in good yields, some of which were unobtainable under atmospheric pressure conditions.


Isoquinuclidine derivatives prepared by Diels-Alder reaction of 1-substituted 2(1*H*)-pyridones with dienophiles are interesting as possible intermediates for iboga alkaloids.¹ We have developed a synthetic route toward this heterocyclic ring system having various substituents.² We now wish to report the reaction of 1-methyl-2(1*H*)-pyridones(**Ia-d**)³ having a phenyl group at 3,4,5, and 6 position with *N*-phenylmaleimide(**II**) under atmospheric and high pressure conditions. Isoquinuclidine derivatives having a phenyl group linked to quaternary carbon derived from **Ia,d** and **II** are expected to possess the interesting pharmacological activities.⁴ Although the high pressure strategy has proven extremely useful to surmount the energy barrier imposed by steric and electronic effects in cycloaddition reaction such as Diels-Alder reaction, an application of the technique in Diels-Alder reaction of 2(1*H*)-pyridones has been reported only by Matsumoto and his co-workers.⁵

First, we examined Diels-Alder reaction of **Ia-d** with **II**(1.2 equiv.) under atmospheric pressure at 110 °C for 72 h in toluene (Table I). The reactions of **Ib,c** with **II** gave only the *endo*

Dedicated to the memory of Dr. Tetsuji Kametani.

Table I. Diels-Alder Reaction of Ia-d with II under Atmospheric and High Pressure

Conditions



Substrate	R ¹	R ²	R ³	R ⁴	Pressure (kbar)	Yield (%) of Products [Yield (%) of III and IV]
Ia	C ₆ H ₅	H	H	H	atmospheric	0
					10	76 [IIIa (26), IVa (50)]
Ib	H	C ₆ H ₅	H	H	atmospheric	50 [IIIb (50)]
					10	13 [IIIb (11), IVb (2)]
Ic	H	H	C ₆ H ₅	H	atmospheric	90 [IIIc (90)]
					10	96 [IIIc (78), IVc (18)]
Id	H	H	H	C ₆ H ₅	atmospheric	0
					10	84 [IIId (76), IVd (8)]

adducts(**IIIb,c**)⁶ in 50% and 90% yields, however, those of **la,d** with **II** did not yield the Diels-Alder adducts (Table I). Next, we carried out high pressure Diels-Alder reaction of **la-d** with **II** (1.2 equiv.) under 10 kbar at 110 °C for 72 h in toluene (Table I). The reactions of **la,d** with **II** afforded a mixture of the *endo* and *exo* adducts in 76%(**IIIa** and **IVa**)⁶ and 84%(**IIId** and **IVd**)⁶ yields, respectively, which were not obtained under atmospheric pressure conditions (Table I). Furthermore, the reactions of **lb,c** with **II** also proceeded to give a mixture of the *endo* and *exo* adducts in 13% (**IIIb** and **IVb**)⁶ and 96%(**IIIc** and **IVc**)⁶ yields, respectively (Table I). The structures of **IIIa-d** and **IVa-d** were confirmed by their spectral analyses. The configuration of two substituents at 5- and 6- positions in **IIIa-d** ($J_{1,6}=J_{4,5}=4\text{Hz}$) and **IVa-d** ($J_{1,6}=2-2.57\text{Hz}$, $J_{4,5}=2-3.3\text{Hz}$) was determined to be *endo* and *exo* from the coupling constant value in their ¹H-nmr spectra, respectively.

Thus, we found that, by using high pressure technique, Diels-Alder reaction of some 1-methyl-2(1*H*)-pyridones having a phenyl group was accelerated, and also gave the *exo* adducts which were not obtained under atmospheric pressure conditions.

Further investigation for the extension of these reactions is now in progress.

REFERENCES AND NOTES

1. H. Tomisawa, H. Hongo, H. Kato, K. Sato, and R. Fujita, *Heterocycles*, 1981, **16**, 1947.
2. a) H. Tomisawa and H. Hongo, *Chem. Pharm. Bull.*, 1970, **18**, 925; b) H. Tomisawa, R. Fujita, K. Noguchi, and H. Hongo, *ibid.*, 1970, **18**, 941; c) H. Hongo, *ibid.*, 1972, **20**, 226; d) H. Tomisawa, H. Hongo, H. Kato, R. Fujita, and A. Sato, *ibid.*, 1978, **26**, 2312; e) H. Tomisawa, H. Hongo, H. Kato, T. Naraki, and R. Fujita, *ibid.*, 1979, **27**, 670.
3. **Ia**: L. Bauer, C. L. Bell, and G. E. Wright, *J. Heterocycl. Chem.*, 1966, **3**, 393; **Ib**: G. P. Gisby, S. E. Royall, and P. G. Sammes, *J. Chem. Soc., Perkin Trans. 1*, 1982, 169; **Ic**: S. Sugawara and M. Kirisawa, *Pharm. Bull.*, 1955, **3**, 187; **Id**: H. Weber, *Arch. Pharmaz.*, 1975, **308**, 331.
4. M. Takeda, H. Inoue, K. Noguchi, Y. Honma, M. Kawamori, G. Tsukamoto, and S. Saito, *Chem. Pharm. Bull.*, 1976, **24**, 1002.
5. K. Matsumoto, K. Hamada, T. Uchida, and H. Yoshida, *Heterocycles*, 1989, **29**, 21, and references cited therein.

6. All new compounds (**IIIa-d** and **IVa-d**) gave satisfactory spectral and analytical data. ^1H -Nmr spectra were recorded on JEOL PMX-60(60 MHz) (**IIIa-d** and **IVa,d**) or JNM-GSX-400 (400 MHz)(**IVb,c**) spectrometers. Selected data are as follows:
- IIIa** : mp 170-173 °C (benzene); ir (Nujol): 1710, 1670 cm^{-1} ; ^1H -nmr (CDCl_3): δ 2.93 (3H, s), 3.67 (1H, dd, $J=4$, 8 Hz), 3.90 (1H, d, $J=8$ Hz), 4.63 (1H, m), 6.67 (1H, dd, $J=6$, 8 Hz), 6.90 (1H, dd, $J=2$, 8 Hz), 7.03-7.73 (10H, m); ms m/z 358 (M^+).
- IIIb** : mp 227-229 °C (benzene); ir (Nujol): 1710, 1670 cm^{-1} ; ^1H -nmr (CDCl_3): δ 3.00 (3H, s), 3.53 (1H, dd, $J=3$, 8 Hz), 3.77 (1H, dd, $J=4$, 8 Hz), 4.57 (1H, m), 4.73 (1H, dd, $J=4$, 6 Hz), 6.70 (1H, dd, $J=2$, 6 Hz), 6.80-7.70 (10H, m); ms m/z 358 (M^+).
- IIIc** : mp 179-182 °C (benzene); ir (Nujol): 1710, 1675 cm^{-1} ; ^1H -nmr (CDCl_3): δ 3.05 (3H, s), 3.55 (1H, dd, $J=3$, 8 Hz), 3.77 (1H, dd, $J=4$, 8 Hz), 4.15 (1H, dd, $J=3$, 6 Hz), 5.10 (1H, dd, $J=2$, 4 Hz), 6.65 (1H, dd, $J=2$, 6 Hz), 6.78-7.65 (10H, m); ms m/z 358 (M^+).
- IIId** : mp 203-205 °C (ether); ir (Nujol): 1710, 1675 cm^{-1} ; ^1H -nmr (CDCl_3): δ 2.35 (3H, s), 3.53 (1H, dd, $J=4$, 8 Hz), 4.13 (1H, m), 4.20 (1H, d, $J=8$ Hz), 6.60 (1H, dd, $J=6$, 8 Hz), 6.85 (1H, dd, $J=2$, 8 Hz), 7.00-7.90 (10H, m); ms m/z 358 (M^+).
- IVa** : mp 205-207 °C (ether); ir (Nujol): 1710, 1660 cm^{-1} ; ^1H -nmr (CDCl_3): δ 2.97 (3H, s), 3.37 (1H, dd, $J=2$, 8 Hz), 3.77 (1H, d, $J=8$ Hz), 4.73 (1H, m), 6.32 (1H, dd, $J=2$, 8 Hz), 6.70 (1H, dd, $J=6$, 8 Hz), 6.97-7.97 (10H, m); ms m/z 359 (M^++1).
- IVb** : mp 279-282 °C (CH_2Cl_2 -ether); ir (Nujol): 1710, 1670 cm^{-1} ; ^1H -nmr (CDCl_3): δ 2.94 (3H, s), 3.32 (1H, dd, $J=3.30$, 8.43 Hz), 3.36 (1H, dd, $J=2.57$, 8.43 Hz), 4.54 (1H, m), 4.82 (1H, dd, $J=2.57$, 5.86 Hz), 6.78 (1H, dd, $J=1.83$, 5.86 Hz), 7.21-7.26 (2H, m), 7.37-7.53 (8H, m); ms m/z 358 (M^+).
- IVc** : mp 195-197 °C (CH_2Cl_2 -ether); ir (Nujol): 1710, 1675 cm^{-1} ; ^1H -nmr (CDCl_3): δ 2.97 (3H, s), 3.32 (1H, dd, $J=3.30$, 8.43 Hz), 3.36 (1H, dd, $J=2.57$, 8.43 Hz), 4.14 (1H, dd, $J=3.30$, 6.23 Hz), 5.17 (1H, m), 6.75 (1H, dd, $J=2.20$, 6.60 Hz), 7.21-7.26 (2H, m), 7.37-7.50 (8H, m); ms m/z 358 (M^+).
- IVd** : mp 238-241 °C (ether); ir (Nujol): 1710, 1670 cm^{-1} ; ^1H -nmr (CDCl_3): δ 2.65 (3H, s), 3.40 (1H, dd, $J=2$, 8 Hz), 4.02 (1H, d, $J=8$ Hz), 4.20 (1H, m), 6.30 (1H, dd, $J=2$, 8 Hz), 6.70 (1H, dd, $J=6$, 8 Hz), 7.03-8.03 (10H, m); ms m/z 358 (M^+).

Received, 30th August, 1989