

WITHANGULATIN A, A NEW WITHANOLIDE FROM *PHYSALIS ANGULATA*¹


Chiu-Ming Chen,* Zong-Tsi Chen, Chiu-Hsiang Hsieh, Wen-Sen Li, and Say-Yee Wen
 Department of Chemistry, National Tsing Hua University,
 Hsinchu, Taiwan 30043, Republic of China

Abstract — A new withanolide as a topoisomerase II inhibitor, withangulatin A was isolated from the whole herb of *Physalis angulata* L. (Solanaceae). The structure of withangulatin A was established as (20S,22R)-15 α -acetoxy-5 β ,6 β -epoxy-4 β ,14 α -dihydroxy-1-oxowitha-2,16,24-trienolide (I) on the basis of spectroscopic and chemical evidence.

Physalis angulata L., known as an antipyretic, diuretic and antitumor folk-medicine,² is one of the common solanaceous plants in Taiwan.³ Various physalins⁴ and withanolides⁵ together with acetylcholine⁶ and chlorogenic acid⁷ had previously been isolated from this species. A thorough search for the bioactive constituents of the whole herb of this plant has now led to the isolation of a new withanolide, named withangulatin A. This compound was found to act on topoisomerase II to induce topoisomerase II-mediated DNA damage *in vitro*.⁸ The present paper describes the structural elucidation of withangulatin A.


Withangulatin A (I), colorless amorphous powder, mp 152-153°C, $[\alpha]_D^{20} +23.9^\circ$ (c=0.018 in CHCl₃), C₃₀H₃₈O₈ (*m/z* M⁺ found 526.2562, calcd 526.2567) was considered to be a sort of withanolides. Its ir spectrum showed the presence of a hydroxyl (3450 cm⁻¹), an acetoxyl (1735 cm⁻¹), an α , β -unsaturated δ -lactone (1710 cm⁻¹) and an α , β -unsaturated ketone (1690 cm⁻¹) functions, while the uv spectrum showed absorption at λ_{max}^{MeOH} 215 nm (ϵ =16500). The mass spectrum displayed characteristic peaks at *m/z* 526 (5%) M⁺, 466 (16) [M-AcOH]⁺, 451 (30) [M-Me-AcOH]⁺, 433 (7) [M-Me-H₂O-AcOH]⁺, 430 (3) [M-AcOH-2H₂O]⁺ and the base peak at *m/z* 125 corresponding to the δ -lactone moiety.⁹ The ¹H-nmr spectrum (400 MHz, CDCl₃) exhibited not only signals due to two tertiary methyls (δ 1.06, 1.40), a secondary methyl (δ 1.07, *d*, *J* = 7.0 Hz) and two vinyl methyls (δ 1.81, 1.90) but also signals at δ 6.14 (1H, *d*, *J* = 10.0 Hz, -CO-CH=CH-, H-2) and δ 6.97 (1H, *dd*, *J* = 10.0, 5.8 Hz, -CH=CH-CH-OH, H-3), which is regarded as a withanolide fingerprint.¹⁰ Moreover, signals at δ 1.91 (3H, *s*) due to an acetoxy group, and at δ 3.76 (1H, *dd*, *J* = 5.8, 2.0 Hz, H-4), 3.34 (1H, *br s*, H-6), 4.21 (1H, *ddd*, *J* = 12.0, 7.0, 3.0 Hz, H-22) and δ 5.19 (1H, *d*, *J* = 2.7 Hz, H-15) ascribed to four oxymethines were observed, in which the strong deshielding doublet (δ 5.19) due to a proton geminal to a secondary

acetoxyl revealed the adjacency of an olefinic proton (δ 5.63, d , $J = 2.7$ Hz, H-16) (see Table 1). Acetylation of I with acetic anhydride in pyridine afforded an acetate (II), mp 134-136°C, ms m/z 568 (32) M^+ , $\nu_{\text{max}}^{\text{KBr}}$ 3450, 1735 cm^{-1} , which remained a tertiary hydroxy group. Its ^1H -nmr spectrum (Table 1) showed a singlet at δ 2.04 due to a new acetoxyl group and the expected downfield shift of the signal assigned to 4 α -proton from δ 3.76 (dd , $J = 5.8, 2.0$ Hz) in I to δ 4.68 (d , $J = 6.0$ Hz) which indicated the existence of an acetoxyl group with β -orientation at C-4 and no hydrogen on C-5.⁹ Withangulatin A, therefore, is unequivocally characterized by the presence of 5,6-epoxy-4 β -hydroxy-2-en-1-one functionality in rings A and B, a typical α,β -unsaturated δ -lactone in C₉-side chain, the same as in usual withanolides,¹¹ and further contains a trisubstituted double bond, an acetoxyl and a tertiary hydroxy groups. In agreement with this statement, the ^{13}C -nmr spectrum (see Table 2) confirmed the presence of 30 carbons and these substitution patterns,¹² and suggested that withangulatin A (I) may possess a structural resemblance to physapubenolide (III)¹³ and withaminimin (IV).¹⁴ According to two dimensional ^1H - ^1H COSY spectrum of I, the allylic proton signal at δ 2.44 (1H, quintet, $J = 7.0$ Hz, H-20) coupling to an oxymethine proton at δ 4.21 (H-22) and methyl protons at δ 1.07 (H-21) was observed. The COSY spectrum also confirmed that an olefinic proton at δ 5.63 (H-16) and an oxymethine proton at δ 5.19 (H-15) were only coupled to each other with a small coupling constant ($J = 2.7$ Hz), indicating H-15 to be a quasi-axial β -proton.¹⁴ Moreover, the ^1H - ^1H NOESY spectrum showed a cross-peak due to H-4 and H-6, and spatial proximity of H-15 with H-16, H-18 and H-8. These observations along with the above data provided powerful support for β -configuration of 5,6-epoxide, the location and α -configuration of the acetoxyl group at C-15, and the location of the tertiary hydroxy group at C-14 and a double bond at C-16(17). These assignments were further confirmed by comparing the ^{13}C -nmr spectral data of I with those of III and IV as shown in Table 2. Since the 14 α -hydroxy group deshielded C-8 through a β -effect, shielded C-7, C-9 and C-12 through γ -effects, and the 14 β -hydroxy group did not cause the shielding of C-12 in III,¹³ the tertiary hydroxy group at C-14 was oriented α -configuration as the only reasonable assignment.¹²




I R=H

II R=Ac



III



IV

Table 1. ^1H -Nmr Data for Withangulatin A (I) and Withangulatin A Acetate (II) (in CDCl_3)^{a)}

Proton	I	II
2	6.14 (<i>d</i> , 10.0)	6.22 (<i>d</i> , 10.0)
3	6.97 (<i>dd</i> , 10.0, 5.8)	7.04 (<i>dd</i> , 10.0, 6.0)
4 α	3.76 (<i>dd</i> , 5.8, 2.0)	4.68 (<i>d</i> , 6.0)
6 α	3.34 (<i>br s</i>)	3.32 (<i>br s</i>)
7 α	2.53 (<i>dt</i> , 14.0, 3.0)	2.54 (<i>ddd</i> , 14.0, 12.0, 2.0)
7 β	1.66 (<i>ddd</i> , 14.0, 12.0, 2.0)	1.66 (<i>ddd</i> , 14.0, 13.0, 2.0)
8	1.84 (<i>m</i>)	1.82 (<i>m</i>)
9	1.80 (<i>m</i>)	1.80 (<i>m</i>)
15	5.19 (<i>d</i> , 2.7)	5.21 (<i>d</i> , 2.7)
16	5.63 (<i>d</i> , 2.7)	5.67 (<i>d</i> , 2.7)
18*	1.06 (<i>s</i>)	1.07 (<i>s</i>)
19*	1.40 (<i>s</i>)	1.38 (<i>s</i>)
20	2.44 (quintet, 7.0)	2.47 (quintet, 7.0)
21*	1.07 (<i>d</i> , 7.0)	1.10 (<i>d</i> , 7.0)
22	4.21 (<i>ddd</i> , 12.0, 7.0, 3.0)	4.22 (<i>ddd</i> , 12.0, 7.0, 3.0)
23 α	2.17 (<i>dd</i> , 17.2, 3.0)	2.16 (<i>dd</i> , 17.2, 3.0)
23 β	2.37 (<i>br dd</i> , 17.2, 12.0)	2.37 (<i>br dd</i> , 17.2, 12.0)
27*	1.81 (<i>s</i>)	1.83 (<i>s</i>)
28*	1.90 (<i>s</i>)	1.91 (<i>s</i>)
4-OH	3.23 (<i>br s</i>)	
4-OAc*		2.04 (<i>s</i>)
15-OAc*	1.91 (<i>s</i>)	1.91 (<i>s</i>)

a) Chemical shifts are in δ units. Coupling constants (in Hz) are given in parentheses.

*Intensity three protons.

Table 2. ^{13}C -Nmr Data for Withangulatin A (I), Physapubenolide (III) and Withaminimin (IV) (δ ppm)

Carbon	I ^{a)}	III ^{b)}	IV ^{c)}
1	202.44 (C) [202.44 (<i>s</i>)]	202.6	204.06
2	131.42 (CH) [131.37 (<i>d</i>)]	131.5	128.71
3	142.75 (CH) [143.85 (<i>d</i>)]	143.5	141.33
4	69.55 (CH) [69.82 (<i>d</i>)]	69.6	36.03
5	63.23 (C) [63.80 (<i>s</i>)]	63.4	77.22
6	63.23 (CH) [62.84 (<i>d</i>)]	62.7	74.26
7	24.66 (CH ₂) [25.05 (<i>t</i>)]	26.1	26.47
8	34.73 (CH) [35.55 (<i>d</i>)]	40.3	35.43
9	39.32 (CH) [40.06 (<i>d</i>)]	36.1	35.43
10	47.61 (C) [48.05 (<i>s</i>)]	47.9	52.20*
11	21.85 (CH ₂) [22.29 (<i>t</i>)]	21.8	23.18
12	37.54 (CH ₂) [38.26 (<i>t</i>)]	41.4	38.82
13	52.05 (C) [52.36 (<i>s</i>)]	46.1	52.23*
14	81.48 (C) [81.92 (<i>s</i>)]	84.1	82.28
15	83.59 (CH) [84.17 (<i>d</i>)]	80.8	83.36
16	120.80 (CH) [120.95 (<i>d</i>)]	33.8	120.37
17	162.41 (C) [163.04 (<i>s</i>)]	52.3	161.30
18	17.69 (CH ₃) [18.07 (<i>q</i>)]	15.6	16.80
19	15.88 (CH ₃) [16.18 (<i>q</i>)]	17.2	15.10
20	35.39 (CH) [35.70 (<i>d</i>)]	37.5	36.11
21	17.32 (CH ₃) [17.60 (<i>q</i>)]	17.2	17.21
22	79.38 (CH) [79.55 (<i>d</i>)]	78.5	78.47
23	33.11 (CH ₂) [33.40 (<i>t</i>)]	31.2	32.35
24	148.40 (C) [148.40 (<i>s</i>)]	149.5	150.25
25	122.19 (C) [122.23 (<i>s</i>)]	121.9	121.44
26	166.38 (C) [166.31 (<i>s</i>)]	166.8	167.52
27	12.35 (CH ₃) [12.40 (<i>q</i>)]	12.4	12.37
28	20.33 (CH ₃) [19.83 (<i>q</i>)]	20.5	20.64
CH ₃ CO	21.28 (CH ₃) [20.95 (<i>q</i>)]	21.5	21.36
CH ₃ CO	169.65 (C) [169.57 (<i>s</i>)]	169.8	170.63

a) Measured at 100 MHz, in CDCl_3 , and benzene- d_6 in brackets. Assignments were made with the aid of DEPT and OFR (in brackets) experiments.b) Solvent CDCl_3 . Data are from ref. 13.c) Solvent CDCl_3 . Data are from ref. 14. *Values are interchangeable.

The cd spectrum of I showed the positive Cotton effect at 250 ($[\theta] +12600$ in MeOH) and 335 ($[\theta] +4200$ in MeOH) nm, indicating I to have a 22R configuration and A/B *cis*-fused ring junction.¹⁵ Additionally, the aforementioned quintet centered at δ 2.44 with $J_{20-21} = J_{20-22} = 7.0$ Hz of 20B-proton signal¹⁴ indicated a 20S configuration in I, the same as in III, IV and related withanolides. In conclusion, the structure including absolute configuration of withangulatin A was established as (20S,22R)-15 α -acetoxy-5 β ,6 β -epoxy-4 β ,14 α -dihydroxy-1-oxowitha-2,16,24-trienolide (I).

EXPERIMENTAL

Melting points were determined on a Yanaco micro-melting point apparatus and are uncorrected. Optical rotations were measured on a JASCO DIP-360 digital polarimeter. Ir spectra were taken on a Perkin Elmer 781 infrared spectrophotometer. Uv spectra were recorded on a Perkin Elmer Lambda 5 uv/vis spectrophotometer. Mass spectra were recorded on either JEOL JMS-D-100 or JMS-HX-110 spectrometer. ¹H- and ¹³C-nmr spectra were recorded on a Bruker AM-400 nmr spectrometer. Preparative tlc was performed on precoated Kieselgel 60 F₂₅₄ plates (1.0 mm, Merck). Hplc was carried out on a Waters model ALC/GPC-244/M-6000A system with a model 440 absorbance detector (set at 254 nm).

Extraction and Isolation

Dried powdered whole herbs of *Physalis angulata* L. (Solanaceae) (1.0 kg), collected from Nantou (Nantou County, Taiwan) in summer, were extracted with MeOH (9 l) at room temperature for one week and the extract was evaporated *in vacuo* to leave a dark greenish residue (95.0 g). The residue was chromatographed repeatedly on silica gel column with CHCl₃ and an increasing content of acetone as eluents, each fraction being monitored by tlc. One of the fractions eluted with CHCl₃-acetone (5:3) was evaporated to give a residue (1.8 g), which was fractionated sequentially by preparative tlc (CHCl₃-MeOH = 12:1) and by hplc on a μ Bondapak C₁₈ column (EtOAc) to give withangulatin A (I, 58 mg).

Withangulatin A (I)

Colorless amorphous powder, mp 152-153°C, $[\alpha]_D^{20} +23.9^\circ$ (c=0.018 in CHCl₃), hrms m/z : 526.2562 (M^+ calcd for C₃₀H₃₈O₈: 526.2567), 511.2310 (calcd for C₂₉H₃₅O₈: 511.2332); ir $\nu_{\max}^{KBr}$: 3450, 1735, 1710, 1690, 1250, 1235, 1140, 1020, 750 cm⁻¹; uv $\lambda_{\max}^{MeOH}$: 215 nm ($\epsilon=16500$); eims m/z (75 eV, rel. int. %): 526 (5), 511 (7), 484 (2), 466 (16), 451 (30), 433 (7), 430 (3), 273 (15), 260 (75), 125 (100); cd: $[\theta]_{250} +12600$, $[\theta]_{335} +4200$ (c=1.88 $\times 10^{-5}$, MeOH); ¹H-nmr: see Table 1; ¹³C-nmr: see Table 2.

Acetylation of Withangulatin A (I)

A mixture of withangulatin A (I, 12 mg), pyridine (1.0 ml) and acetic anhydride (1.0 ml) was kept at room temperature for 10 h, and treated as usual. The reaction product was purified by preparative tlc (*n*-hexane-EtOAc = 4:1) to give colorless amorphous powder (II, 8 mg), mp 134-136°C, ir $\nu_{\max}^{KBr}$: 3450, 1735, 1710, 1690, 1240, 1110, 1090, 1020, 960 cm⁻¹; eims m/z (75 eV, rel. int. %): 568 (32) M^+ , 553 (17), 508 (11), 448 (14), 320 (18), 260 (35), 125 (100); ¹H-nmr: see Table 1.

ACKNOWLEDGEMENTS

We are grateful to the National Science Council, Republic of China, for financial support of this work, and to Prof. Chang-Sheng Kuoh, Department of Biology, National Cheng Kung University, for authentication of plant material and to Prof. Tian-Shung Wu, Department of Chemistry, National Cheng Kung University, for measurement of cd spectrum.

REFERENCES

1. Part IV on "Constituents of Formosan Antitumor Folk Medicine". Part III: C. M. Chen, Z. T. Chen, and Y. L. Hong, *Phytochemistry*, 1990, **29**, 980.
2. N. Y. Chiu and K. H. Chang, "The Illustrated Medicinal Plants of Taiwan", Southern Materials Center, Inc., Taipei, 1986, Vol. 2, p. 191.
3. Y. C. Liu and C. H. Ou, in "Flora of Taiwan", ed. by H. L. Li, T. S. Liu, T. S. Huang, T. Koyama, and C. E. DeVol, Epoch Publishing Co. Ltd., Taipei, 1978, Vol. IV, p. 535.
4. (a) L. R. Row, K. S. Reddy, N. S. Sarma, T. Matsuura, and R. Nakashima, *Phytochemistry*, 1980, **19**, 1175; (b) L. R. Row, N. S. Sarma, T. Matsuura, and R. Nakashima, *Phytochemistry*, 1978, **17**, 1641; (c) L. R. Row, N. S. Sarma, K. S. Reddy, T. Matsuura, and R. Nakashima, *Phytochemistry*, 1978, **17**, 1647; (d) S. S. Sankara and P. D. Sethi, *Indian J. Pharm.*, 1970, **32**, 163.
5. (a) O. E. Vasina, V. A. Maslennikova, N. D. Abdullaev, and N. K. Abubakirov, *Khim. Prir. Soedin.*, 1986, 596 (*Chem. Abstr.*, 1987, **106**, 135237k); (b) O. E. Vasina, N. D. Abdullaev, and N. K. Abubakirov, *Khim. Prir. Soedin.*, 1987, 856 (*Chem. Abstr.*, 1988, **108**, 201718n).
6. A. Cesario de Mello and P. Afiatpour, *Cienc. Cult. (Sao Paulo)*, 1985, **37**, 799 (*Chem. Abstr.*, 1985, **103**, 85099g).
7. J. Politis, *Compt. rend.*, 1948, **226**, 692 (*Chem. Abstr.*, 1948, **42**, 4648i).
8. J. K. Juang, H. W. Huang, C. M. Chen, and H. J. Liu, *Biochem. Biophys. Res. Commun.*, 1989, **159**, 1128.
9. S. M. Kupchan, W. K. Anderson, P. Bollinger, R. W. Doskotch, R. M. Smith, J. A. S. Renauld, H. K. Schnoes, A. L. Burlingame, and D. H. Smith, *J. Org. Chem.*, 1969, **34**, 3858.
10. A. Bagchi, P. Neogi, M. Sahai, A. B. Ray, Y. Oshima, and H. Hikino, *Phytochemistry*, 1984, **23**, 853.
11. I. Kirson and E. Glotter, *J. Nat. Prod.*, 1981, **44**, 633.
12. H. E. Gottlieb and I. Kirson, *Org. Magn. Reson.*, 1981, **16**, 20.
13. E. Glotter, M. Sahai, and I. Kirson, *J. Chem. Soc., Perkin Trans. 1*, 1985, 2241.
14. H. E. Gottlieb, M. Cojocar, S. C. Sinha, M. Sahai, A. Bagchi, A. Ali, and A. B. Ray, *Phytochemistry*, 1987, **26**, 1801.
15. E. Glotter, I. Kirson, A. Abraham, and D. Lavie, *Tetrahedron*, 1973, **29**, 1353.

Received, 7th May, 1990