

PHOTOINDUCED FRAGMENTATION OF PYRIDO[2,1-f]-as-TRIAZINIUM-4-OLATE AND ITS BENZOLOGUE. A CONTRIBUTION TO THE MECHANISM OF CURTIUS REARRANGEMENT¹


Sándor Bátori,^a András Messmer,^{a*} and Hans-Joachim Timpe^b

a) Central Research Institute for Chemistry, Hungarian Academy of Sciences, H-1525 Budapest, P.O.Box 17. Hungary; b) Technical University "Carl Schorlemmer", Leuna-Merseburg, GDR

Abstract — The zwitterionic pyrido[2,1-f]-as-triazinium-4-olates (**1a-b**) and as-triazino[6,1-a]isoquinolinium-1-olate (**10**) were photolysed to give acid amides (**5**) and (**11**), respectively. Formation of **5** through acylnitrene (**4**) was supported in separate experiment.


Numerous publications deal with the photochemical behaviour of five-membered betaines,² however, the photochemistry of six-membered zwitterions has been studied to a lesser extent.³⁻⁸

The photolysis of s-triazolopyridazinium-4-olates was earlier reported⁹ and it was established that, upon the photolysis in methanol, opening of the pyridazine ring proceeded via the formation of an intermediate diaziridine ring and, subsequently, 3-aminotriazolo derivatives were formed.



As a continuation of our investigations concerning the synthesis and reaction of pyrido [2,1-*f*]-*as*-triazinium-4-olates (1a-b)^{10,11} we examined the photochemical behaviour of these zwitterionic compounds. The photolysis of the 4-olates (1a-b) (2 g of each) was performed in methanol (600 ml) with a medium pressure mercury lamp (750 W, 1.5 h, 20°C) immersed into the solution. A work-up of the photolysed solution by column chromatography (Al₂O₃; ethyl acetate : methanol = 95:5) afforded 2-pyridinecarboxamide (5)^{12a} (58 and 60%, respectively).


In the case of 1a, the formation of HCN was detected^{12b} and in the reaction of 1b (R=C₆H₅) benzonitrile (3b) was isolated in 30% yield. Their formation allows to propose a possible mechanism for the photoinduced process: a fission of the N-N bond¹³ by electronical excitation takes place followed by elimination of RCN (3a-b) from nitrenes (2a-b) to give the same acyl-nitrene (4) which is finally stabilised by hydrogen-abstraction from the solvent yielding 2-pyridinecarboxamide (5).



The supposed existence of 4^{14a} was supported by the photolysis of picolinoyl azide (6)^{14b} which is also a possible source of 4. Thus, azide (6) when photolysed under the same conditions described above, afforded the amide (5) as major product (23%) and urethane (7)¹⁵ as minor product (11%). However, the latter compound (7) was the only product (75%) when 6 (0.7 g) was refluxed in methanol (70 ml; 4.5 h).

As a further example, we synthesized the *as*-triazino[6,1-*a*]isoquinolinium-1-olate (10) (one of the possible three benzologues of olate (1a)) to extend

our studies. Thus, 2-amino-1-ethoxycarbonylisoquinolinium tosylate (8)¹⁶ was treated with formamide in the presence of phosphorous oxychloride (see procedure given for 1a;¹⁰ CAUTION: HCN gas evolution!) to afford 1(2H)-oxo-as-triazino[6,1-a]isoquinolinium perchlorate (9)¹⁷ in 65% yield. This salt was deprotonated by triethylamine to give 1-olate (10)¹⁸ in 70% yield.



The zwitterionic system (10) was photolysed under the same reaction conditions as given for 1a-b to afford 1-isoquinolinecarboxamide (11) in 52% yield (mp 169-171°C; lit.,¹⁹ mp 168-169°C). The reaction proceeded much slower (within 8 h).

The uv spectra of olate (10) and product (11) differ from each other significantly (Figure 1), therefore the photolytic decomposition can be followed by uv-maxima (Figure 2). On this basis, the overall quantum yield was estimated by spectroscopic measurement²⁰ and proved to be $\Phi = 3.1 \times 10^{-3}$ ($\pm 5\%$). This value was found to be stable throughout the photolysis. To the best of our knowledge these findings represent the first photoinduced fragmentation of cyclic zwitterionic systems which - unlike other cyclic zwitterions - did not start with rearrangement but with direct ring opening.

The results of the photolysis of picolinoyl azide (6) and bicyclic olates (1a-b) allow to make a concluding remark on the mechanism of the Curtius rearrangement. Lwowski's study²¹ showed that the Curtius rearrangement of the photolysis of pivaloyl azide could not involve any nitrene intermediate and revealed two simultaneous reaction pathways: (i) formation of isocyanate ("Curtius" and subsequent products), and (ii) formation of a nitrene intermediate and subsequent products which are, however, different from those formed through path (i).




Figure 1. UV spectra of as-triazino[6,1-a]isoquinolinium-1-olate (10) (--) and 1-isoquinolinecarboxamide (11) (—)




Figure 2. Change of the uv spectrum of 10 in the course of photolysis recorded in 15 min intervals (Methanol, $C_0 = 7.1 \times 10^{-5} \text{ mol.l}^{-1}$, $T = 20^\circ\text{C}$)

Similarly, in case of the photolysis of picolinoyl azide (6) we observed the formation of two compounds: (i) N-2-pyridylmethylurethane (7) as the product of reaction of 2-pyridylisocyanate (the result of the Curtius rearrangement) with methanol; and (ii) picolinoyl amide (5) which was obviously formed in reaction of the nitrene intermediate (4) and methanol. Since the photolysis of olates (1a-b) led to the same picolinoyl amide (5) and gave - unlike the analogous reaction of azide (6) - no "Curtius" product (7), we believe that these findings provide a novel support of the synchronous mechanism of the Curtius rearrangement involving no nitrene intermediate as suggested by Lwowski.

REFERENCES AND NOTES

1. Condensed as-Triazines. Part XI. For Part X see ref. 11.
2. This is a well reviewed topic; see e.g.: a) W. D. Ollis and C. A. Ramsden, "Advances in Heterocyclic Chemistry", Vol. 19. ed. by A. B. Katritzky and A. J. Boulton, Academic Press Inc., New York, 1976, p. 1; b) C. G.

- Newton and C. A. Ramsden, Tetrahedron, 1982, 38, 2965; c) A. Lablanche-Combiér "Photochemistry of Heterocyclic Compounds", ed. by O. Buchardt, J. Wiley and Sons, New York, 1976. p. 1.
3. For reviews, see: a) C. A. Ramsden, "Advances in Heterocyclic Chemistry", Vol. 26, ed. by A. R. Katritzky and A. J. Boulton, Academic Press, New York, 1980, p. 1; b) S. T. Reid, ibid., Vol. 30, 1982, p. 230.
4. A. R. Katritzky and H. Wilde, J. Chem. Soc., Chem. Commun., 1975, 1170.
5. Y. Maki, M. Suzuki, T. Furuta, T. Hiramitsu, and M. Kuzuya, Tetrahedron Lett., 1974, 4107.
6. J. M. Dunston and P. Yates, Tetrahedron Lett., 1964, 505.
7. E. F. Ullman and W. A. Henderson, Jr., J. Am. Chem. Soc., 1967, 89, 4390.
8. R. Y. Ning, J. F. Blount, W. Y. Chen, and P. B. Madan, J. Org. Chem., 1975, 40, 2201.
9. a) H. G. O. Becker, D. Nagel, and H.-J. Timpe, Z. Chem., 1973, 13, 213;
b) H.-J. Timpe, R. Burggraf, and U. Lammel, J. prakt. Chem., 1981, 323, 627.
10. S. Bátori, Zs. Juhász-Piedl, P. Sándor, and A. Messmer, J. Heterocycl. Chem., 1986, 23, 375.
11. S. Bátori and A. Messmer, J. Heterocycl. Chem., 1988, 25, 435.
12. a) H. Meyer, Monatsh., 1894, 15, 172.
b) Qualitative observation: benzidine test.
13. A similar fission of the N-N bond followed by formation of benzamide from the resulting acylnitrene has been recently observed in the photochemical reactions of the acyclic trialkylammonio-N-benzoylimides.
a) S. Freeman and M. J. P. Harger, J. Chem. Res. (S), 1988, 192;
b) S. Freeman and M. J. P. Harger, ibid., 1989, 354.
14. a) The attempted trapping of nitrene (4) in benzene and in DMSO solution gave indefinite products.
b) H. Meyer and J. Mally, Monatsh., 1912, 33, 397.

15. mp 132-133°C (methanol). Spectral data of 7: Ir (KBr): 3180, 3130, 3050, 2950, 2820, 1720, 1580, 1530, 1420, 1300, 1210 cm⁻¹; ¹H nmr (CDCl₃): δ 10.60 (s, 1H, NH); 8.41 (d, J_{5,6} = 4.8 Hz, 1H, H-6); 8.13 (d, J_{3,4} = 7.8 Hz, 1H, H-3); 7.72 (dd, J_{3,4} = 7.8 Hz, J_{4,5} = 7.0 Hz, 1H, H-4); 7.00 (dd, J_{4,5} = 7.0 Hz, J_{5,6} = 4.8 Hz, 1H, H-5); 3.88 (s, 3H, CH₃).
16. S. Bátori, Gy. Hajós, P. Sándor and A. Messmer, J. Org. Chem., 1989, 54, 3062.
17. All isolated compounds gave satisfactory elemental analyses, mp 272-274°C (from acetonitrile-ethyl acetate). Spectral data of 9: Ir (KBr): 3080, 3050, 3040, 2750, 2650, 1720, 1640, 1610, 1590, 1070 cm⁻¹; ¹H nmr (TFA): δ 9.10 (d, J_{6,7} = 6.6 Hz, 1H, H-6); 8.82 (d, J_{6,7} = 6.6 Hz, 1H, H-7); 8.80 (s, 1H, H-3); 8.60-8.20 (m, 4H, H-8,9,10,11).
18. mp 199-201°C (acetonitrile). Spectral data of 10: Ir (KBr): 3120, 3090, 3070, 3040, 3010, 1640, 1615, 1550, 1510, 1490, 1450 cm⁻¹; ¹H nmr (DMSO-d₆): δ 8.96 (d, J_{6,7} = 6.8 Hz, 1H, H-6); 8.75 (s, 1H, H-3); 8.66 (d, J_{6,7} = 6.8 Hz, 1H, H-7); 8.55-8.10 (m, 4H, H-8,9,10,11).
19. A. Reissert, Ber., 1905, 38, 3429.
20. H. Baumann, W. Lindenlaub, and H.-J. Timpe, J. prakt. Chem., 1978, 320, 836.
21. a) G. T. Tissue, S. Linke, and W. Lwowski, J. Am. Chem. Soc., 1967, 89, 6303.
b) S. Linke, G. T. Tissue, and W. Lwowski, ibid., 1967, 89, 6303.

Received, 19th November, 1990