

STUDIES IN THE FIELD OF PYRIDAZINE COMPOUNDS, 28.¹ SYNTHESIS OF
5-SUBSTITUTED 6(5H)-1,2,4-TRIAZOLO[4,3-b]PYRIDAZINONES

Judit Kosáry

Department of Chemistry and Biochemistry of the University for Horticulture
and Food Industry, Budapest, P.O. Box 53, H-1502 Hungary

Abstract - The unknown 5-hydroxy-6(5H)-1,2,4-triazolo[4,3-b]pyridazinones (4) were prepared by rearrangement of 6-hydroxyalkoxy-1,2,4-triazolo[4,3-b]pyridazines (1) in polyphosphoric acid. The reaction was accompanied by the formation of bis(triazolopyridazine) derivatives (2). The products (4) were transformed to chloro derivatives (5).

5-Substituted 6(5H)-1,2,4-triazolo[4,3-b]pyridazinones could not be synthesized by direct alkylation of 6-hydroxy-1,2,4-triazolo[4,3-b]pyridazine. According to spectral evidence the latter compound exists only in the hydroxy form and therefore its alkylation by conventional methods afforded only O-alkylated derivatives.² It was reported that 1-hydroxyethoxyphthalazine³ and 3-hydroxyethoxypyridazine⁴ derivatives in the presence of thionyl chloride in chloroform resulted N-chloroethylphthalazinones and pyridazinones by an O→N alkyl rearrangement via an oxazolinium intermediate which was subsequently attacked by the chloride ion. In the case of pyridazine derivatives⁴ the mechanism of the rearrangement was followed by ¹H nmr spectroscopy and an intramolecular process was confirmed.

The synthesis of 1 was described earlier.⁵ 6-Chloro-1,2,4-triazolo[4,3-b]pyridazine reacted with 1,2-ethane- or 1,3-propanediol in the presence of sodium hydride to give 6-hydroxyalkoxy-1,2,4-triazolo[4,3-b]pyridazines (1) and by-products: the bis(triazolopyridazine) derivatives (2) with recovery of the starting material. However, the formation of 2 was suppressed (1-2 % yields) under diluted conditions.⁵

In avoiding direct alkylation we attempted to apply the above O→N rearrangement to 6-hydroxyalkoxy-1,2,4-triazolo[4,3-b]pyridazines (1) but only 6-chloroalkoxy derivatives


(3) were isolated. On the basis of Hückel⁶ and CNDO/2⁷ calculations of 1,2,4-triazolo[4,3-b]pyridazine ring system the electron density of N-5 seems not to be enough to stabilize an oxazolo- or 1,3-oxazinotriazolopyridazinium ion intermediate (n=2 or n=3). It was found that on heating in polyphosphoric acid at 120 °C 1 gave 5-substituted 6(5H)-1,2,4-triazolo[4,3-b]pyridazinones (4) and the bis-derivatives (2) (Scheme 1). On the basis of ir, ¹H nmr and uv data the N- and O-hydroxyalkyl derivatives could be well distinguished. The spectral data of 1b and 4b were compared in detail. Rearrangement of 1b to 4b caused a red shift of the absorption maximum in uv spectra from λ_{\max} 272 nm to 279 nm, smaller than that observed with pyridazine derivatives.⁴ Compound (4b) showed a characteristic carbonyl stretching band at 1660 cm⁻¹. ¹H Nmr spectra also supported the constitutions of 1b and 4b: a Het-NCH₂ triplet at 4.35 ppm (4b) and a Het-OCH₂ triplet at 4.50 ppm (1b) were characteristic (Table).

The low yields of 4 were attributed to the formation of the by-products (2). Poor solubility of compounds 4 in organic solvents made their isolation difficult. Compounds (4) were transformed by thionyl chloride to chloro derivatives (5).



The first step of the rearrangement of 6-hydroxyalkoxy-1,2,4-triazolo[4,3-b]pyridazines (1) to 5-hydroxyalkyl-6(5H)-1,2,4-triazolo[4,3-b]pyridazinones (4) in polyphosphoric acid seems to involve an usual protonation and dehydration process⁸ with the formation of an alkyl cation. The stabilization of the alkyl cation might be carried out in two different ways: "a" - a nucleophilic attack by N-5, hydration and deprotonation to the formation of 4; "b" - a nucleophilic attack by oxygen of another hydroxyalkoxy derivative (1), hydration and deprotonation leads to the formation of 2 and alkanediol (Scheme 2). Heating of the mixture of 1b and 1c in polyphosphoric acid resulted in the formation of only 4b and 4c and no mixed 5-substituted triazolopyridazinones were detected. This crossover reaction suggests that the formation of the compounds (4) is an intramolecular reaction.

EXPERIMENTAL

Melting points were determined on a Boethius apparatus. Melting points are not corrected. The following apparatus were used to obtain spectral data: ir: Perkin-Elmer 577; ¹H nmr: Varian EM-390 90 MHz, using TMS as an internal reference; uv: Cary 1180 instrument.



- a $\tilde{}$ R = H n = 2
- b $\tilde{}$ R = H n = 3
- c $\tilde{}$ R = Me n = 2
- d $\tilde{}$ R = Me n = 3




Scheme 1

Table
 Characteristic ir (KBr, cm^{-1}), ^1H nmr (CDCl_3 , δ ppm) and uv (9% ethanol) data of compounds (1, 2, 3, 4 and 5)

Compd	ir	^1H nmr										uv λ_{max} nm
		ArOCH ₂ t(2H)	NCH ₂ t(2H)	CH ₂ CH ₂ CH ₂ m(2H)	CH ₂ OH t(2H)	CH ₂ Cl t(2H)	H-3 s(1H)	H-7 ABq(2H)	H-8 s(3H)	Me s(3H)		
1a	3235, 3100, 1630, 1550, 860, 840	4.50 (J=6 Hz)	-	-	3.90 (J=6 Hz)	-	9.30	7.05 (J=10 Hz)	8.25 (J=10 Hz)	-	272	
1b	3230, 3100, 1620, 1547, 888, 835	4.50 (J=6 Hz)	-	2.20	3.45 (J=6 Hz)	-	8.90	6.85 (J=10 Hz)	8.02 (J=10 Hz)	-	272	
1c	3227, 3095, 1539, 1502, 889, 870	4.45 (J=6 Hz)	-	-	3.60 (J=6 Hz)	-	9.01	-	8.00 ^a (J=0 Hz)	2.10	271	
1d	3230, 3090, 1600, 1545, 880, 840	4.50 (J=6 Hz)	-	1.80	3.40 (J=6 Hz)	-	8.90	-	8.10 ^a (J=0 Hz)	2.10	270	
2a	3123, 1620, 1548, 858, 825	4.75 ^b (J=0 Hz)	-	-	-	-	8.90 ^c	6.90 ^d (J=10 Hz)	8.01 ^d (J=10 Hz)	-	272	
2b	3100, 1623, 1540, 860, 830	4.75 ^b (J=0 Hz)	-	2.00	-	-	8.90 ^c	6.85 ^d (J=10 Hz)	8.05 ^d (J=10 Hz)	-	273	
2c	3100, 1600, 1570, 858, 830	4.75 ^b (J=0 Hz)	-	-	-	-	9.00 ^c	-	8.00 ^c (J=0 Hz)	2.05 ^e	272	
2d	3110, 1630, 1600, 860, 840	4.80 ^b (J=0 Hz)	-	1.90	-	-	8.90 ^c	-	7.90 ^c (J=0 Hz)	2.10 ^e	270	
3a	3136, 1618, 1547, 887, 831	4.65 (J=5 Hz)	-	-	-	3.90 (J=5 Hz)	8.90	6.90 (J=10 Hz)	8.05 (J=10 Hz)	-	273	
3b	3126, 1622, 1558, 1543, 854, 827	4.52 (J=5 Hz)	-	2.25	-	3.73 (J=5 Hz)	8.95	6.80 (J=10 Hz)	8.02 (J=10 Hz)	-	273	
4a	3230, 3095, 1657, 1625, 1530, 850	-	4.25 (J=7 Hz)	-	4.00 (J=7 Hz)	-	9.30	7.01 (J=10 Hz)	8.15 (J=10 Hz)	-	279	
4b	3235, 3090, 1660, 1610, 1520, 840	-	4.35 (J=7 Hz)	1.80	3.55 (J=6 Hz)	-	9.40	6.75 (J=10 Hz)	8.10 (J=10 Hz)	-	279	
4c	3225, 3090, 1656, 1615, 1530, 850	-	4.45 (J=7 Hz)	-	3.60 (J=7 Hz)	-	9.20	-	8.00 ^a (J=0 Hz)	2.05	279	
5a	3130, 1665, 1610, 1540, 860, 825	-	4.40 (J=6 Hz)	-	-	4.10 (J=6 Hz)	8.90	6.95 (J=10 Hz)	8.10 (J=10 Hz)	-	280	
5b	3126, 1660, 1550, 865, 830	-	4.42 (J=6 Hz)	1.95	-	3.83 (J=5 Hz)	9.00	6.90 (J=10 Hz)	8.02 (J=10 Hz)	-	279	

^as(1H); ^bs(4H); ^cs(2H); ^dABq(4H); ^es(6H)



Scheme 2

Recrystallizations were carried out using ethanol.

Preparation of compounds 1

To a suspension of NaH (0.5 g, 50 % dispersion in mineral oil, 0.01 mol) in 1,2-ethane- or 1,3-propanediol (10 ml), 6-chloro-1,2,4-triazolo[4,3-b]pyridazine (1.54 g, 0.01 mol) was added in small portions at 5 °C. The reaction mixture was stirred for 24 h at room temperature. The precipitate was filtered off. Chromatography on silica gel (eluant ethyl acetate) gave 1 along with 1-2 % of 2.

Compound 1a: mp 172 °C; yield 54 %. Anal. Calcd for $C_7H_8N_4O_2$: C, 46.66; H, 4.48; N, 31.10. Found: C, 46.60; H, 4.50; N, 31.10.

Compound 1b: mp 135 °C; yield 72 %. Anal. Calcd for $C_8H_{10}N_4O_2$: C, 49.48; H, 5.19; N, 28.85. Found: C, 49.36; H, 5.22; N, 28.77.

Compound 1c: mp 180 °C; yield 64 %. Anal. Calcd for $C_8H_{10}N_4O_2$: C, 49.48; H, 5.19; N, 28.85. Found: C, 49.29; H, 5.30; N, 28.80.

Compound 1d: mp 152 °C; yield 70 %. Anal. Calcd for $C_9H_{12}N_4O_2$: C, 51.91; H, 5.81; N, 26.91. Found: C, 51.84; H, 5.80; N, 26.99.

Preparation of compounds 2 and 4

1 (0.01 mol) was heated at 120 °C in polyphosphoric acid (9 g) for 1 h. The reaction mixture was poured onto crushed ice, neutralized with solid $KHCO_3$ and the aqueous solution was extracted with chloroform. Evaporation and chromatography of the solid residue on silica gel (eluant methanol and ethyl acetate 1:9) gave 2 and 4.

Compound 2a: mp 254 °C; yield 21 %. Anal. Calcd for $C_{12}H_{10}N_8O_2$: C, 48.32; H, 3.38; N, 37.57. Found: C, 48.30; H, 3.42; N, 37.70.

Compound 2b: mp 197 °C; yield 19 %. Anal. Calcd for $C_{13}H_{12}N_8O_2$: C, 50.00; H, 3.87; N, 35.88. Found: C, 49.97; H, 3.89; N, 35.75.

Compound 2c: mp 199 °C; yield 20 %. Anal. Calcd for $C_{14}H_{14}N_8O_2$: C, 51.53; H, 4.32; N, 34.34. Found: C, 51.37; H, 4.30; N, 34.40.

Compound 2d: mp 176 °C; yield 5 %. Anal. Calcd for $C_{15}H_{16}N_8O_2$: C, 52.93; H, 4.74; N, 32.93. Found: C, 52.88; H, 4.70; N, 33.07.

Compound 4a: mp 186 °C; yield 15 %. Anal. Calcd for $C_7H_8N_4O_2$: C, 46.66; H, 4.48; N, 31.10. Found: C, 46.62; H, 4.50; N, 31.15.

Compound 4b: mp 166 °C; yield 15 %. Anal. Calcd for $C_8H_{10}N_4O_2$: C, 49.48; H, 5.19; N, 28.85.

Found: C, 49.40; H, 5.25; N, 29.01.

Compound 4c: mp 180 °C; yield 11 %. Anal. Calcd for $C_8H_{10}N_4O_2$: C, 49.48; H, 5.19; N, 28.85.

Found: C, 49.34; H, 5.12; N, 29.10.

Preparation of compounds 3 and 5

To a solution of 1a,b or 4a,b (5 mmol) in chloroform (50 ml), thionyl chloride (3.06 g, 0.026 mol) was added dropwise at 10 °C. The reaction mixture was refluxed for 1 h then poured onto crushed ice, neutralized by 10 % Na_2CO_3 and extracted with chloroform. The solvent was evaporated in vacuo and the residue was triturated with ether.

Compound 3a: mp 118 °C; yield 73 %. Anal. Calcd for $C_7H_7N_4OCl$: C, 42.33; H, 3.55; N, 28.21.

Found: C, 42.20; H, 3.63; N, 28.00.

Compound 3b: mp 97 °C; yield 57 %. Anal. Calcd for $C_8H_9N_4OCl$: C, 45.19; H, 4.27; N, 26.35.

Found: C, 45.01; H, 4.35; N, 26.32.

Compound 5a: mp 128 °C; yield 65 %. Anal. Calcd for $C_7H_7N_4OCl$: C, 42.33; H, 3.55; N, 28.21.

Found: C, 42.20; H, 3.62; N, 28.24.

Compound 5b: mp 104 °C; yield 59 %. Anal. Calcd for $C_8H_9N_4OCl$: C, 45.19; H, 4.27; N, 26.35.

Found: C, 45.10; H, 4.31; N, 26.30.

ACKNOWLEDGEMENT

Thanks are due to Dr. G. Jerkovich and Dr. A. Szabó for recording the spectra and to Mrs. E. Németh for technical assistance.

REFERENCES

1. Part 27.: E. Kasztreiner, K. Czakó, J. Kosáry, E. Diesler, G. Rabloczky, L. Jaszlits and A. Jednákovits, Eur. J. Med. Chem. Chim. Ther., 1991, 26, 739.
2. S. Linholter and R. Rosenorn, Acta Chem. Scand., 1962, 16, 2389.
3. B. G. Pring and C.-G. Swahn, Acta Chem. Scand., 1973, 27, 1891.
4. P. Mátyus, N. Makk, E. Kasztreiner and G. Jerkovich, Heterocycles, 1989, 29, 67.
5. J. Kosáry, Magyar Kém. Folyóirat, 1980, 86, 282 (Chem. Abstr., 1981, 94, 47247s).
6. M. Japelj, B. Stanovnik and M. Tisler, J. Heterocycl. Chem., 1969, 6, 558.
7. J. Kosáry and Z. Gabányi, Chem. Ber., 1983, 116, 3513.
8. R. K. Hill and O. T. Chortyk, J. Am. Chem. Soc., 1962, 84, 1064.

Received, 5th November, 1991