

TOTAL SYNTHESIS OF (+)-MENTHOFURAN[#]

Kozo Shishido,* Takeshi Takata, Koji Umimoto, and Masayuki Shibuya


Faculty of Pharmaceutical Sciences, University of Tokushima

Sho-machi 1, Tokushima 770, Japan

Abstract - Total synthesis of optically active menthofuran, a perfumy monoterpene, has been accomplished employing the intramolecular [3+2] cycloaddition-based methodology for the construction of fused furans.

We have recently devised a new and facile construction of disubstituted fused furans based on the intramolecular [3+2] cycloaddition of nitrile oxide and reported its application to the synthesis of furanosesquiterpene (+)-pallascensin A.¹ Our main objective in this area centers not only on the extension of this methodology to more complex systems but also on the application to the synthesis of biologically active natural products.

In this communication, we wish to report a total synthesis of monoterpene (+)-menthofuran (**1**),² one of the important perfumeries isolated from peppermint oil (*Mentha piperita*),³ starting from (+)-citronellal (**2**) employing the extended strategy for the construction of trisubstituted fused furans (Scheme 1).




Scheme 1

Reduction of (+)-citronellal (**2**) { $[\alpha]_D^{+14.28^\circ}$; lit.,⁴ $[\alpha]_D^{+17.38^\circ}$ } with sodium borohydride followed by protection of the primary alcohol provided the t-butyldimethylsilyl ether (**4**). The allylic hydroxylation of **4** was best carried out in 62 % yield using an excess of t-butyl hydroperoxide in the presence of a catalytic amount of selenium dioxide.⁵ The allyl alcohol thus obtained was

[#] Dedicated to Dr. Masatomo Hamana on the occasion of his 75th birthday

acetylated and desilylated successively to give the alcohol (5), which was then transformed into the oxime (6) by sequential Swern oxidation and oxime formation in 84 % overall yield from 4. Treatment of 6 with 7% aqueous sodium hypochlorite⁶ in methylene chloride at room temperature provided the isoxazoline (7) as an inseparable mixture of diastereoisomers (ca. 8:1 from ¹H nmr) in 76 % yield. Although the exact configuration of the newly developed stereogenic centers of the cycloadduct was not determined by the ¹H nmr, the stereostructure of the major diastereoisomer was tentatively assigned as depicted in 7 from the mechanistic point of view.⁷ Alternatively, the isoxazoline (7) was also assembled by treatment of the nitroalkane (8), prepared from the alcohol (5) via the standard transformations, with *p*-chlorophenyl isocyanate⁸ in 91 % yield. Conversion of 7 into the target molecule was achieved in three-step sequence in 75 % overall yield. Thus, treatment of 7 with lithium hydroxide provided the alcohol (3) which was exposed to the conditions of reductive hydrolysis followed by immediate treatment of the crude product⁹ with *p*-toluenesulfonic acid to give (+)-menthofuran (1)¹¹ { [α]_D+75.45°; lit.,¹⁰ [α]_D+87.46° }, identical with an authentic sample by ¹H nmr and ir spectrometric comparisons (Scheme 2).



Reagents : i, SeO₂, ^tBuOOH ; ii, Ac₂O, pyridine ; iii, ⁿBu₄NF ; iv, (COCl)₂, DMSO, Et₃N ; v, NH₂OH·HCl, AcONa ; vi, 7% aq. NaOCl ; vii, LiOH·H₂O ; viii, Raney Ni, H₂, (MeO)₃B ; ix, PTSA ; x, MsCl, Et₃N ; xi, NaI ; xii, AgNO₂ ; xiii, *p*-ClC₆H₄NCO, Et₃N.

The synthesis described above represents an application of the trisubstituted fused furan construction to the synthesis of (+)-menthofuran. Further work is in progress to extend this methodology to the synthesis of other types of biologically active furanoids.

ACKNOWLEDGEMENT

We are grateful to Professor S. Tsuboi, Okayama University, for his kind gift of authentic menthofuran and to Professor K. Ogasawara, Tohoku University, for providing (+)-citronellal.

REFERENCES AND NOTES

1. K. Shishido, K. Umimoto, and M. Shibuya, Heterocycles, 1990, **31**, 597.
2. R. H. Eastman, J. Am. Chem. Soc., 1950, **72**, 5313.
3. Recent chiral synthesis : (a) R. R. Juo and W. Herz, J. Org. Chem., 1985, **50**, 700; (b) S. G. Hedge, D. Beckwith, R. Doti, and J. Wolinsky, J. Org. Chem., 1985, **50**, 894.
4. S. Takano, T. Sugihara, and K. Ogasawara, Synlett, 1991, 279.
5. M. A. Umbreit and K. B. Sharpless, J. Am. Chem. Soc., 1977, **99**, 5526.
6. G. A. Lee, Synthesis, 1982, 508.
7. K. Shishido, Y. Tokunaga, N. Omachi, K. Hiroya, K. Fukumoto, and T. Kametani, J. Chem. Soc., Perkin Trans. 1, 1990, 2481.
8. A. P. Kozikowski and P. D. Stein, J. Am. Chem. Soc., 1982, **104**, 4023.
9. The tlc of crude product showed two spots. The faster spot is menthofuran and the slower one may be the β,γ -dihydroxy ketone which converged on the former after treatment of the mixture with PTSA.
10. L. H. Zalkow, J. W. Ellis, and S. M. R. Brennan, J. Org. Chem., 1963, **28**, 1705.
11. Colorless oil; bp₂₅ 85 °C (Kugelrohr) (lit.^{2b} bp_{2.8} 48 °C); ¹H nmr (200 MHz, CDCl₃) δ : 1.07 (3H, d, J=6.3 Hz), 1.92 (3H, d, J=1.0 Hz), 2.65 (1H, dd, J=15.5 and 5.3 Hz), 7.03 (1H, br s).

Received, 17th October, 1991