

TWO COMPLEX FLAVONOIDS IN THE FARINOSE EXUDATE OF *PITYROGRAMMA CALOMELANOS*

Fujio Asai^a, Munekazu Iinuma^{b*}, Toshiyuki Tanaka^b and Mizuo Mizuno^b

^aDepartment of Liberal Science, Aichi Gakuin University, Iwasaki Nisshin-cho, Aichi 470-01, Japan. ^bDepartment of Pharmacognosy, Gifu Pharmaceutical University, 6-1 Mitahora-higashi 5 chome, Gifu 502, Japan

Abstract ----- From the farinose exudate of *Pityrogramma calomelanos*, two new complex flavonoids named calomelanols D and E were isolated. Their structures were determined by spectroscopic analysis.

Pityrogramma species (Polypodiaceae) occasionally spout farinose on the surface of frond, and some of them are commonly called as "Gold Fern" or "Silver Fern" by the difference of farinose color.¹ Several new compounds with a novel C6-C3-C6-C3-C6 skeleton (complex flavonoids) were isolated in addition to simple chalcones or dihydrochalcones. The complex flavonoid is regarded as a flavonoid condensed with A neoflavonoid via common phloroglucinol moiety (A ring). Among them, eight complex flavonoids, D-1, D-2a and D-2b in *P. calomelanos*,² T-1, T-2 and T-3 in *P. trifoliata*,³ and X-1 and X-2 in *P. calomelanos* var. *aureoflava*⁴ were isolated and their structures were determined, and some of them were confirmed by synthesis.⁵⁻⁷ Besides these flavones, the presence of minor cognates in the farinose exudate of *P. calomelanos* has been suggested.⁸ To clarify the structures of them, the farinose exudate of *P. calomelanos* in Indonesia was examined. In succession to our preceding paper⁹ reporting three new 8-(3-phenylpropionyl)dihydrochalcones, named calomelanols A, B and C, the isolation and structure determination of other two new complex flavonoids are described in this paper.

Farinose exudate of *P. calomelanos* was rinsed with acetone, and the concentrated extract was purified with silica gel chromatography to lead the isolation of two compounds.

Compound (1), calomelanol D, yellow needles, gave M^+ at m/z 416.0889 in the hrms and the empirical formula corresponds to $C_{24}H_{16}O_7$. The uv spectral data suggested that **1** has a flavonol core, and the behaviors caused by addition of shift reagents were closely similar to those of kaempferol.¹⁰ In the 1H -nmr spectrum, a set of three-proton double doublets at δ 3.05 ($J = 14, 2$ Hz), 3.43 ($J = 14, 7$ Hz) and 5.12 ($J = 7, 2$ Hz) indicated the presence of a partial structure of $PhCHCH_2CO$. Furthermore, a set of two-proton doublets [δ 7.00 and 8.01 (each $J = 9$ Hz)], an aromatic proton singlet (δ 6.56), and five-proton multiplet which is assigned to a mono-substituted benzene ring, and three hydroxyl groups [δ 8.34, 9.06 and 12.30 (chelated)] were observed. In the eims, prominent fragment peaks were observed at m/z 373, 339 and 121 (Figure 1). These results showed that the partial structure, $PhCHCH_2CO$, was fused with a kaempferol skeleton on the A ring. The behavior of uv and the appearance of a chelated hydroxyl group in 1H -nmr




Figure 1

spectra permitted two possible structures (**1a**) or (**1b**) (Figure 2). The confirmation of structure for **1** was depended on 2D nmr experiments. That is, in the 1H - ^{13}C COSY (Figure 3), an aromatic singlet proton caused a cross peak with a carbon signal at δ 99.8. In the long range 1H - ^{13}C COSY (Figure 3), the carbon at





Figure 2

Figure 3



δ 99.8 also caused a cross peak with a chelated hydroxyl group. Consequently, **1a** was preferable to **1b** as a structure of calomelanol D. The stereochemistry of C-7'' is now being investigated.

Compound (2), calomelanol E, a colorless powder, gave M^+ m/z 418.1083 in the hrms, and the empirical formula is $C_{24}H_{18}O_7$. In the 1H -nmr spectrum, two sets of three-proton double doublets were observed [δ

2.89 ($J=17, 2$ Hz), 3.25 ($J=17, 6$ Hz), 4.48 ($J=6, 2$ Hz); δ 2.81 ($J=17, 3$ Hz), 3.39 ($J=17, 13$ Hz), 5.61 ($J=13, 3$ Hz)]. The former set suggested the presence of a partial structure of PhCHCH_2CO . The latter was assigned to H-3 and H-2 of a flavanone skeleton. Furthermore, the spectrum also showed the presence of two *p*-substituted benzene rings [δ 6.71, 6.92 (each $J=8$ Hz) and 6.82, 7.35 (each $J=9$ Hz)], an aromatic proton (δ 6.37) and three hydroxyl groups [δ 9.40, 9.65 and 12.41 (chelated)]. In the eims, significant fragments were appeared at m/z 299, 298, 257 and 120. The fragment ions could be drawn as in Scheme 1.



These spectral data mentioned above suggested that the structure of **2** was either **2a** or **2b** shown in Figure 4. In the ^1H - ^{13}C COSY, an aromatic proton based on the A ring caused a cross peak with a carbon at δ 96.1. In the long range ^1H - ^{13}C long COSY, a chelated hydroxyl group caused cross peaks with three quaternary carbons assigned to C-5 (δ 159.0), C-6 (δ 105.9) and C-10 (δ 104.8) (Figure 5). Consequently, the structure of calomelanol E was determined to be **2a**.



EXPERIMENTAL

Plant material and extraction were described in preceding paper.⁸

Isolation of compounds 1 and 2. A more polar fraction containing D-1 and calomelanols A-C was purified by silica gel column chromatography (eluent: *n*-hexane-acetone = 3:1 and C₆H₆-EtOAc = 5:1) and recrystallization to give **1** (45 mg) and **2** (60 mg) in pure form.

Compound 1 (calomelanol D): mp 235 °C (dec.) (MeOH); yellow needles; $[\alpha]_D^{20}$ (c 0.06, MeOH); ir (KBr) 1760, 1660, 1630, 1618 cm⁻¹; hrms *m/z* 416.0889 (Calcd 416.0869 for C₂₄H₁₆O₇); eims *m/z* (rel. int.): 416 (M⁺, 100), 388 (18), 373 (38), 339 (22), 241 (1), 239 (7), 208 (4), 205 (7), 184 (11), 121 (24); uv (nm, MeOH): 270, 325sh, 373, +NaOMe: 283, 325, 418, +AlCl₃: 265, 295sh, 360, 428; ¹H-nmr (acetone-*d*₆) δ: 3.05 (1H, dd, *J* = 14, 2 Hz, H-8"), 3.43 (1H, dd, *J* = 14, 7 Hz, H-8"), 5.12 (1H, br dd, *J* = 7, 2 Hz, H-7"), 6.56 (1H, s, H-6), 7.00 (2H, d, *J* = 9 Hz, H-3' and 5'), 7.20-7.28 (5H, m, H-2"-6"), 8.01 (2H, d, *J* = 9 Hz, H-2' and 6'), 8.34, 9.06 (1H each, br s, OH), 12.30 (1H, s, C5-OH); ¹³C-nmr (acetone-*d*₆): all carbons were assigned by the aids of C-H and long range C-H COSY δ: 148.2 (C-2), 137.6 (C-3), 177.1 (C-4), 161.4 (C-5), 99.8 (C-6), 158.0 (C-7), 105.1 (C-8), 153.3 (C-9), 107.4 (C-10), 123.1 (C-1'), 130.7 (C-2' and 6'), 116.6 (C-3' and 5'), 160.6 (C-4'), 142.5 (C-1"), 127.7 (C-2" and 6"), 130.1 (C-3" and 5"), 128.3 (C-4"), 35.8 (C-7"), 37.7 (C-8"), 166.0 (C-9").

Compound 2 (calomelanol E): mp 265 °C (dec.) (MeOH); a colorless powder, $[\alpha]_D^{25}$ (c 0.06, MeOH); ir (KBr): 1760, 1640, 1590 cm⁻¹, hrms *m/z* 418.1083 (Calcd 418.1053 for C₂₄H₁₈O₇); eims *m/z* (rel. int.): 418 (M⁺, 100), 402 (20), 390 (9), 375 (7), 325 (9), 324 (8), 323 (7), 299 (22), 298 (18), 297 (15), 283 (9), 280 (29), 270 (29), 257 (11), 255 (20), 252 (33), 205 (38), 147 (11), 120 (53), 107 (22); uv (nm, MeOH): 280, 340, +NaOMe: 285sh, 332, +AlCl₃: 280, 340; ¹H-nmr (acetone-*d*₆) δ: 2.92 (4H, m, H-3 and 8"), 4.59 (1H, br dd, *J* = 6, 2 Hz, H-7"), 5.62 (1H, dd, *J* = 13, 3 Hz, H-2), 6.27 (1H, s, H-8), 6.77 (2H, d, *J* = 8 Hz, H-3" and 5'), 6.92 (2H, d, *J* = 9 Hz, H-3' and 5'), 7.02 (2H, d, *J* = 8 Hz, H-2" and 6"), 7.43 (2H, d, *J* = 9 Hz, H-2' and 6'), 8.30, 8.58 (1H each, br s, OH), 12.49 (1H, s, C5-OH); (DMSO-*d*₆) (assigned by the H-H COSY) δ: 2.81 (1H, dd, *J* = 17, 3 Hz, H-3_{eq}), 2.89 (1H, dd, *J* = 17, 2 Hz, H-8"), 3.25 (1H, dd, *J* = 17, 6 Hz, H-8"), 3.39 (1H, dd, *J* = 17, 13 Hz, H-3_{ax}), 4.48 (1H, br dd, *J* = 6, 2 Hz, H-7"), 5.61 (1H, dd, *J* = 13, 3 Hz, H-2), 6.37 (1H, s, H-8), 6.71 (1H, d, *J* = 8 Hz, H-3" and 5"), 6.82 (2H, d, *J* = 9 Hz, H-3' and 5'), 6.92 (2H, d, *J* = 8 Hz, H-2" and 6"), 7.35 (2H, d, *J* = 9 Hz, H-2' and 6'), 9.40 (1H, br s, C4'-OH), 9.65 (1H, br s, C4'-OH), 12.41 (1H, s, C5-OH); ¹³C-nmr (DMSO-*d*₆): all carbons were assigned by the

aids of C-H and long range C-H COSY) δ : 78.7 (C-2), 41.9 (C-3), 198.2 (C-4), 159.0 (C-5), 105.9 (C-6), 158.4 (C-7), 96.1 (C-8), 161.4 (C-9), 104.8 (C-10), 128.3 (C-1'), 128.5 (C-2' and 6'), 115.2 (C-3' and 5'), 157.9 (C-4'), 131.2 (C-1''), 127.5 (C-2'' and 6''), 115.5 (C-3'' and 5''), 156.4 (C-4''), 32.4 (C-7''), 36.7 (C-8''), 166.7 (C-9'').

REFERENCES

1. D. L. Jones, 'Encyclopedia of Ferns', Lothian Publishing Company (PTY) LTD, Melbourne, Sydney, Auckland, 1987, p. 8.
2. H. Wagner, O. Seligmann, O. Chari, E. Wollenweber, V. H. Dietz, D. M. X. Donnelly, M. J. Meegan, and B. O'Donnell, *Tetrahedron Lett.*, **1979**, 4296.
3. V. H. Dietz, E. Wollenweber, J. Favre-Bonvin, and L. D. Gometz, *Z. Naturforsch.*, **1980**, **53c**, 36.
4. J. Favre-Bonvin, M. Jay, E. Wollenweber, and V. H. Dietz, *Phytochemistry*, **1980**, **19**, 2043.
5. M. Iinuma, S. Matsuura, and F. Asai, *Heterocycles*, **1983**, **20**, 1923.
6. M. Iinuma, K. Hamada, M. Mizuno, F. Asai, and E. Wollenweber, *Z. Naturforsch.*, **1986**, **41c**, 681.
7. F. Asai, M. Iinuma, T. Tanaka, and M. Mizuno, *Chem. Pharm. Bull.*, **1990**, **38**, 1079.
8. E. Wollenweber and V. H. Dietz, *Biochem. System. Ecol.*, **1980**, **8**, 22.
9. F. Asai, M. Iinuma, T. Tanaka, and M. Mizuno, *Phytochemistry*, **1991**, **30**, 3091.
10. T. J. Mabry, K. R. Markham, and M. B. Thomas, 'The Systematic Identification of Flavonoids', Springer-Verlag, Berlin, Heidelberg, New York, 1970, pp. 41-103.

Received, 18th October, 1991