

THERMAL REACTION OF BENZIMIDAZOLIUM *N*-ALLYLIDES†

Yoshiro Matsuda*, Makoto Yamashita, Kimitoshi Takahashi,
Shizuki Ide, Katsura Torisu, and Kazuki Furuno

*School of Pharmaceutical Sciences, Nagasaki University,
1-14 Bunkyo-machi Nagasaki 852, Japan*


Abstract – The reaction of benzimidazolium salt (**3**) and polarized olefins (**1a,b,d** and **2c**) with K_2CO_3 in $CHCl_3$ gave the corresponding benzimidazolium *N*-allylides (**4a-d**). Thermolyses of *N*-allylides (**4a,b**) in refluxing xylene afforded the 1,5-dipolar cyclization product, benzopyrroloimidazole (**5a**), whereas heating of *N*-allylides (**4c,d**) resulted in back-donated 1,6-cyclization to give the mesomeric betaines, benzimidazopyridiniumides (**7a,b**). Treatment of **3** and **2a** with K_2CO_3 in $CHCl_3$ yielded directly benzopyrroloimidazole (**5b**), while similar treatment of **3** with **1c** or **2b** gave benzopyrrolopyrazines (**6a,b**).

1,5-Dipolar cyclization of *N*-allylides or *N*-vinylimino ylides, which were obtained by the reaction of *N*-ylides or *N*-imines with acetylenes and polarized olefins (ketene dithioacetals and ethoxymethylene compounds) is useful for the syntheses of heterocyclic compounds.¹⁻⁹ With regard to benzimidazolium *N*-ylides, 1,5-dipolar cyclization was observed for the reaction of benzimidazolium *N*-ylides with acetylenes, giving the corresponding benzopyrroloimidazoles or benzopyrrolo-

† Dedicated to Prof. Masatomo Hamana on the occasion of his 75th birthday.


pyrazines *via* benzimidazolium *N*-allylides.⁹ However, the literature was devoid of the reaction of benzimidazolium *N*-ylides with polarized olefins. As part of our continuing interest in ylide chemistry, we briefly reported on the synthesis of the mesomeric betaine, benzimidazo[1,2-*a*]pyridiniumide (7b) which was obtained by the reaction of benzimidazolium salt (3) with diethyl ethoxymethylenemalonate (2c) in the presence of K₂CO₃ *via* a novel back-donated 1,6-cyclization.¹⁰ The purpose of the present investigation is to see whether benzimidazolium *N*-allylides (4) undergo 1,5-dipolar cyclization to afford benzopyrrolo[1,2-*a*]imidazoles (5) or back-donated 1,6-cyclization to give the mesomeric betaines, 4-oxobenzimidazo[1,2-*a*]pyridiniumides (7). The polarized olefins (1,2) used in the present work were shown below. (Scheme 1)

Scheme 1




The reaction of benzimidazolium salt (3) with polarized olefins (1a,b) in the presence of K₂CO₃ in CHCl₃ for three days gave benzimidazolium *N*-allylides (4a,b) in good yield. A solution of the *N*-allylide (4a) in xylene was refluxed for 24 h to afford benzopyrroloimidazole (5a), a 1,5-dipolar cyclization product in moderate yield. In our previous paper,¹¹ we reported that thermolysis of imidazolium *N*-(3,3-dicyano-1-ethoxycarbonyl-2-methylthio)allylide afforded the back-donated 1,6-cyclization product, 7-iminoimidazopyridiniumide derivative. However, heating of benzimidazolium *N*-allylide (4b) in refluxing xylene gave the 1,5-dipolar cyclization product (5a). Treatment of 3 with polarized olefin (2a) in CHCl₃ did not give *N*-allylide but afforded directly benzopyrroloimidazole (5b) in 25% yield. Interestingly, the reaction of 3 with 1c or 2b gave benzopyrrolo[1,2-*a*]pyrazine derivatives (6a,b). The reaction of 3 with 1d in the presence of K₂CO₃ in CHCl₃ gave benzimidazolium *N*-allylide (4c). As reported in our previous paper,¹⁰ when the salt (3) and 2c were

treated with K_2CO_3 in $CHCl_3$ for a week, the mesomeric betaine, benzimidazopyridiniumide (**7b**) was obtained in 16% yield. However, the mixture of **3** and **2c** with K_2CO_3 in $CHCl_3$ was stirred at room temperature for three days to give *N*-allylide (**4d**) in 68% yield. It should be noted that heating of **4c,d** in refluxing xylene resulted in the back-donated 1,6-cyclization giving rise to the mesomeric betaines (**7a,b**) in 87% and 64% yields, respectively. (Scheme 2)




The formation of compounds (**5a,6a**) may be rationalized as outlined in Scheme 3. In the case of benzopyrroloimidazole (**5a**), as described in the previous papers,¹⁻¹⁰ the initial step may be 1,5-cyclization to give **9**. This step is then followed by elimination of the phenylsulfonyl group that leads to **5a**, whereas, as pointed out by Meth-Cohn,⁹ the resonance structure (**8**) may cyclize to give benzopyrrolopyrazine (**6a**) *via* **9'**. The formation of benzimidazopyridiniumide (**7a**) may be clarified as described in our previous paper.¹⁰ The resonance structure (**11**) may cyclize to give **7a** *via* **12**. (Scheme 4)

Scheme 3



Scheme 4



EXPERIMENTAL

Melting points were determined with a Mitamura Mel-Temp and are uncorrected. Ir spectra were recorded as KBr pellets on a JASCO IRA-2 spectrophotometer. Uv spectra were recorded on a Hitachi 323 spectrophotometer. ^1H -Nmr spectra were obtained on a JNM-FX-90Q (90 MHz) spectrometer with tetramethylsilane as internal standard. Chemical shifts are reported in parts per million (δ). Elemental analyses (C,H,N) of all compounds described here were performed on a Yanagimoto MT-2 CHN recorder.

Benzimidazolium *N*-(1-ethoxycarbonyl)allylides (4a-d)

A mixture of **3** (1.20 g, 4 mmol), polarized olefins (**1a,b,d,2c**) (4 mmol) and K_2CO_3 (1.10 g, 8 mmol) in CHCl_3 (40 ml) was stirred at room temperature for three days and the mixture was then evaporated under reduced pressure. To the residue was added ice-water (100 ml). The precipitate was filtered, washed with water, dried and recrystallized from CHCl_3 -MeOH to give products (**4a-d**), respectively.

4a: mp 196°C ; yield 1.44 g (79%); ir(KBr) ν_{max} 2150(CN), 1660(CO) cm^{-1} ; uv(EtOH) λ_{max} (log ϵ) 243(4.06), 272(3.99), 279(3.99), 303(3.93), 381(4.27) nm; ^1H -nmr (CDCl_3) δ 1.24(3H, t, $J=7$ Hz, CH_2CH_3), 2.72(3H, s, SCH₃), 4.08(3H, s, NCH₃), 4.11(2H, q, $J=7$ Hz, CH_2CH_3), 7.25-7.78(9H, m, Ar-H), 9.57(1H, s, C₂-H). *Anal.* Calcd for $\text{C}_{22}\text{H}_{21}\text{N}_3\text{O}_4\text{S}_2$: C, 58.00; H, 4.65; N, 9.22. Found: C, 57.89; H, 4.68; N, 9.29.

4b: mp 180°C ; yield 0.72 g (53%); ir(KBr) ν_{max} 2190(CN), 2170(CN), 1660(CO) cm^{-1} ; uv(EtOH) λ_{max} (log ϵ) 242(3.65), 265(3.59), 272(3.65), 280(3.65), 302(3.55), 370(4.06) nm; ^1H -nmr (CDCl_3) δ 1.24(3H, t, $J=7$ Hz, CH_2CH_3), 2.61(3H, s, SCH₃), 4.13(3H, s, NCH₃), 4.19(2H, q, $J=7$ Hz, CH_2CH_3), 7.27-7.70(4H, m, Ar-H), 8.74(1H, s, C₂-H). *Anal.* Calcd for $\text{C}_{17}\text{H}_{16}\text{N}_4\text{O}_2\text{S}$: C, 59.98; H, 4.74; N, 16.46. Found: C, 59.88; H, 4.74; N, 16.37.

4c: mp 185°C ; yield 1.38 g (93%); ir(KBr) ν_{max} 2170(CN), 1680(CO), 1660(CO) cm^{-1} ; uv(EtOH) λ_{max} (log ϵ) 203(4.16), 235sh(3.84), 263(3.59), 276(3.59), 289(3.59) nm; ^1H -nmr (CDCl_3) δ 1.22(3H, t, $J=7$ Hz, CH_2CH_3), 2.46(3H, s, SCH₃), 3.50(3H, s, OCH₃), 4.14(2H, q, $J=7$ Hz, CH_2CH_3), 7.40-7.72(4H, br s, Ar-H), 8.92(1H, s, C₂-H). *Anal.* Calcd for $\text{C}_{18}\text{H}_{19}\text{N}_3\text{O}_4\text{S}$: C, 57.90; H, 5.13; N, 11.25. Found: C, 57.87; H, 5.11; N, 11.14.

4d: mp 200°C; yield 1.06 g (68%); ir(KBr) $\nu_{\max}$ 1690(CO), 1660(CO), 1640(CO) cm^{-1} ; uv(EtOH) $\lambda_{\max}$ (log ϵ) 264(4.13), 270(4.13), 278(4.04), 344(4.56) nm; ^1H -nmr(CDCl_3) δ 0.95(6H, t, $J=7$ Hz, $2\times\text{CH}_2\text{CH}_3$), 1.28(3H, t, $J=7$ Hz, CH_2CH_3), 3.73(4H, q, $J=7$ Hz, $2\times\text{CH}_2\text{CH}_3$), 4.07(3H, s, NCH_3), 4.19(2H, q, $J=7$ Hz, CH_2CH_3), 7.39-7.49(4H, m, Ar-H), 8.57(1H, s, C2-H), 8.59(1H, s, C2'-H). *Anal.* Calcd for $\text{C}_{20}\text{H}_{24}\text{N}_2\text{O}_6$: C, 61.85; H, 6.23; N, 7.21. Found: C, 61.84; H, 6.20; N, 7.24.

3-Cyano-1-ethoxycarbonyl-4-methyl-2-methylthio-4H-benzo[b]pyrrolo-[1,2-a]imidazole (5a)

A solution of **4a** or **4b** (2 mmol) in xylene (60 ml) was refluxed for 24 h and the solution was then evaporated under reduced pressure. To the residue was added ice-water (50 ml). The precipitate was filtered, washed with water, dried and recrystallized from EtOH to give **5a**.

5a: mp 180°C; yield 0.40 g (64%) from **4a** (0.91 g), 0.36 g (57%) from **4b** (0.68 g); ir(KBr) $\nu_{\max}$ 2200(CN), 1670(CO) cm^{-1} ; uv(EtOH) $\lambda_{\max}$ (log ϵ) 217(4.32), 245(4.54), 267(4.18), 282(4.37), 305(4.06), 331(4.21) nm; ^1H -nmr(CDCl_3) δ 1.48(3H, t, $J=7$ Hz, CH_2CH_3), 2.74(3H, s, SCH_3), 3.94(3H, s, NCH_3), 4.47(2H, q, $J=7$ Hz, CH_2CH_3), 7.15-7.40(3H, m, Ar-H), 8.59-8.70(1H, m, C8-H). *Anal.* Calcd for $\text{C}_{16}\text{H}_{15}\text{N}_3\text{O}_2\text{S}$: C, 61.32; H, 4.82; N, 13.41. Found: C, 61.53; H, 4.88; N, 13.33.

1-Ethoxycarbonyl-3-methoxycarbonyl-4-methyl-4H-benzo[b]pyrrolo-[1,2-a]imidazole (5b)

A mixture of **3** (1.20 g, 4 mmol), **2a** (0.52 g, 4 mmol), and K_2CO_3 (1.10 g, 8 mmol) in CHCl_3 (40 ml) was stirred at room temperature for 6 days and the mixture was added to ice-water (100 ml) and extracted with CHCl_3 (2×30 ml). The combined extracts were washed with water (50 ml), dried (Na_2SO_4), and evaporated under reduced pressure. The residue was submitted to column chromatography on silica gel. From a benzene- CHCl_3 (20:1) fraction, product (**5b**) was obtained.

5b: mp 138°C; yield 0.30 g (25%); ir(KBr) $\nu_{\max}$ 1700(CO), 1680(CO) cm^{-1} ; uv(EtOH) $\lambda_{\max}$ (log ϵ) 214(4.36), 248(4.47), 277(4.12), 293(4.37), 320(4.22), 330(4.34) nm; ^1H -nmr(CDCl_3) δ 1.41(3H, t, $J=7$ Hz, CH_2CH_3), 3.87(3H, s, OCH_3), 4.22(3H, s, NCH_3), 4.37(2H,

q, $J=7$ Hz, CH_2CH_3), 7.19-7.46(3H, m, Ar-H), 7.67(1H, s, $\text{C}_2\text{-H}$), 8.75-8.86(1H, m, $\text{C}_8\text{-H}$). *Anal.* Calcd for $\text{C}_{16}\text{H}_{16}\text{N}_2\text{O}_4$: C, 63.99; H, 5.37; N, 9.33. Found: C, 63.83; H, 5.39; N, 9.23.

4,5-Dihydro-5-methyl-4-oxobenzo[*c*]pyrrolo[1,2-*a*]pyrazines (6a,b)

Benzopyrrolopyrazines (6a,b) were prepared by the reaction of 3 (1.20 g, 4 mmol) with polarized olefins (1c,2b) (4 mmol) in the presence of K_2CO_3 (1.10 g, 8 mmol) in CHCl_3 (40 ml) using procedure above for the synthesis of 5b. From a benzene- CHCl_3 (20:1) fraction, products (6a,b) were obtained, respectively.

6a: mp 236°C; yield 1.08 g (93%); ir(KBr) ν_{max} 1660(CO) cm^{-1} ; uv(EtOH) λ_{max} (log ϵ) 220(4.43), 245 (4.23), 275(4.20), 327(4.18), 340(4.08) nm; $^1\text{H-nmr}$ (CDCl_3) δ 2.61(3H, s, SCH_3), 3.67(3H, s, NCH_3), 7.20-7.78(4H, m, Ar-H), 8.45(1H, s, $\text{C}_1\text{-H}$). *Anal.* Calcd for $\text{C}_{13}\text{H}_{11}\text{N}_3\text{O}_3\text{S}$: C, 53.96; H, 3.80; N, 14.53. Found: C, 53.86; H, 3.90; N, 14.48.

6b: mp 292°C; yield 0.65 g (72%); ir(KBr) ν_{max} 2220(CN), 1650(CO) cm^{-1} ; uv(EtOH) λ_{max} (log ϵ) 240(4.41), 260(3.83), 302(3.63), 313(3.79), 326(3.68) nm; $^1\text{H-nmr}$ (CDCl_3) δ 3.68(3H, s, NCH_3), 7.26-8.05(6H, m, Ar-H). *Anal.* Calcd for $\text{C}_{13}\text{H}_9\text{N}_3\text{O}$: C, 69.95; H, 4.06; N, 18.82. Found: C, 70.22; H, 4.22; N, 18.62.

1,4-Dihydro-1-ethoxycarbonyl-5-methyl-4-oxobenz[*b*]imidazo[1,2-*a*]-pyridin-9b-ium-1-ides (7a,b)

Compound (7a,b) was obtained by heating of 4c,d (2 mmol) using procedure above for the synthesis of 5a.

7a: mp 247°C; yield 87%; ir(KBr) ν_{max} 2200(CN), 1710(CO) cm^{-1} ; uv(EtOH) λ_{max} (log ϵ) 230(4.29), 263(4.27), 280(4.38), 364(3.93), 382(4.19), 404(4.23) nm; $^1\text{H-nmr}$ (CDCl_3) δ 1.48(3H, t, $J=7$ Hz, CH_2CH_3), 2.58(3H, s, SCH_3), 4.60(2H, q, $J=7$ Hz, CH_2CH_3), 4.69(3H, s, NCH_3), 7.46-7.79(4H, m, Ar-H). *Anal.* Calcd for $\text{C}_{17}\text{H}_{15}\text{N}_3\text{O}_3\text{S}$: C, 59.81; H, 4.43; N, 12.31. Found: C, 59.77; H, 4.47; N, 12.31.

7b¹: mp 278°C; yield 64%.

REFERENCES

1. K. Matsumoto, *J. Syn. Org. Chem. Japan*, 1974, **32**, 731.
2. T. Uchida and K. Matsumoto, *Synthesis*, 1976, 209.
3. F. J. Swinbourne, J. H. Hunt, and G. Klinkert, "Advances in Heterocyclic Chemistry", Vol. 23, ed. by A. R. Katritzky and A. J. Boulton, Academic Press, New York, 1978, p. 103.
4. R. M. Acheson and N. F. Flmore, *ibid.*, Vol. 23, 1978, p. 263.
5. Y. Tamura and M. Ikeda, *ibid.*, Vol. 29, 1981, p. 71.
6. E. C. Taylor and I. J. Turchi, *Chem. Rev.*, 1979, **79**, 181.
7. Y. Tamura, Y. Sumida, S. Haruki, and M. Ikeda, *J. Chem. Soc., Perkin Trans. I*, 1975, 575.
8. Y. Tominaga and Y. Matsuda, *J. Syn. Org. Chem. Japan*, 1985, **43**, 669.
9. O. Meth-Cohn, *Tetrahedron Lett.*, 1975, 413.
10. Y. Matsuda, H. Gotou, K. Katou, H. Matsumoto, M. Yamashita, K. Takahashi, and K. Ide, *Heterocycles*, 1990, **31**, 997.
11. Y. Matsuda, H. Gotou, K. Katou, H. Matsumoto, M. Yamashita, K. Takahashi, K. Ide, K. Furuno, and K. Torisu, *Heterocycles*, 1991, **32**, 2217.

Received, 25th October, 1991