

SYNTHETIC CONNECTIONS TO THE DIRECTED *ortho* METALATION REACTION. 3,4-PYRIDINES FROM 4-TRIALKYLSILYL-3-PYRIDYL TRIFLATES


Masao Tsukazaki and Victor Snieckus*

Guelph-Waterloo Centre for Graduate Work in Chemistry, University of Waterloo, Waterloo, Ontario, Canada N2L 3G1

Abstract— 4-Trialkylsilyl-3-pyridyl triflates (**11a-b**), derived from **9** and **12** by directed *ortho* metalation chemistry, serve as useful precursors of 3,4-pyridynes (**6a-b**) and lead by cycloaddition and nucleophilic trapping reactions to products (**13a-b**, **14a-b**, **15**, **16a-b**, and **17a-b**).

In the field of heteraryne chemistry,¹ the 3,4-pyridyne species has attracted considerable theoretical² and synthetic³ interest. The classical methods of 3,4-pyridyne generation *via* thermolysis of diazonium carboxylate⁴ and oxidative degradation of aminotriazole⁵ have been supplemented by new procedures based on the original observations of halopyridine metal-halogen exchange and directed *ortho* metalation by Kauffmann,⁶ Gribble,^{3a} and Quéguiner.^{3b-c,7} Noteworthy is the lack of general approaches for substituted 3,4-pyridynes which is mainly due to poor accessibility of appropriately substituted pyridine precursors and/or their instability. The insightful studies by Kobayashi and coworkers⁸ on the fluoride-induced generation of benzyne from *o*-silylaryl triflates⁹ triggered our work¹⁰ on the generation of substituted benzyne (**2**), derived from precursors (**1**) *via* directed *ortho* metalation strategies,¹¹ and their cycloaddition (**3**) and nucleophilic (**4**) reactivities (Scheme 1).

Scheme 1




This paper is dedicated to Professor Emeritus Masatomo Hamana, a heterocyclic chemist *par excellence*, in celebration of his 75th birthday.



Herein we report our preliminary results of a sequel study concerning the use of *o*-silylpyridyl triflates **5** for the generation of the corresponding 3,4-pyridynes (**6**) and their cycloaddition (**7**) and nucleophilic (**8**) reactions respectively.¹²

Adopting the earlier general methodology for the synthesis of substituted pyridines *via* the powerful carbamate directed metalation group,¹³ the 3-pyridyl carbamate (**9**) was sequentially lithiated with LDA and quenched with TMSCl or TESCl (triethylsilyl chloride) to furnish regiospecifically the 4-silylated products (**10a**) and (**10b**) in good to excellent yields. Since silylpyridines are sensitive to base-induced cleavage,¹⁴ carbamate hydrolysis of **10a-b** under basic conditions¹⁵ was precluded. Although reductive carbamate cleavage of **10b** was achieved using LAH, the resulting 3-hydroxy-4-trimethylsilylpyridine was unstable to desilylation under workup conditions. The TES derivative (**10a**), on the other hand, did not suffer from this problem and the analogous pyridinol was readily obtained and converted directly into the triflate (**11a**). The corresponding picolinamide (**11b**) was obtained by taking advantage of the previously established anionic ortho-Fries rearrangement.¹³ Thus LDA metalation of **10b** at -78°C and slow warming to room temperature afforded the migration product (**12**)¹⁶ which was converted into the triflate (**11b**).


Scheme 2



Treatment of **11a** with an excess of furan in the presence of TBAF in acetonitrile at ice bath temperatures led to the cycloaddition product (**13a**) in modest yield (54%). Under CsF conditions in the presence of 18-crown-6, the cycloaddition of **11a** with 1,3-diphenylisobenzofuran to give adduct (**14a**) proceeded in a similar yield (58%). In contrast to these reactions with electron-rich dienes, the analogous transformation of **11a** with tetracyclone, an electron-deficient diene, led to the isoquinoline (**15**) in low yield (26%). Parallel observations were made in the reactions with the picolinamide 3,4-pyridyne precursor (**11b**) to give adducts (**13b**) (44%) and (**14b**) (17%) in yields which are lower presumably owing to steric effects from the carboxamide substituent.

Nucleophilic reactivity of **11a** and **11b** was probed with phenol and thiophenol. While phenol was ineffective in these reactions (<15% yields of products), thiophenol afforded good yields of substituted products (**16a-b**) and (**17a-b**). In consonance with previous observations,^{6,17} a 1:1 mixture of **16a** (42%) and **17a** (42%) was obtained from the reaction of **11a** with thiophenol. Also as expected,¹⁸ the analogous picolinamide (**11b**) underwent regioselective addition to give **16b** in 55% yield. The isomer (**17b**) was not detected (<2%).

Scheme 3



In summary, silylpyridyl triflates (**11a-b**), prepared by taking advantage of 3-pyridyl carbamate directed ortho metalation chemistry, have been shown to serve as useful precursors for 3,4-pyridynes (**6a-b**) under neutral fluoride-induced conditions from which result synthetically useful yields of cycloaddition (**13a-b**, **14a-b**, **15**) and nucleophilic addition (**16a-b**, **17a-b**) products. The use of 4-trialkylsilyl-3-pyridyl triflates for 3,4-pyridyne generation has advantage over 4-trimethylsilyl-3-bromopyridine¹² in that using the latter intermediate a) yields of cycloaddition products (e.g. **13a**) are lower and b) its synthesis is complicated by elimination and halogen dance reactions. Furthermore, the connection to aromatic metalation chemistry for the generation of a variety of other substituted 3,4-pyridyne precursors may be feasible by the described methodology.¹⁹

ACKNOWLEDGEMENT

We are grateful to NSERC Canada and the University of Waterloo for the support of this research.

REFERENCES AND NOTES

1. M. G. Reinecke, *Tetrahedron*, **1982**, *38*, 427.
2. H. H. Nam and G. L. Leroi, *J. Am. Chem. Soc.*, **1988**, *110*, 4096 and refs cited therein.
3. a) G. W. Gribble and M. G. Saulnier, *Tetrahedron Lett.*, **1980**, *21*, 4137; b) F. Marsais, B. Laperdrix, T. Güngör, M. Mallet, and G. Quéguiner, *J. Chem. Res. (M)*, **1982**, 2863; c) F. Marsais and G. Quéguiner, *Tetrahedron*, **1983**, *39*, 2009; d) S. P. Modi and S. Archer, *J. Org. Chem.*, **1989**, *54*, 5189; e) B. Jamart-Gregoire, C. Leger, and P. Caubere, *Tetrahedron Lett.*, **1990**, *31*, 7599; f) See also N. Furukawa, T. Shibutani, and H. Fujihara, *Tetrahedron Lett.* **1987**, *28*, 2727.
4. T. Kauffmann and F.-P. Boettcher, *Chem. Ber.*, **1962**, *95*, 949.
5. T. Sasaki, K. Kanematsu, and M. Uchide, *Bull. Chem. Soc. Japan*, **1971**, *44*, 858.
6. T. Kauffmann, *Angew. Chem. Int. Ed. Engl.*, **1965**, *4*, 543.
7. The theoretically predicted less stable 2,3-pyridyne (M. J. S. Dewar and G. P. Ford, *J. Chem. Soc., Chem. Commun.*, **1977**, 539), although generated *via* aminotriazole and anhydride intermediates (G. W. Fleet and I. Fleming, *J. Chem. Soc. C*, **1969**, 1758; E. K. Fields and S. Meyerson, *J. Org. Chem.*, **1966**, *31*, 3307), has enjoyed little synthetic impact.
8. Y. Himeshima, T. Sonoda, and H. Kobayashi, *Chemistry Lett.*, **1983**, 1211; see also the original report by R. F. Cunico and E. M. Dexheimer, *J. Organometal. Chem.*, **1973**, *59*, 153.
9. For use of *o*-haloaryl tosylates and aryl triflates themselves for benzyne formation, see W. H. Darlington and J. Szmuszkowicz, *Tetrahedron Lett.*, **1988**, *29*, 1883; K.-y. Jung and M. Koreeda, *J. Org. Chem.*, **1989**, *54*, 5667; P. P. Wickham, K. H. Hazen, H. Guo, G. Jones, K. H. Reuter, and W. J. Scott, *J. Org. Chem.*, **1991**, *56*, 2045; T. Matsumoto, T. Hosoya, M. Katsuki, and K. Suzuki, *Tetrahedron Lett.*, **1991**, *32*, 6735.
10. a) K. Shankaran and V. Snieckus, *Tetrahedron Lett.*, **1984**, *25*, 2827; b) E. G. Doadt, Ph. D. Thesis, University of Waterloo, **1988**.
11. V. Snieckus, *Chem. Rev.*, **1990**, *90*, 879.
12. Complementary work on *o*-silylbromopyridines but under *t*-BuOK induction has been recently reported: E. Effenberger and W. Daub, *Chem. Ber.*, **1991**, *124*, 2119.
13. M. A. J. Miah and V. Snieckus, *J. Org. Chem.*, **1985**, *50*, 5436.
14. A. Fischer, M. W. Morgan, and C. Eaborn, *J. Organometal. Chem.*, **1977**, *136*, 323.
15. M. P. Sibi and V. Snieckus, *J. Org. Chem.*, **1983**, *48*, 1935.
16. 2-Deprotonation of **10b** has been also achieved, a result which allows a general protocol for the synthesis of 2-substituted 3-pyridinols *via* 4-silyl protection: M. Tsukazaki and V. Snieckus, in preparation.
17. a) J. A. Zoltewicz and C. Nisi, *J. Org. Chem.*, **1969**, *34*, 765; b) W. J. van Zoest and H. J. den Hertog, *Rec. Trav. Chim. Pays-Bas*, **1974**, *93*, 166.
18. R. W. Hoffman, "Dehydrobenzene and Cycloalkynes," Academic Press, New York, **1967**, p. 99; E. K. Fields, "Organic Reactive Intermediates," ed. S. P. McManus, Academic Press, New York, **1973**, p. 449.
19. All new compounds show analytical and spectral (ir, nmr, ms, hrms) data in agreement with the assigned structures.

Received, 28th November, 1991