

STUDY OF THIAZOLO[4,5-*d*]PYRIMIDINES : THE SYNTHESIS
OF THIAZOLO[4,5-*d*]PYRIMIDINE-2,7-DIONES AND NOVEL
RING OPENING TO 2,4-THIAZOLIDINEDIONE

Katsuhiko Nagahara* , Masae Sekine, and Atsushi Takada

School of Pharmaceutical Sciences, Kitasato University,
5-9-1, Shirokane, Minato-ku, Tokyo 108, Japan

Howard B. Cottam and Roland K. Robins


Department of Medicine 0663, University of California,
San Diego, 9500 Gilman Drive, La Jolla, California 92093,
U. S. A.

Abstract --- Treatment of 7-oxo-3-phenylthiazolo[4,5-*d*]-pyrimidine-2(6*H*)-thione (**1**) with dimethyl sulfate afforded 6-methyl-3-phenylthiazolo[4,5-*d*]pyrimidine-2,7-dione (**2**), 3-phenylthiazolo[4,5-*d*]pyrimidine-2,7(6*H*)-dione (**3**) and/or 5-*N*-methylcarbamoyl-3-phenyl-2,4-thiazolidinedione (**5**), depending on reaction conditions. Furthermore, reaction of **3** with dimethyl sulfate caused the ring opening to give corresponding **5**. Also, treatment of **5** with phosphorus oxychloride gave 4-chloro-2-oxo-3-phenylthiazolidine-5-*N*-methylcarboxamide (**6**).

We have recently described the synthesis and immunopotentiating properties

of purine nucleoside analogues in the thiazolo[4,5-*d*]pyrimidine ring system.^{1,2} We now report the reaction of dimethyl sulfate with a substituted thiazolo[4,5-*d*]pyrimidine-2-thione. Gewald reported that the reaction of 7-oxo-3-phenylthiazolo[4,5-*d*]pyrimidine-2(6*H*)-thione (**1**) with dimethyl sulfate gave only 6-methyl-3-phenylthiazolo[4,5-*d*]pyrimidine-2,7-dione (**2**).³ However, we found that the treatment of **1** with dimethyl sulfate under the same reaction conditions (at 130°C) afforded **2** and 3-phenylthiazolo[4,5-*d*]pyrimidine-2,7(6*H*)-dione (**3**).⁴ Also, heating of **1** with dimethyl sulfate at 110-120°C or 150-160°C gave only **3** or the corresponding **2** and a small amount of a new 2,4-thiazolidinedione (**5**), respectively. Furthermore, the reaction of **3** with aqueous dimethyl sulfate at 150-160°C caused the novel ring cleavage of thiazolo[4,5-*d*]pyrimidine-2,7-dione to give **5**.

Treatment of **1** (1.9 mmol) with 10-fold molar amount of dimethyl sulfate under heat for 1 h at 130°C afforded the corresponding **2** (25%) and **3** (27%), respectively. Next, stirring of **1** with an equivalent amount of dimethyl sulfate in an aqueous solution of 1N NaOH at room temperature for 20 min gave a




Scheme 1

75% yield of 6-methyl-2-thioxo-3-phenylthiazolo[4,5-*d*]pyrimidin-7-one (**4**).⁵

The 6-methyl-2,7-dione type compound (**2**) was identical to the product of the reaction of **4** with dimethyl sulfate under heating for 1 h at 130°C (Scheme 1).


Interestingly, the reaction of **1** with dimethyl sulfate under heating at 110-120°C gave only **3** in high yield (93%). Also, heating of **1** with dimethyl sulfate at 150-160°C afforded **2** (43%) and 5-*N*-methylcarbamoyl-3-phenyl-2,4-thiazolidinedione (**5**) (8%),⁶ respectively. Moreover, the reaction of **3** with dimethyl sulfate at 150-160°C for 1 h caused the ring opening to **5** (56%). Heating of **2** with dimethyl sulfate at 150-160°C, however, under the similar reaction conditions resulted in **2** being recovered unchanged.

Next, treatment of **5** (3.3 mmol) with 10-fold molar amount of phosphorus oxychloride and an equivalent amount of *N,N*-dimethylaniline under reflux at 130°C for 3 h afforded 4-chloro-2-oxo-3-phenylthiazolidine-5-*N*-methylcarbamamide (**6**) (49%)⁷ (Scheme 2).



Scheme 2

The novel ring cleavage of **1** to **5** probably proceeds through initial hydrolysis of the thione moiety on the compound (**1**), followed by the ring opening of the pyrimidine ring and methylation, and loss of *N,N*-dimethylformamide by hydrolysis (Scheme 3).



Scheme 3

REFERENCES AND NOTES

1. K. Nagahara, J. D. Anderson, G. D. Kini, N. K. Dalley, S. B. Larson, D. F. Smee, A. Jin, B. S. Sharma, W. B. Jolley, R. K. Robins, and H. B. Cottam, *J. Med. Chem.*, 1990, **33**, 407.
2. G. D. Kini, J. D. Anderson, Y. S. Sanghvi, A. F. Lewis, D. F. Smee, G. R. Revankar, R. K. Robins, and H. B. Cottam, *ibid.*, 1991, **34**, 3006.
3. K. Gewald, *J. Prakt. Chem.*, 1966, **32**, 26.
4. **3** : mp > 300°C; EI-ms : m/z 245 (M^+); ir : ν cm^{-1} 3420 (NH), 1700,

1640 (C=O); ^1H -nmr (DMSO- d_6) δ : 8.16 (1H, s, C-H), 13.01 (1H, s, N-H, exchangeable with D_2O).

5. **4** : mp 250-251°C; EI-ms : m/z 275 (M^+); ir : ν cm^{-1} 1670 (C=O).
6. **5** : mp 218-219°C; EI-ms : m/z 250 (M^+); ir : ν cm^{-1} 3320 (NH), 1740, 1700, 1670 (C=O); ^1H -nmr (DMSO- d_6) δ : 2.69 (3H, d, J = 4.5 Hz, Me), 5.24 (1H, s, C-H, exchangeable with D_2O), 7.26-7.56 (5H, m, Ph), 8.58 (1H, d, J = 4.5 Hz, NH, exchangeable with D_2O); ^{13}C -nmr (DMSO- d_6) δ : 26.4 (Me), 52.2 (C-5, exchangeable with D_2O), 127.8, 129.2, 129.3, 133.1 (Ph), 164.6 (-NHCO-), 169.6 (C-2), 170.9 (C-4).
7. **6** : mp 178-180°C; EI-ms : m/z 268 (M^+), 270 ($\text{M} + 2$); ir : ν cm^{-1} 3420, 3400 (NH), 1690, 1660, 1630 (C=O); ^1H -nmr (DMSO- d_6) δ : 2.76 (3H, d, J =4.0 Hz, Me), 7.24-7.57 (5H, m, Ph), 8.03 (1H, d, J = 4.5 Hz, NH, exchangeable with D_2O); ^{13}C -nmr (DMSO- d_6) δ : 26.6 (Me), 108.1 (C-5), 120.8 (C-4), 129.0, 129.1, 129.9, 134.0 (Ph), 158.9 (-NHCO-), 166.7 (C-2).

The compounds (3, 4, 5, 6) were confirmed by the elemental analyses.

Received, 7th September, 1992