

**INVERSE [4+1] CYCLOADDITIONS OF 3-METHYL-2,3-DIHYDRO-
1,3-BENZOTHAZOLE-2-YLIDENE WITH 1,2,4,5-TETRAZINES**

Peter Imming, Andreas Kümmell, and Gunther Seitz* ¹


Institut für Pharmazeutische Chemie der Philipps-Universität, Marbacher Weg 6
3550 Marburg/Lahn, Germany

Abstract - [4+1] Cycloadditions of the nucleophilic singlet carbene 3-methyl-2,3-dihydro-1,3-benzothiazole-2-ylidene (**2**) with the electron-deficient s-cis-fixed diazadiene systems of a series of substituted 1,2,4,5-tetrazines (**4a-c**), with CF₃, SCH₃ and C₆H₅-groups are described together with some unexpected follow-up reactions.

In the course of our work¹⁻⁴ on cheletropic [4+1] cycloadditions of nucleophilic singlet carbenes⁵⁻⁹ to cyclic diazadiene systems we have recently included 3-methyl-2,3-dihydro-1,3-benzothiazole-2-ylidene (**2**)¹⁰⁻¹⁴ in our studies. The present paper reports successful transformations of this heterocyclic amino-thio-carbene with the differently substituted 1,2,4,5-tetrazines (**4a-c**).


We used the cyano-substituted benzothiazoline (**1a**)¹⁵ as a safe carbene source. **1a** is readily accessible from 3-methyl-benzothiazolium iodide and potassium cyanide, and easy to handle. On adding a saturated solution of **1a** to **4a** in boiling benzene, decolourization of the red solution occurs rapidly. Work-up by column chromatography yields an orange substance whose analytical and spectroscopic data are in accord with constitution (**7a**). The formation of **7a** seems plausible if one assumes the [4+1] cycloadduct (**5a**) and the strained spiro compounds (**6a**) (formed by elimination of elemental nitrogen) to be non-isolable intermediates. **6a** finally stabilizes itself by a [1,5] sigmatropic rearrangement of the thiazole sulfur atom, giving **7a**. The reaction of **1** with **4b** is interpreted analogously to the stage of the tricyclic heterocycle (**7b**). After that, the pronounced tendency of the angular methylthio group to migrate occasions another sigmatropic shift, yielding the sulfenic acid amide (**8**).

¹Dedicated to Prof. E. C. Taylor on the occasion of his 70th birthday.



8, in turn, is probably hydrolyzed during chromatography on silica gel and transformed to the stable pyrazole derivative (9b), isolated as a beige solid in 44% yield. The reaction of 1a with the phenyl-substituted tetrazine (4c) took a surprising turn. The main product was the phenyl thiocyanate (10), isolated as colourless crystals in 73% yield. In addition, small amounts of the disulfide (11) were isolated as a pale yellow crystalline powder. The structures of both compounds agree with their analytical and spectroscopic data.¹⁶ The unusual formation of the phenyl thiocyanate (10) is explainable if one assumes 6c to form intermediately. 6c then reacts with the hydrogen cyanide liberated from 1a. The cyanide ion presumably attacks the bivalent sulfur atom occasioning ring opening as indicated by the arrows, and subsequent protonation of the nitrogen atom of the pyrazole ring. The formation of the disulfide (11) is complex and cannot be explained without having isolated intermediate products. Interestingly, 11

was the only product isolated (yield, 78%) from the reaction of **1b** with **4c**, taking no account of the piperidine eliminated from **1b**.



We thank the Deutsche Forschungsgemeinschaft and the Fonds der Chemischen Industrie for financial support; the Hoechst AG, Bayer AG, and Solvay, Fluor and Derivate GmbH for generous gifts of chemicals.

EXPERIMENTAL PART

Hplc was done with a Merck-Hitachi Liquid Chromatograph. Melting points are uncorrected. Ir spectra were recorded with Perkin-Elmer spectrophotometers 257 and 398, nmr spectra with Varian T60 and XL100 and Jeol JNM-FX100 and -GX400, mass spectra with Vacuum Generators 7070 (70 eV) spectrometers.

4-Methyl-3,9a-bis(trifluoromethyl)-9aH-pyrazolo[3,4-b][1,4]benzothiazine (7): A solution of **4a** (0.65 g, 3 mmol) in 10 ml of dry benzene is heated to reflux while a saturated solution of **1a** (0.53 g, 3 mmol) is added until the red colour of the tetrazine is faded (after about two thirds of the stoichiometric amount). The solvent is evaporated in vacuo, and the residue was chromatographed on silica gel (column, 30 x 3 cm; eluent: n-hexane/ethyl

acetate 8+2). **7** is eluted as an orange-yellow fraction and crystallized from n-hexane, yielding 0.24 g (31%, related to **1**), mp 113°C. Ir (KBr): $\nu = 3105 \text{ cm}^{-1}$, 1600, 1580, 1570, 1480, 1440, 1425, 1400, 1215, 1195, 1175, 1125, 1110, 760. Uv (CH_2Cl_2): λ_{max} (lg ϵ) = 261 nm (3.701), 280 (3.783), 285 (3.832), 341 (3.581), 416 (3.809). $^1\text{H-Nmr}$ (400 MHz, CDCl_3): $\delta = 7.46\text{--}7.23$ (4 H, m, phenyl H), 3.73 (3 H, q, CH_3 , $J_{\text{HF}} = 2.0 \text{ Hz}$). $^{13}\text{C-Nmr}$ (CDCl_3): $\delta = 147.3$ (s, broad, C-3a, $J_{\text{CF}} = 1 \text{ Hz}$), 137.9 (dd; C-4a, $^3J_{\text{CH}} = 7/5 \text{ Hz}$), 128.0/127.9 (2d; 2 phenyl CH, $^1J_{\text{CH}} = 168/170 \text{ Hz}$), 127.4 (q, C-3, $^2J_{\text{CF}} = 38 \text{ Hz}$), 126.2 (d, phenyl CH, $^1J_{\text{CH}} = 172 \text{ Hz}$), 122.1 (q, CF_3 , $^1J_{\text{CF}} = 287 \text{ Hz}$), 121.5 (q, CF_3 , $^1J_{\text{CF}} = 269 \text{ Hz}$), 116.1 (d, C-5, $^1J_{\text{CH}} = 160 \text{ Hz}$), 111.4 (dd, C-8a, $^3J_{\text{CH}} = 10/7 \text{ Hz}$), 83.8 (q, C-9a, $^2J_{\text{CF}} = 30 \text{ Hz}$), 40.0 (qq, CH_3 , $^1J_{\text{CH}} = 142 \text{ Hz}$, $J_{\text{CF}} = 4 \text{ Hz}$). Ms (70 eV, 30°C): m/z (%) = 339 (100) [M^+], 341 (14). Anal. Calcd for $\text{C}_{12}\text{H}_7\text{N}_3\text{F}_6\text{S}$: C, 42.48; H, 2.08; N, 12.39. Found: C, 42.74; H, 2.11; N, 12.40.

4-Methyl-3-methylthio-1H-pyrazolo[4,5-*b*][1,4]benzothiazine (9): A solution of **1b** (0.47 g, 2 mmol) and **4b** (0.70 g, 2 mmol) in 5 ml of dry benzene is heated to reflux under nitrogen for 30 min. The solvent is removed in vacuo, and the residue was chromatographed on silica gel (column, 40 x 3 cm). n-Hexane/dichloromethane (1+1) elutes **1b**, ethyl acetate impure **9**, which after a second chromatographic purification (column, 30 x 3 cm; eluent: n-hexane/ethyl acetate 6 + 4) is isolated as a yellow oil that solidifies on drying in a vacuum. After several washings with a mixture of n-pentane and ether (9 + 1) a beige solid is obtained. Yield, 0.22 g (44%), mp 49–52°C. Ir (KBr): $\nu = 3140 \text{ cm}^{-1}$, 2960, 1555, 1475, 1410, 1310, 1285, 755. $^1\text{H-Nmr}$ (400 MHz, DMSO-d_6): $\delta = 12.81$ (s, broad, 1 H, NH), 7.12 (1H, m), 7.04 (1H, m), 6.85–6.79 (2H, m, 4 phenyl H), 3.43 (3 H, s, NCH_3), 2.38 (3 H, s, SCH_3). $^{13}\text{C-Nmr}$ (CDCl_3): $\delta = 144.7$ (s, C-4a), 135.3 (s*) / 129.9 (s, 2 phenyl C), 127.8/127.4 (2d, C-6/8, $^1J_{\text{CH}} = 160 \text{ Hz}$ each), 121.2 (d, C-7, $^1J_{\text{CH}} = 162 \text{ Hz}$), 120.1 (s, C-8a), 118.8 (s*, phenyl C), 113.6 (d, C-5, $^1J_{\text{CH}} = 158 \text{ Hz}$), 35.4 (q, NCH_3 , $^1J_{\text{CH}} = 138 \text{ Hz}$), 21.6 (q, SCH_3 , $^1J_{\text{CH}} = 141 \text{ Hz}$). *Signals show marked line broadening due to tautomeric equilibrium of the pyrazole system. Ms (70 eV, 130°C): m/z (%) = 249 (100) [M^+], 234 (54) [$\text{M}^+ - \text{CH}_3$]. Anal. Calcd for $\text{C}_{11}\text{H}_{11}\text{N}_3\text{S}_3$: C, 52.99; H, 4.45; N, 16.85; S, 25.72. Found: C, 52.96; H, 4.50; N, 16.44; S, 26.00.

Reaction of 1a and 4c: A solution of **4c** (0.47 g, 2 mmol) in 6 ml of dry toluene is heated to reflux under nitrogen. **1a** (0.35 g, 2 mmol), dissolved in toluene (10 ml), is added dropwisely. After refluxing for 1 h, the solvent is evaporated and the residue was chromatographed on silica gel (column, 30 x 3 cm). With dichloromethane

unreacted tetrazine was eluted. A mixture of n-hexane and ethyl acetate (1:1) eluted the phenyl isothiocyanate (**10**) (pale yellow solid), pure ethyl acetate the disulfide (**11**) (pale yellow powder).

Methyl(3,5-diphenyl-1H-pyrazol-4-yl)-(2-thiocyanatophenyl)amine (10): Yield 0.53 g (73%), colourless crystals, mp 182°C (acetone). Ir (KBr): $\nu = 3310 \text{ cm}^{-1}$ (NH), 2165 (SCN), 1585, 1480, 1455, 1445, 960. Uv (CH_2Cl_2): λ_{max} (lg ϵ) = 231 nm (4.471), 281 (sh, 4.012). $^1\text{H-Nmr}$ (400 MHz, DMSO-d_6): $\delta = 13.42$ (1 H, s, broad, NH), 7.77 (2H, s, broad, phenyl H), 7.49 (2H, s, broad, phenyl H), 7.36-7.07 (9H, m, phenyl H), 6.74 (1H, m, 4'-H), 3.22 (3 H, s, CH_3). $^{13}\text{C-Nmr}$ * (DMSO-d_6): $\delta = 147.9, 146.0, 137.6, 132.5, 131.3, 129.6, 128.3, 127.7, 126.4, 124.2, 120.7, 117.6, 111.9, 111.2, 41.7$. Ms (70 eV, 60°C): m/z (%) = 382 (13) [M^+], 355 (33), 354 (29), 252 (68), 149 (100). *The phenyl substituents are recorded isochronically because of the tautomeric equilibrium of the pyrazole system. Anal. Calcd for $\text{C}_{23}\text{H}_{18}\text{N}_4\text{S}$: C, 72.23; H, 4.74; N, 14.65; S, 8.38. Found: C, 71.96; H, 4.83; N, 14.44; S, 8.04.

Bis{2-[(3,5-Diphenyl-1H-pyrazol-4-yl)methylamino]}diphenyl disulfide (11): Yield 174 mg (24%) pale yellow crystals, mp 257°C [decomp. n-hexane/ethyl acetate 1+1]. Ir (KBr): $\nu = 3390 \text{ cm}^{-1}$ (broad, NH), 3190, 1585, 1470, 1450, 1330, 1270, 755. $^1\text{H-Nmr}$ (400 MHz, DMSO-d_6)*: $\delta = 13.18$ (1H, s, broad, NH), 7.80 (2H, s, broad, phenyl H), 7.46 (2H, s, broad, phenyl H), 7.30-7.20 (6H, m, phenyl H), 6.99 (1H, m, 4'-H), 6.89 (1H, m, 3'-H), 6.63 (1H, m, 6'-H), 6.47 (1H, m, 5'-H), 3.14 (3 H, s, CH_3). *Relative integrals; multiply by two to get absolute number of protons. $^{13}\text{C-Nmr}$ (DMSO-d_6): $\delta = 146.5, 146.2, 137.7, 133.0, 128.2, 127.4, 126.7, 126.5, 126.0, 125.5, 123.3, 119.6, 116.2, 41.8$. The phenyl substituents are recorded isochronically because of the tautomeric equilibrium of the pyrazole system, but show considerable line broadening. Ms (70 eV, 250°C): m/z (%) = 357 (100) [$1/2 \text{ M}^+ + \text{H}$]. Anal. Calcd for $\text{C}_{44}\text{H}_{36}\text{N}_6\text{S}_2$: C, 74.13; H, 5.09; N, 11.79; S, 9.00. Found: C, 73.50; H, 5.05; N, 11.63; S, 9.04.

Reaction of 3-methyl-2-piperidinobenzothiazoline (1b) with 4c: A solution of **1b** (0.56 g, 2.4 mmol) and **4c** (0.47 g, 2.0 in 10 ml of dry xylene is heated to reflux for 1 h. The residue after removal of the solvent in vacuo is separated chromatographically on silica gel (column, 40 x 3 cm). The fractions eluted with dichloromethane contain residual xylene, **4c**, and piperidine and are discarded. Ethyl acetate elutes **11** that is purified by crystallization as described above. Yield, 556 mg (78%).

REFERENCES

1. A. Kümmell and G. Seitz, *Tetrahedron Lett.*, 1991, 32, 2743.
2. P. Imming, R. Mohr, E. Müller, W. Overheu, and G. Seitz, *Angew. Chem.*, 1982, 94, 291; *Angew. Chem., Int. Ed. Engl.*, 1982, 21, 284.
3. G. Seitz, S. Dietrich, R. Dhar, W. Massa, and G. Baum, *Arch. Pharm. (Weinheim)*, 1986, 319, 798.
4. X.-G. Yang, R. John, and G. Seitz, *Arch. Pharm. (Weinheim)*, 1991, 324, 923.
5. R.W. Hoffmann, K. Steinbach, and W. Lilienblum, *Chem. Ber.*, 1976, 109, 1759.
6. W. Lilienblum and R.W. Hoffmann, *Chem. Ber.*, 1977, 110, 3405.
7. K. Burger, U. Wassmuth, and S. Penninger, *J. Fluorine Chem.*, 1982, 20, 813.
8. H. Möhrle and H. Dwuletzi, *Chemiker-Ztg.*, 1987, 111, 9.
9. Reviews: Methoden Org. Chem. (Houben-Weyl) 4th ed., Vol. E19b, part 1, p. 1 and part 2, p. 1628, 1746; R.A. Moss, *Acc. Chem. Res.*, 1980, 13, 58ff; C. Wentrup, *Reactive Molecules*, Wiley, New York 1984, p. 237f.
10. J.-J. Vorsanger, *Bull. Soc. Chim. Fr.*, 1964, 119.
11. J. Metzger, H. Larive, R. Dennilaule, R. Baralle, and C. Gaurat, *Bull. Soc. Chim. Fr.*, 1964, 2857.
12. H. Quast and S. Hünig, *Angew. Chem.*, 1964, 76, 989; *Angew. Chem., Int. Ed. Engl.*, 1964, 3, 800.
13. H. Wanzlick, H.-J. Kleiner, I. Lasch, and H.U. Földner, *Angew. Chem.*, 1966, 78, 115; *Angew. Chem., Int. Ed. Engl.*, 1966, 5, 126 and references quoted there.
14. R.W. Hoffmann, B. Hagenbruch, and D.M. Smith, *Chem. Ber.*, 1977, 110, 23.
15. H. Wahl and J.-J. Vorsanger, *Bull. Soc. Chim. Fr.*, 1965, 3359.
16. For the nmr spectroscopical structure elucidation with the aid of COLOC and CH-COSY spectra see A. Kümmell, Dissertation, Universität Marburg (Germany), 1990.

Received, 19th October, 1992