

REGIOSELECTIVITY IN THE 1,3-DIPOLAR CYCLOADDITION REACTION
OF DIAZOMETHANE TO 7-OXANORBORNENE DERIVATIVES

Odón Arjona*, Cristina Manzano, and Joaquín Plumet*


Departamento de Química Orgánica I, Facultad de Química, Universidad
Complutense, 28040 Madrid, Spain

Dedicated to Prof. Edward C. Taylor

Abstract—The regioselectivity of the 1,3-dipolar cycloaddition reaction of diazomethane to 7-oxabicyclo[2.2.1]hept-5-ene derivatives substituted on C-2 is discussed. The observed ratio of regioisomers depends upon the nature of the substituents on C-2 and may be accounted for in terms of TS asynchrony for concerted processes.

The regioselectivity of electrophilic additions to bicyclo[2.2.1]hept-5-ene systems is controlled by the nature of the substituents at C-2.¹ On the other hand, Diels-Alder² and 1,3-dipolar cycloaddition reactions of nitrile betaines³ to 7-oxabicyclo[2.2.1]hept-5-ene derivatives⁴ displayed only moderate regioselectivity. In these last cases kinetic parameters indicate a concerted mechanism. However, the cycloadditions of diazonium betaines with oxanorbornenic compounds are unprecedented in the literature.⁵ In this report we wish account for our finding in this field.

Diazomethane reacts with oxanorbornenic derivatives (1a) and (1b) affording adducts (2) and (3) in quantitative yield.⁶ Isomeric ratios were determined by integration of the 360 MHz ¹H-nmr spectra of the crude reaction mixtures. No isomer from *endo* face attack to the carbon-carbon double bond could be detected.⁷



All attempts to carry out the cycloaddition of diazomethane with the related 7-oxabicyclo[2.2.1]hept-5-ene-2-one led to inseparable mixtures formed by products of homology and/or cycloaddition. It has been reported that the reaction between diazomethane and norbornenic ketones leads to cycloaddition products when a mixture of MeOH-Et₂O is used as a solvent, at -25°C,⁸ and Adam *et al.* have prepared these adducts in CCl₄ at 20°C.⁹ Homologation procedures of the saturated 7-oxanorbornanone employing diazomethane have been described.¹⁰ In our case, all efforts to obtain exclusively cycloadditions products in the conditions reported for norbornenic ketones were fruitless. Nevertheless, products formally arising from cycloaddition of diazomethane to the double bond of 7-oxabicyclo[2.2.1]hept-5-ene-2-one may be obtained by treatment of derivatives (2a) and (3a) with MeONa/MeOH-formalin.¹¹

Adducts (2) and (3) were separated by fractional crystallization (EtOH) of the reaction crude mixture and their structures were determined by NOE experiments in their 360 MHz ¹H-nmr spectra. All signals for both regioisomers were assigned in this fashion.¹² Ratio of regioisomers was evaluated by integration of the signals corresponding to the H-3 *endo* protons.

Table 1

FMO's energies and coefficients for 1a and 1b


Compound	$\epsilon(\text{HOMO})$	$\epsilon(\text{LUMO})$	C_5HO	C_6HO	C_5LU	C_6LU
<u>1a</u>	-7.97eV	7.80eV	0.564	0.561	-0.749	0.716
<u>1b</u>	-7.37eV	8.09eV	0.410	0.383	0.775	-0.766
			CCH_2HO	$\text{C}_\text{N}\text{HO}$	CCH_2LU	$\text{C}_\text{N}\text{LU}$
<u>Diazomethane</u>	-6.75eV	7.86eV	-0.74	-0.659	0.662	0.612

The PMO theory provides a qualitative answer to regioselectivity problems in 1,3-dipolar cycloadditions.¹³ Using MNDO¹⁴ optimised geometries we have calculated the *ab initio* STO-3G¹⁵ MO's of 2-*endo*-hydroxy-7-oxabicyclo[2.2.1]hept-5-ene-2-carbonitrile (1, X=CN, Y=OH a reasonable model for 1a), (1b) and

diazomethane¹⁶ (Table 1).

The observed regioselectivity is qualitatively in agreement with a HOMO-dipole controlled reaction.¹⁷ However, the differences in regioselectivity founded between (1a) and (1b) are not explained by this simple PMO's approach. Inspection of the energies and coefficients for both HO-LU interactions indicates that the regioselectivity should be very similar for both compounds (1a) and (1b).

Diazomethane is a good example of ambivalence of 1,3-dipoles,¹⁸ possessing the nucleophilic terminus on the carbon atom and the electrophilic site on the terminal nitrogen. In terms of asynchronicity of TS of 1,3-dipolar cycloaddition reactions, significant contribution of structure (4) is then possible.



MNDO calculations¹⁴ of model cations and anions in the 7-oxanorbornene system (Table 2) indicate that a homoconjugated ethylenedioxy functionality stabilizes both positive and negative charges developing on C-6 (structures 5 and 6).

Table 2

MNDO calculations of heats of formation of model cations and anions (Kcal/mol)

		<u>Cation (ΔH_0°)</u>		<u>Anion (ΔH_0°)</u>	
<u>X</u>	<u>Y</u>	<u>C₅</u>	<u>C₆</u>	<u>C₅</u>	<u>C₆</u>
F	CN	204.7	204.6	-26.6	-32.3
Cl	CN	238.7	239.3	6.74	1.21
OCH ₂ CH ₂ O		115.6	111.9	-89.8	-92.4

The contributions of these structures to the nature of the transition state do not modify the regioselectivity predicted by application of the PMO theory. On the other hand, electron deficient groups (F, Cl, CN) on C-2

stabilize a negative charge on C-6.

Thus, an increased contribution of structure (5) to the description of TS in the case of (1a) could be responsible for the difference in regioselectivity observed. In conclusion, the reaction of diazomethane and 7-oxanorbornenic derivatives is, at least in part, controlled by remote substitution on C-2 in the bicyclic system.

ACKNOWLEDGEMENTS

We thank Dr. Carrupt (Lausanne) for the MNDO calculations of heats of formation (Table 2). We acknowledge support of this research by the CICYT (Grant nº 9035, Ministerio de Educación y Ciencia, Spain) and by PharmaMar S.A. One of us (C. M.) acknowledge the Ministerio de Educación y Ciencia of Spain for a fellowship.

REFERENCES AND NOTES

1. See: O. Arjona, A. Mallo, L. García, and J. Plumet, *J. Chem. Soc., Perkin Trans. II*, 1989, 1315; O. Arjona, R Fernández de la Pradilla, J. Plumet, and A. Viso, *Tetrahedron*, 1989, **45**, 4565; O. Arjona, R Fernández de la Pradilla, I. Pita-Romero, J. Plumet, and A. Viso, *Tetrahedron*, 1990, **46**, 8199; O. Arjona, R Fernández de la Pradilla, J. Plumet, and A. Viso, *J. Org. Chem.*, 1991, **56**, 6227 and references quoted in the paper.
2. K. A. Black and P. Vogel, *J. Org. Chem.* 1986, **51**, 5341.
3. J. Plumet, G. Escobar, C. Manzano, O. Arjona, P. A. Carrupt, and P. Vogel, *Heterocycles*, 1986, **24**, 1535; O. Arjona, C. Dominguez, R Fernández de la Pradilla, A. Mallo, C. Manzano, and J. Plumet, *J. Org. Chem.*, 1989, **54**, 5883; O. Arjona, A. de Dios, R Fernández de la Pradilla, A. Mallo, and J. Plumet, *Tetrahedron*, 1990, **46**, 8179.
4. For other 1,3-dipolar cycloaddition reactions to 7-oxanorbornene systems, see: D. Cristina, M. de Amici, C. de Micheli, and R. Gandolfi, *Tetrahedron*, 1981, **37**, 1349; L. Fisera, J. Kovác, J. Patús, and D. Pavlovic, *Collect. Czech. Chem. Comm.*, 1983, **48**, 1048; L. Fisera and D. Pavlovic, *Collect. Czech. Chem. Comm.*, 1984, **49**, 1990; O. Arjona, R. F. de la Pradilla, R. A. Pérez, and J. Plumet, *Tetrahedron*, 1988, **44**, 7199.
5. For a review of 1,3-dipolar cycloaddition reactions of diazoalkanes, see: M. Regitz and H. Heydt, '1,3-Dipolar Cycloadditions Chemistry,' Vol. 1, ed. by A. Padwa, John Wiley, 1983, Chapt. 4.
6. All reactions were performed with 5 equivalents of diazomethane, in Et₂O at room temperature.
7. Exo selectivity is the rule in the 1,3-dipolar cycloaddition reactions to bicyclic systems, see, R. Huisgen, *Pure*

- Appl. Chem.*, 1981, **53**, 171. For an exception, see, M. Burdiso, A. Gamba, R. Gandolfi, and P. Pevarello, *Tetrahedron*, 1987, **43**, 1835.
8. R. S. Bly, F. Barton Culp, and R. K. Bly, *J. Org. Chem.*, 1970, **35**, 2235.
9. W. Adam, N. Carballeira, E. Crämer, V. Lucchini, E. M. Peters, K. Peters, and H. G. von Schnering, *Chem. Ber.*, 1987, **120**, 695.
10. Y. Auberson, R. M. Bimwala, and P. Vogel, *Tetrahedron Lett.*, 1991, **32**, 1637.
11. K. A. Black and P. Vogel, *Helv. Chim. Acta*, 1984, **67**, 1612.
12. For instance, isomer **2a**: mp 153-155 °C; ir, (KBr) $\nu_{\max}$, 2270, 1740; ^1H -nmr (CDCl_3), δ 1.91 (1H, d, J_{xn} 15.1 Hz, H-3n), 2.23 (3H, s, CH_3), 2.25 (1H, ddd, $J_{5,6}$ 7.5 Hz, $J_{5,\alpha}$ 9.6 Hz, $J_{5,\beta}$ 3.6 Hz, H-5), 2.71 (1H, dd, J_{xn} 15.1 Hz, $J_{\alpha,4}$ 6.0 Hz, H-3x), 4.41 (1H, d, $J_{\alpha,4}$ 6.0 Hz, H-4), 4.41 (1H, ddd, $J_{\beta,\alpha}$ 18.3 Hz, $J_{\beta,5}$ 3.5 Hz, $J_{\beta,6}$ 2.7 Hz, H- β), 4.68 (1H, ddd, $J_{\alpha,\beta}$ 18.3 Hz, $J_{\alpha,5}$ 9.6 Hz, $J_{\alpha,6}$ 1.2 Hz, H- α), 5.23 (1H, ddd, $J_{5,6}$ 7.5 Hz, $J_{6,\alpha}$ 1.2 Hz, $J_{6,\beta}$ 2.7 Hz, H-6), 5.38 (1H, s H-1); ^{13}C -nmr (CDCl_3), δ 20.3, 38.6, 43.3, 73.4, 81.5, 82.2, 83.0, 90.5, 117.6, 168.8. Anal. Calcd for $\text{C}_{10}\text{H}_{11}\text{O}_3\text{N}_3$, C, 54.30; H, 4.98; N, 19.00. Found, C, 54.26; H, 4.77; N, 18.78. Isomer **2a**: mp 122-123 °C; ir, (KBr) $\nu_{\max}$, 2250, 1735; ^1H -nmr (CDCl_3), δ 2.09 (1H, d, J_{xn} 15.1 Hz, H-3n), 2.18 (3H, s, CH_3), 2.46 (1H, ddd, $J_{5,6}$ 7.5 Hz, $J_{6,\alpha}$ 9.6 Hz, $J_{6,\beta}$ 3.6 Hz, H-6), 2.82 (1H, dd, J_{xn} 15.1 Hz, $J_{\alpha,4}$ 6.0 Hz, H-3x), 4.41 (1H, ddd, $J_{\beta,\alpha}$ 18.3 Hz, $J_{\beta,6}$ 3.6 Hz, $J_{\beta,5}$ 2.7 Hz, H- β), 4.68 (1H, ddd, $J_{\alpha,\beta}$ 18.3 Hz, $J_{\alpha,6}$ 9.6 Hz, $J_{\alpha,5}$ 1.2 Hz, H- α), 4.79 (1H, s H-1), 5.00 (1H, ddd, $J_{5,6}$ 7.5 Hz, $J_{5,\alpha}$ 1.2 Hz, $J_{5,\beta}$ 2.7 Hz, H-5), 5.30 (1H, d, $J_{\alpha,4}$ 6.0 Hz, H-4), ; ^{13}C -nmr (CDCl_3), δ 20.4 (q, 1J 133 Hz), 33.0 (d, 1J 148 Hz), 42.0 (t, 1J 140 Hz), 78.1 (d, 1J 155 Hz), 82.0 (t, 1J , 142 Hz), 85.8 (d, 1J 175 Hz), 95.0 (d 1J , 150 Hz), 117.8 (s), 168.8(s). Anal. Calcd. for $\text{C}_{10}\text{H}_{11}\text{O}_3\text{N}_3$, C, 54.30; H, 4.98; N, 19.00. Found, C, 54.20; H, 4.75; N, 18.80.
13. R. Huisgen in ref. 5, Vol. 1, Chp. 1.
14. M. J. S. Dewar and W. Thiel, *J. Am. Chem. Soc.*, 1977, **99**, 4907.
15. Program MONSTERGAUSS 81, R. A. Poirier and R. M. Peterson, University of Toronto.
16. FMO's energies and coefficients of diazomethane have been previously evaluated by Houk *et al.*, $E_{\text{HO}} = -9.0$ eV (PES), $E_{\text{LU}} = 1.8$ eV (estimated). See, K. N. Houk and K. Yamaguchi, Ref. 5, Vol. 2, p. 423 and references therein.
17. J. Geitner, R. Huisgen, and R. Sustmann, *Tetrahedron Lett.*, 1977, 881 and references therein.
18. Ref. 13, p. 31 and references therein.

Received, 2nd November, 1992