

FIVE PHENOLIC COMPOUNDS IN THE UNDERGROUND PARTS OF *VANCOUVERIA HEXANDRA*

Munekazu Iinuma^{a*}, Yoko Kanie^a, Toshiyuki Tanaka^a, Mizuo Mizuno^a
and Frank A. Lang^b

^aDepartment of Pharmacognosy, Gifu Pharmaceutical University, 6-1
Mitahora-higashi 5 chome, Gifu 502, Japan. ^bDepartment of Biology,
Southern Oregon State College, 1250 Siskiyou, Boulevard, Ashland, OR
97520-5071, U.S.A.

Abstract ----- From the underground parts of *Vancouveria hexandra*, a new
2-phenoxybenzochromone and four new prenylated flavones were
isolated. Their structures were determined by the spectral analysis.

In our previous papers, the occurrence of new glycosides of flavonol with a γ,γ -dimethylallyl group in some *Epimedium* species,¹⁻⁵ and the presence of new cognate glycosides in *Vancouveria hexandra*⁶⁻⁸ were described. In continuation of our study on the chemotaxonomy of Epimediaceae (Berberidaceae), in particular, between *Epimedium* and *Vancouveria*, we tried to isolate of non-glycosidic phenolic compounds from a lesser polar fraction in the underground parts of *V. hexandra*, which resulted in the isolation of five new compounds. We deal with the structural elucidation of these compounds in this paper.

The underground parts of *Vancouveria hexandra* C. Morr. & Decne. were extracted with MeOH. After partition of the extract with EtOAc and *n*-BuOH, the *n*-BuOH soluble portion was subjected to silica gel column chromatography (Si CC). The fraction eluted with CHCl₃-MeOH (5 : 1) was further purified with Si CC and preparative tlc to give five new compounds (1-5).

Compound (1) was obtained as a yellow amorphous, and gave M^+ at m/z 300. In the ¹H nmr spectrum, two hydroxyl groups [δ 9.91 and 12.90 (chelated)] and a methoxyl group (δ 3.92) were observed. A set of *meta*








Figure 1



A = 2-hydroxy-3-methylbut-4-enyl

Table 1 ¹³C nmr spectral data of 1^a

carbon No.	ppm
2	168.6
3	98.8
4	183.6
5	161.6
6	93.2
7	165.4
8	87.3
9	155.3
10	103.4
1'	143.5
2', 6'	122.3
3', 5'	117.0
4'	156.6
OMe	56.6

a : Measured in DMSO-*d*₆, and assignment was referred to ref. 12.

coupled one-proton doublets [δ 6.47 and 6.71 (each $J = 2$ Hz)], a set of *ortho*-coupled two-proton doublet [6.95 and 7.26 (each $J = 9$ Hz)], and a one-proton singlet (δ 5.16) were further observed. Among them, the *ortho*-coupled protons were assigned to those on a *p*-oxygenated benzene ring. The singlet at δ 5.16 was assignable to H-3 in 2-phenoxybenzochromone.⁹ These above data suggested that **1** was a 5,7,4'-trioxygenated 2-phenoxybenzochromone. The problem was the position of the methoxyl group, which was made clear as follows. In the eims, a fragment ion at m/z 167 (Figure 1) which corresponds to an [A1⁺ + H] fragment after retro Diels-Alder cleavage such as flavonoid compound suggested that the methoxyl group and one hydroxyl group were substituted on the A ring. Furthermore, the signals based on a chelated hydroxy

Table 2 ^{13}C nmr spectral data of compounds 2-5

carbon No.	2*	3**	carbon No.	4***	5***
2	162.5	163.5	2	165.5	164.7
3	104.0	103.3	3	104.2	103.6
4	182.4	181.6	4	183.2	182.7
5	162.3	161.4	5	160.9	160.3
6	99.5	98.8	6	99.9	109.7
7	164.1	164.1	7	165.5	160.4
8	94.2	93.8	8	95.6	94.2
9	157.8	157.2	9	158.0	156.7
10	105.3	103.7	10	105.6	104.7
1'	123.5	122.7a	1'	123.2	122.3
2'	124.5	122.0	2'	128.0	127.0
3'	121.5	121.4a	3'	131.1	130.5
4'	156.5	153.7	4'	163.4	162.9
5'	117.0	129.3	5'	117.7	117.3
6'	127.6	127.3	6'	127.5	127.4
4''	121.4	120.6	1'', 1'''	38.9	38.8b
5''	131.7	131.6	2'', 2'''	77.0	76.3, 76.4
6''	76.5	77.2	3'', 3'''	148.4	147.6, 147.7
Me	28.3 (x 2)	28.6	4'', 4'''	111.2	110.2, 110.7
1''		35.6	5'', 5'''	18.6	18.3, 18.4
2''		73.4			
3''		148.0			
4''		109.8			
5''		17.7			

*CDCl₃, **DMSO-d₆, ***acetone-d₆

a: Maybe interchangeable, b: Another carbon was buried in solvent signals.

group and on a typical set of one-proton *meta*-coupled doublets in the ^1H nmr spectrum indicated that the A ring was oxygenated at C-5 and C-7. When **1** was compared with another known 2-phenoxybenzochromone possessing a 5,7-dihydroxyl A ring isolated from *Artemisia capillaris*,⁹ the protons at C-6 and C-8 of **1** were shifted in a lower field than those of known one. These data suggested that the A ring was a 5-hydroxy-7-methoxyl substitution, which was finally estimated by an NOE experiment. On irradiation of the methoxyl group, NOEs were observed at both H-6 (19%) and H-8 (20%), respectively. The structure of **1** was, consequently, concluded to be 2-(*p*-hydroxyphenoxy)-5-hydroxy-7-methoxybenzochromone. To the best of our knowledge, 2-phenoxy-benzochromones were rare phenolics in plant kingdom and a few instances have been confirmed in Compositae⁹⁻¹¹ and Rosaceae.¹² 2-Phenoxybenzochromone with a 5-hydroxy-7-methoxy moiety has been synthetically derived, however, it is a first isolation of such a derivative as natural product. The ^{13}C nmr spectral data are shown in Table 1.

Compounds (2-5) were suggested to be a flavone compound by their uv spectral data. Compound (2) was obtained as a yellow amorphous, and gave M^+ at m/z 336. In the 1H nmr spectrum, a set of *ortho*-coupled olefinic protons [δ 5.71, 6.38 (each $J = 10$ Hz)] and two methyls (δ 1.48) were observed, indicating the presence of a dimethylpyran ring. The base peak at m/z 321 in the Eims showed the ring was fused on the ring B. In the 1H nmr spectrum, two hydroxyl groups [δ 10.20 and 13.00 (chelated)], a set of *meta*-coupled one-proton doublets were supported the A ring of 2 to be a 5,7-dihydroxyl substitution. Furthermore, a typical ABX system at δ 6.83 (1H, d, $J = 9$ Hz), 7.48 (1H, d, $J = 2$ Hz) and 7.63 (1H, dd, $J = 9, 2$ Hz) was observed in the 1H nmr spectrum. These data indicated that the dimethylchromene ring was fused on the B ring and formed the ring between C-3' and 4'-OH. Consequently, the structure of 2 was determined to be 5,7-dihydroxy-6",6"-dimethylpyrano(2",3" : 4',3')flavone.

Compound (3) was obtained as a yellow amorphous, and gave M^+ at m/z 420. All spectral data suggested that 3 had a same A ring and such a dimethylchromene ring system as 2. The presence of another 2-hydroxy-3-methylbut-4-enyl group was indicated by the following signals in the 1H nmr spectrum : δ 1.77 (3H, s, Me), 2.70 (1H, dd, $J = 14, 9$ Hz), 3.02 (1H, dd, $J = 14, 5$ Hz), 4.18 (1H, m), 4.70 and 4.80 (1H each, br s, =CH₂). Further protons of the ring B were observed as two *meta*-coupled doublets [δ 7.66 and 7.70 ($J = 2$ Hz)]. These data suggested another 2-hydroxy-3-methylbut-4-enyl group was attached at C-5' of 2. Consequently, the structure of 3 was confirmed to be 5,7-dihydroxy-5'-(2-hydroxy-3-methylbut-4-enyl)-6",6"-dimethylpyrano(2",3" : 4',3')flavone.

Compounds (4) and (5) were obtained as a yellow amorphous. Compound (4) gave M^+ at m/z 354, and the uv and 1H nmr spectra indicated that 4 was an apigenin with a substituent at C-3'. In the 1H nmr spectrum, the substituent was regarded as a 2-hydroxy-3-methylbut-4-enyl group. The structure of 4 was then determined to be 5,7,4'-trihydroxy-3'-(2-hydroxy-3-methylbut-4-enyl)flavone. Compound (5) was a flavone substituted further with another 2-hydroxy-3-methylbut-4-enyl group on the A ring compared with 4. The position of the group was determined to be at C-6 on the A ring, because the chemical shift of the chelated hydroxyl group at C-5 was appeared at δ 13.47 which is in a lower field than those of 2-4.¹³ The structure of 5 was, therefore, characterized as 5,7,4'-trihydroxy-6,3'-di(2-hydroxy-3-methylbut-4-enyl)flavone.

EXPERIMENTAL

Plant material and extraction were described in a preceding paper.⁷

Isolation of compounds 1-5. The CHCl₃-MeOH (5:1) fraction was further chromatographed on silica gel using some CHCl₃-MeOH solvent systems and finally purified by use of preparative tlc to give **1** (14 mg), **2** (22 mg), **3** (8 mg), **4** (12 mg) and **5** (10mg) as pure form.

Compound 1 [2-(*p*-hydroxyphenoxy)-5-hydroxy-7-methoxybenzochromone]: A pale yellow amorphous; Hrms Calcd for C₁₆H₁₂O₆: 300.0634. Found: 300.0612; Eims *m/z* (rel. int.): 300 (M⁺, 100), 271 (20), 208 (2), 179 (2), 167 (27); uv (nm, MeOH): 229, 283; ¹H nmr (DMSO-*d*₆) δ: 3.92 (3H, s, OMe), 5.16 (1H, s, H-3), 6.47 (1H, d, *J* = 2 Hz, H-6), 6.71 (1H, d, *J* = 2 Hz, H-8), 6.95 (2H, d, *J* = 9 Hz, H-3' and 5'), 7.26 (2H, d, *J* = 9 Hz, H-2' and 6'), 9.91 (1H, s, C4'-OH), 12.90 (1H, s, C5-OH).

Compound 2 [5,7-dihydroxy-6",6"-dimethylpyrano(2",3" : 4',3')flavone]: A yellow amorphous; Hrms Calcd for C₂₀H₁₆O₅: 336.1004. Found: 336.1025; Eims *m/z* (rel. int.): 336 (M⁺, 27), 321 (100), 300 (11), 272 (84), 169 (8), 152 (7); uv (nm, MeOH): 268, 342, +NaOMe: 226, 275, 312, 362, +AlCl₃: 279, 300sh, 362, 389, +AlCl₃/HCl: 279, 300sh, 357, 388, +NaOAc: 264, 274sh, 310, 369, +NaOAc/H₃BO₃: 269, 345; ¹H nmr (DMSO-*d*₆) δ: 1.48 (6H, s, 2 x Me), 5.71 (1H, d, *J* = 10 Hz, H-5"), 6.31 (1H, d, *J* = 2 Hz, H-6), 6.38 (1H, d, *J* = 10 Hz, H-4"), 6.45 (1H, d, *J* = 2 Hz, H-8), 6.54 (1H, s, H-3), 6.83 (1H, d, *J* = 9 Hz, H-5'), 7.48 (1H, d, *J* = 2 Hz, H-2'), 7.63 (1H, dd, *J* = 9, 2 Hz, H-6'), 10.20 (1H, s, C7-OH), 13.00 (1H, s, C5-OH).

Compound 3 [5,7-dihydroxy-5'-(2-hydroxy-3-methylbut-4-enyl)-6",6"-dimethylpyrano(2",3" : 4',3')flavone]: A yellow amorphous; Hrms Calcd for C₂₅H₂₄O₆: 420.1550. Found: 420.1578; Eims *m/z* (rel. int.): 420 (M⁺, 47), 405 (51), 350 (100), 335 (64), 153 (24); uv (nm, MeOH): 242, 269, 347, +NaOMe: 269, 315, 372, +AlCl₃: 277, 360, 387, +AlCl₃/HCl: 278, 360, 387, +NaOAc: 274sh, 318, 361, +NaOAc/H₃BO₃: 269, 351; ¹H nmr (DMSO-*d*₆) δ: 1.43 (6H, s, 2 x Me), 1.77 (3H, s, H-5"), 2.70 (1H, dd, *J* = 14, 9 Hz, H-1"), 3.02 (1H, dd, *J* = 14, 5 Hz, H-1"), 4.18 (1H, m, H-2"), 4.70 (1H, br s, H-4"), 4.80 (2H, br s, H-4" and OH), 5.84 (1H, d, *J* = 10 Hz, H-5"), 6.19 (1H, d, *J* = 2 Hz, H-6), 6.47 (1H, d, *J* = 2 Hz, H-8), 6.49 (1H, d, *J* = 10 Hz, H-4"), 6.76 (1H, s, H-3), 7.66, 7.70 (1H each, d, *J* = 2 Hz, H-2' and 6'), 10.80 (1H, s, C7-OH), 12.92 (1H, s, C5-OH).

Compound 4 [5,7,4'-trihydroxy-3'-(2-hydroxy-3-methylbut-4-enyl)flavone]: A yellow amorphous; Hrms Calcd for C₂₀H₁₈O₆: 354.1103. Found: 354.1124; Eims *m/z* (rel. int.): 354 (M⁺, 36), 336 (9), 321 (22), 284 (100), 155 (16), 153 (22); uv (nm, MeOH): 268, 340, +NaOMe: 256, 323, 397, +AlCl₃: 251, 276, 301,

352, 400, +AlCl₃/HCl: 276, 300, 351, 400, +NaOAc: 275, 301, 379; ¹H nmr (acetone-*d*₆) δ: 1.20 (3H, br s, H-5"), 2.70 (1H, dd, *J*= 15, 9 Hz, H-1"), 3.02 (1H, dd, *J*= 15, 4 Hz, H-1"), 4.48 (1H, m, H-2"), 4.80, 5.00 (1H each, br s, H-4"), 6.25 (1H, d, *J*= 2 Hz, H-6), 6.53 (1H, d, *J*= 2 Hz, H-8), 6.63 (1H, s, H-3), 7.00 (1H, d, *J*= 9 Hz, H-5'), 7.82 (1H, dd, *J*= 9, 2 Hz, H-6'), 7.88 (1H, d, *J*= 2 Hz, H-2'), 9.80 (1H, s, C7-OH), 13.00 (1H, s, C5-OH).

Compound 5 [5,7,4'-trihydroxy-6,3'-di(2-hydroxy-3-methylbut-4-enyl)flavone]: A yellow amorphous ; Hrms Calcd for C₂₅H₂₆O₇: 438.1678. Found: 438.1648; Eims *m/z* (rel. int.): 438 (M⁺, 7), 420 (11), 405 (6), 387 (367 (100), 350 (27), 206 (18), 286 (18), 165 (22), 123 (13); uv (nm, MeOH): 274, 336, +NaOMe: 275, 330, 395, +AlCl₃: 285, 310, 370, +AlCl₃/HCl: 276, 310, 370, +NaOAc: 270, 309, 370, +NaOAc/H₃BO₃: 271, 336 ; ¹H nmr (acetone-*d*₆) δ: 1.19 (6H, br s, 2 x Me, H-5" and 5""), 2.85-3.05 (4H, m, H-1" and 1""), 4.42 (2H, m, H-2" and 2""), 4.77, 4.79, 4.95, 4.99 (1H each, br s, H-4" and 4""), 6.50 (1H, s, H-8), 6.58 (1H, s, H-3), 6.96 (1H, d, *J*= 9 Hz, H-5'), 7.78 (1H, dd, *J*= 9, 2 Hz, H-6'), 7.82 (1H, d, *J*= 2 Hz, H-2'), 13.47 (1H, s, C5-OH).

ACKNOWLEDGMENT

This study was supported by Grant in Aid No. 03042017 from the Ministry of Education, Arts and Sciences, Japan.

REFERENCES

1. M. Mizuno, M. Iinuma, T. Tanaka, and N. Sakakibara, *J. Nat. Prod.*, 1990, **53**, 744.
2. M. Mizuno, M. Iinuma, T. Tanaka, N. Sakakibara, M. Nishi, A. Inada, and T. Nakanishi, *Phytochemistry*, 1989, **28**, 2527.
3. M. Mizuno, M. Iinuma, T. Tanka, N. Sakakibara, T. Nakanishi, A. Inada, and M. Nishi, *Chem. Pharm. Bull.*, 1989, **37**, 2241.
4. M. Mizuno, M. Iinuma, T. Tanaka, S. Iwashima, N. Sakakibara, X. Liu, D. Shi, and H. Müssel, *Asian J. Plant Sci.*, 1989, **1**, 1.
5. M. Mizuno, M. Iinuma, T. Tanaka, N. Sakakibara, S. Hanioka, and X. Liu, *Chem. Pharm. Bull.*,

1988, **36**, 3487.

6. M. Mizuno, M. Iinuma, T. Tanaka, N. Sakakibara, J. Murata, H. Murata, and F.A. Lang, *Phytochemistry*, 1990, **29**, 1277.
7. M. Mizuno, Y. Kanie, M. Iinuma, T. Tanaka, and F.A. Lang, *Phytochemistry*, 1991, **30**, 2765.
8. M. Mizuno, Y. Kanie, M. Iinuma, T. Tanaka, and F.A. Lang, *Phytochemistry*, 1992, **31**, 297.
9. T. Komiya, Y. Naruse, and H. Oshio, *Yakugaku Zasshi*, 1976, **96**, 855.
10. T. Komiya, M. Tsukui, and H. Oshio, *Yakugaku Zasshi*, 1976, **96**, 841.
11. E.K. Adesogan and A.L. Okunade, *J. Chem. Soc., Chem. Comm.*, 1978, 152.
12. H. Hashidoko, S. Tahara, and J. Mizutani, *Phytochemistry*, 1991, **30**, 3837.
13. T. Fukai and T. Nomura, *Heterocycles*, 1991, **32**, 499.

Received, 12th November, 1992