

**DIELS-ALDER CYCLOADDITION OF 2(1*H*)-PYRIDONES
HAVING TWO ELECTRON-WITHDRAWING GROUPS WITH
ELECTRON DEFICIENT DIENOPHILE**

Hiroto Nakano and Hiroshi Hongo*

Tohoku College of Pharmacy


4-4-1 Komatsushima, Aoba-ku, Sendai 981, Japan

Abstract- The first examples are presented of Diels-Alder cycloadditions of 2(1*H*)-pyridones having two electron-withdrawing groups on the ring, with electron deficient dienophiles, and the adducts obtained may act as dienophiles leading to richly functionalized, bridged, and polycyclic aza-compounds, under high pressure conditions.

2(1*H*)-Pyridones are classified as aromatic heterocycles, and generally do not undergo efficient Diels-Alder cycloaddition. Nevertheless, the Diels-Alder adducts from the reactions of 2(1*H*)-pyridones with dienophiles have an isoquinuclidine skeleton, and therefore have potential as synthetic intermediates.^{1h} Recently we reported the first examples of Diels-Alder cycloadditions using, as dienes, 2(1*H*)-pyridones substituted by an electron-withdrawing group on the ring, with electron deficient dienophiles.^{1l} Furthermore, the *endo* Diels-Alder adducts obtained may act as dienophiles leading to bridged and polycyclic aza-compounds.^{1l} Although inverse-electron-demand [2+4] cycloadditions of 1,3-disulfonyl-2(1*H*)-pyridones with electron rich dienophiles were reported only by Posner and his co-worker,² there are few publications relating to [4+2] cycloadditions of 2(1*H*)-pyridones possessing two electron-withdrawing groups on the ring, with electron deficient dienophiles. Herein we wish to report Diels-Alder cycloaddition of 1-methyl-2(1*H*)-pyridones (**1a-d**) having two methoxycarbonyl groups on the ring, with *N*-

Dedicated to Prof. E. C. Taylor on the occasion of his 70th birthday.


phenylmaleimide (**2**), and these reactions are the first examples of [4+2] cycloadditions of 2(1*H*)-pyridones substituted by two electron-withdrawing groups, with electron deficient dienophiles. The Diels-Alder adducts (**3a-d**) obtained are potentially valuable synthetic intermediates. In fact, the compounds (**3dn** and **3dx**) act as dienophiles under high pressure conditions³ leading to bridged and polycyclic aza-compounds (**4a** and **4b**), which are unobtainable under atmospheric pressure conditions.



Scheme 1


A mixture of 2(1*H*)-pyridones (**1a-c**) and the dienophile (**2**) (5 equiv.) was heated at 110°C for 72 h, and the *endo* adducts (**3a-c**) were stereoselectively obtained in 23, 71, and 17 % yields, respectively. However, the same reaction of **1d** with **2** afforded a mixture of the *endo* [**3dn**(43%)] and the *exo* [**3dx**(33%)] adducts. The stereochemistry of the *endo* **3a-dn** ($J_{1,6}=J_{4,5}=4$ Hz) and the *exo* **3dx** ($J_{1,6}=J_{4,5}=2$ Hz) adducts follows from the coupling constants. In general, the coupling constants of *endo*-H and *exo*-H coupling with bridgehead protons are 3.5 Hz and under ¹d,k,l,m and 3.5-4.5 Hz, ¹d,k,l,m respectively, in isoquinuclidine derivatives (Scheme 1). Diels-Alder cycloadditions of the adducts (**3dn** and **3dx**) with 2,3-dimethylbuta-1,3-diene (**5**) (3

equiv.) under high pressure conditions (10 Kbar, 70°C, 48 h, methylene dichloride) stereoselectively afforded the corresponding *cis-anti-cis* adducts [**4a** (54% ; mp 177-179°C) and **4b** (96% ; mp 230-232°C)] (Scheme 2).



Scheme 2

Considering that the adducts (**4a** and **4b**) were stereoselectively obtained, the *cis-anti-cis* stereochemistry of **4a,b** was deduced from the basis of steric factor ; the diene (**5**) attacked from less hindered side¹¹ of the *endo* adduct (**3dn**) (upper side a) and the *exo* adduct (**3dx**) (under side b), respectively (Scheme 3).



Scheme 3

The structures of **3a-d** and **4a,b** were confirmed by ir and ^1H -nmr spectroscopy and gave satisfactory elemental analyses.

REFERENCES

1. a) E. B. Sheinin, G. E. Wright, C. L. Bell, and L. Bauer, *J. Heterocycl. Chem.*, 1968, **5**, 859; b) H. Tomisawa and H. Hongo, *Tetrahedron Lett.*, 1969, 2465; c) U. Heep, *Tetrahedron*, 1975, **31**, 77 ; d) K. Somekawa, T. Watanabe, and S. Kumamoto, *Nippon Kagaku Kaishi*, 1978, 412; e) P. S. Mariano, P. L. Huesman, and R. L. Beamer, *Tetrahedron*, 1978, **34**, 2617; f) G. P. Gisby, S. E. Royall, and P. G. Sammes, *J. Chem. Soc., Chem. Commun.*, 1979, 501; g) K. Matsumoto, Y. Ikemi, T. Uchida, and R. M. Acheson, *J. Chem. Soc., Chem. Commun.*, 1979, 1091; h) H. Tomisawa, H. Hongo, H. Kato, K. Sato, and R. Fujita, *Heterocycles*, 1981, **16**, 1947; i) G. P. Gisby, S. E. Royall, and P. G. Sammes, *J. Chem. Soc., Perkin Trans. 1*, 1982, 169; j) M. Kuzuya, A. Noguchi, E. Mano, and T. Okuda, *Bull. Chem. Soc. Jpn.*, 1985, **58**, 1149; k) H. Tomisawa, H. Nakano, and H. Hongo, *Heterocycles*, 1990, **30**, 359. l) H. Nakano, H. Tomisawa, and H. Hongo, *J. Chem. Soc., Chem. Commun.*, 1990, 1775. m) G. H. Posner, V. Vinader, and K. Afarinkia, *J. Org. Chem.*, 1992, **57**, 4088.
2. G. H. Posner and C. Switzer, *J. Org. Chem.*, 1987, **52**, 1642.
3. a) K. Matsumoto, A. Sera, and T. Uchida, *Synthesis*, 1985, 1; b) K. Matsumoto and A. Sera, *ibid.*, 1985, 999; c) W. J. le. Noble (ed.), "Organic High Pressure Chemistry" Elsevier, Amsterdam, 1988.

Received, 7th September, 1992