


3-ARYL-3H-1,2-BENZODITHIOLES. SYNTHESIS AND REACTION WITH DICHLOROCARBENE AND CARBENOID

Shin-ichi Chida and Ryu Sato*

Department of Applied Chemistry and Molecular Science, Faculty of Engineering,
Iwate University, Morioka 020, Japan

Abstract - New stable 3H-1,2-benzodithiole derivatives, 3-aryl-3H-1,2-benzodithioles, were synthesized. While the reaction of the 3-aryl-3H-1,2-benzodithioles with $\text{CH}_2\text{I}_2\text{-Et}_2\text{Zn}$ gave methylene-inserted products 4-aryl-2,4-dihydro-1,3-benzodithiins, the reaction with dichlorocarbene afforded unexpected products 2-chloro-3-arylbenzo[*b*]thiophenes.

Recently, the attention of many organic sulfur chemists has been directed to study of the biological activity as well as the unique reactivity of cyclic polysulfides such as benzopentathiepin.¹ We have succeeded in the synthesis of a variety of cyclic polysulfides by the reaction of the corresponding dimercaptanes with elemental sulfur / ammonia and have examined their reactivities to give many heterocycles involving sulfur atoms.² Our current interest has focused on the chemistry of new types of cyclic polysulfides containing a benzylmethylene group in the polysulfides ring, e.g., 6H-benzo[*g*]-1,2,3,4,5-pentathiocin (**BPTC**), 5H-benzo[*f*]-1,2,3,4-tetrathiepin (**BTTP**), and 3H-benzo-1,2-dithiole (**1a**).³ Here we wish to report a synthesis of new stable 3H-1,2-benzodithiole derivatives having an aryl substituent at 3-position⁴ and the reactions with electrophilic agents such as dichlorocarbene and carbenoid.




Scheme 1

Treatment of the alcohol (**4**), which was derived from thiophenol (**3**) by ortho-lithiation⁵ followed by reaction with aldehyde, with P_2S_5 afforded dimercaptanes (**2b-d**). The desired 3-aryl-3H-1,2-benzodithioles (**1b-d**) were


obtained by cyclization of **2b-d** with S_8 / NH_3 in CH_2Cl_2 ² (Scheme 2). In contrast to the reaction of 2-mercaptomethylbenzenethiol (**2a**) with S_8 / NH_3 to give a mixture of **BTTP**, **BPTC**, and **1a**³ (Scheme 1), the same treatment of **2b-d** afforded only 1,2-benzodithioles (**1b-d**). These results suggest that in the reaction the

Scheme 2



introduction of a suitable substituent at the benzyl position in **2a** controls the ring size of the cyclic polysulfides produced by the reaction of **2** with S_8 / NH_3 . The structures of all new 3*H*-1,2-benzodithioles (**1b-d**) were assigned on the basis of ¹H nmr, ir, mass spectrometry, and elemental analysis.⁶ Since the repulsion between sulfur-sulfur lone pair electrons in the 1,2-dithiole ring leads to easy cleavage of the sulfur-sulfur bond, compound (**1a**) is unstable even at room temperature and easily polymerizes. Therefore, it is difficult to isolate **1a** in pure state⁷ and the chemistry of **1a** has not been studied in detail so far.⁸ Interestingly, in contrast to **1a**, 3*H*-1,2-benzodithioles (**1b-d**) are extremely stable cyclic disulfides. Consequently, **1b-d** are suitable compounds to study the chemistry of the 3*H*-1,2-benzodithioles.⁹ To examine the reactivity toward electrophiles, **1b-d** were exposed to dichlorocarbene and carbenoid.¹⁰ Thus, treatment of **1b-d** with excess NaOH and trioctylmethylammonium chloride (TOMAC) in $CHCl_3$ at room temperature for 0.5 h gave the unexpected product, 2-chloro-3-aryl-benzo[*b*]thiophenes (**5b-d**) (Scheme 3). **5b**; pale yellow oil; ¹H nmr ($CDCl_3$) δ 6.90-7.90 (m); ir (neat) 2950, 1425, 1010, and 750 cm^{-1} ; ms m/z calcd for $C_{14}H_9ClS$ 244.0113, obsd 244.0085.


Scheme 3



On the other hand, treatment of **1b-d** with CH_2I_2 and Et_2Zn (0.98 M *n*-hexane solution) in ether afforded methylene-inserted products 4-aryl-1,3-benzodithianes (Scheme 4). **6b**; pale yellow oil; ¹H nmr ($CDCl_3$) δ 3.82


(s, 2H, $-\text{CH}_2-$), 5.23 (s, 1H, $-\text{CH}<$), 6.73-7.63 (m, 9H, arom); ir (neat) 2920, 1439, and 706 cm^{-1} ; ms m/z calcd for $\text{C}_{14}\text{H}_{12}\text{S}_2$ 244.0380, obsd 244.0328.

Scheme 4



From these results, a plausible reaction pathway to form **5** from **1** with dichlorocarbene is shown in Scheme 5.

Scheme 5



The dichlorocarbene-inserted product (**A**), which was formed from **1** with dichlorocarbene, is deprotonated with excess NaOH followed by ring-contraction accompanied with elimination of chloride and aromatization accompanied with desulfurization to give **5**.

Further study on the reactivity of the 3*H*-1,2-benzodithioles (**1b-d**) is now in progress.

ACKNOWLEDGEMENT

This work was supported by a Grant-in-Aid for Scientific Research No. 04640483 from the Ministry of Education, Science and Culture.

REFERENCES

- For example, B. L. Chenard and T. J. Miller, *J. Org. Chem.*, 1984, **49**, 1222; B. L. Chenard, R. L. Harlow, A. L. Johnson, and S. A. Vladuchick, *J. Am. Chem. Soc.*, 1985, **107**, 3871; M. Litaudon and M. Guyot, *Tetrahedron Lett.*, 1991, **32**, 911; B. S. Davidson, T. F. Molinski, L. R. Barrows, and C. M. Ireland, *J. Am. Chem. Soc.*, 1991, **113**, 4709.

2. R. Sato, T. Goto, and M. Saito, *J. Synth. Org. Chem. Jpn.*, 1990, **48**, 797, and references cited in this review; R. Sato and K. Chino, *Tetrahedron Lett.*, 1991, **32**, 6345; R. Sato, J. Kumagai, K. Yonesaka, and S. Satoh, *ibid.*, 1992, **33**, 947.
3. R. Sato, S. Satoh, and M. Saito, *Chem. Lett.*, 1990, 139; R. Sato and S. Satoh, *Synthesis*, 1991, 785; S. Satoh and R. Sato, *Tetrahedron Lett.*, 1992, **33**, 2517.
4. M-F. Chan and M. E. Garst, *J. Chem. Soc., Chem. Commun.*, 1991, 540.
5. G. D. Figuly, C. K. Loop, and J. C. Martin, *J. Am. Chem. Soc.*, 1989, **111**, 654.
6. **1b** (Ar=Phenyl); Yellow oil; ^1H nmr (CDCl_3) δ 5.96 (s, 1H, Ph-CH) and 6.86-7.60 (m, 9H, arom); ir (neat) 3000, 1570, 1440, 730, and 690 cm^{-1} ; ms (m/z) 230(M^+); uv λ_{max} (MeCN) 367.5 nm (ϵ 407); *Anal.* Calcd for $\text{C}_{13}\text{H}_{10}\text{S}_2$ C, 67.58; H, 4.67. Found: C, 67.71; H, 4.33. **1c** (Ar=*o*-Methoxyphenyl); Yellow needles; mp 97°C (from n-hexane); ^1H nmr (CDCl_3) δ 3.88(s, 3H, -OCH₃), 6.30(s, 1H, -CH-), and 6.60-7.45(m, 8H, arom); ir (neat) 2925, 1489, 1461, 1248 and 747 cm^{-1} ; ms (m/z) 260(M^+); uv λ_{max} (MeCN) 364.5 nm (ϵ 322); *Anal.* Calcd for $\text{C}_{14}\text{H}_{12}\text{OS}_2$ C, 64.58; H, 4.65. Found: C, 64.44; H, 4.67. **1d** (Ar=Mesityl); Yellow oil; ^1H nmr (CDCl_3) δ 2.15(s, 3H, *o*-CH₃), 2.25(s, 3H, *p*-CH₃), 2.42(s, 3H, *o*-CH₃), and 6.50-7.20(m, 5H, Ms-CH- and arom); ir (neat) 2918, 1609, 1438, and 743 cm^{-1} ; uv (MeCN) λ_{max} 393.5 nm (ϵ 507); ms m/z calcd for $\text{C}_{16}\text{H}_{16}\text{S}_2$ 272.0693, obsd 272.0696.
7. A. Lüttringhaus and K. Hägele, *Angew. Chem.*, 1955, **67**, 304.
8. A. G. Hortmann, A. J. Aron, and A. K. Bhattacharya, *J. Org. Chem.*, 1978, **43**, 3374.
9. To obtain the structural information, the theoretical calculations of **1a-d** were carried out using MM2(77) and MOPAC Ver.5.01 [MOPAC Ver. 5.00 (QCPE No. 445), J. J. P. Stewart, QCPE Bull., 1989, 9, 10; T. Hirano, *JCPE Newsletter*, 1989, **1**, 36; Revised as Ver. 5.01 by J. Toyoda, for Apple Macintosh] programs. The calculation results are shown below.

Dithiole	Calcd. S-S bond length (Å)	Calcd. CSSC dihedral angle (deg.)	Energy level of HOMO (eV)
1a	2.017	36.9	-8.59
1b	2.015	36.6	-8.53
1c	2.014	36.2	-8.39
1d	2.011	32.1	-8.40

10. S. Tamagaki and S. Oae, *Bull. Chem. Soc. Jpn.*, 1972, **45**, 960; W. Ando, Y. Kumamoto, and T. Takata, *Tetrahedron Lett.*, 1985, **26**, 5187; T. Ghosh, *J. Org. Chem.*, 1990, **55**, 1146.

Received, 6th January, 1993