

**PHOTOCHEMICAL REACTION OF 6-ALKOXY-7-(2-OXOALKYL)-
QUINOLINE-5,8-QUINONES**

Katsuhiko Nakagawa^{*}, Hirofumi Kaneko, Takashi Yano,
Shigehiro Inai, Kenzo Tabuchi, and Shozo Shimada^{**}


Department of Industrial Chemistry, Niihama National
College of Technology, Niihama, Ehime 792, Japan

^{**}SUNFUCO Ltd., Osaka, 530 Japan


Abstract- Photochemical reaction of 6-alkoxy-7-(2-oxoalkyl)quinoline-5,8-quinones in benzene gave a novel heterocyclic quinones, 2-alkyl-4-methylquino[6,7-d]-[1,3]dioxepine-6,11-diones via intramolecular γ -H-abstraction.

Heterocyclic quinones have attracted considerable interests from both biological and photochemical view points.¹ Among these, synthetic studies of quinones containing the quinoline-5,8-quinone moiety like streptonigrin, lavendamycin, and streptonigrone have been extensively investigated² while very little work has been done on the introduction of heterocyclic ring to the quinoline-5,8-quinone moiety. Recently, we have developed a photochemical introduction of 1,3-dioxepine³ and 1,3-oxathiepine⁴ ring into naphthoquinone nuclei. In continuation with these studies, we now report here that the synthesis of 6-alkoxy-7-(2-oxoalkyl)quinoline-5,8-quinones (**2**) and photochemical reaction of **2** in benzene gave the heterocycle-fused quinoline-5,8-quinone (**3**). The title compounds (**2**)⁵ were prepared in 77% (**2a**) and 70% (**2b**) resp. by the photochemical reaction of the corresponding 6-alkoxyquinoline-

5,8-quinones (1)⁶ with 2-methoxypropene in acetonitrile with a CuSO_4 filtered light from 400 W high-pressure Hg lamp in the presence of potassium carbonate under a nitrogen and subsequent cerium(IV) ammonium nitrate (CAN) oxidation of the photo-products in aqueous acetic acid.³




Irradiation of a benzene solution of 2a ($1.6 \times 10^{-4} \text{ mol dm}^{-3}$) with a Pyrex-filtered light from 400W high-pressure Hg lamp under a nitrogen for 3 h gave a novel cyclization product, 4-methylquino[6,7-d][1,3]-dioxepine-6,11-dione (3a) in 15% yield with 92% conversion of 2a. The structure of 3a was assigned on the basis of its spectroscopic analysis.⁷ The uv spectrum of 3a in dichloromethane showed a characteristic band at 450 nm due to 1,3-dioxepin ring-fused quinoline-5,8-quinone, in analogy with 1,3-dioxepin ring-fused 1,4-naphthoquinone.³



Similarly, irradiation of 2b gave the cyclized product (3b) in 60% yield with 86% conversion of 2b. Photochemical reactivity increased in order of $2a < 2b$ which reflected the γ -C-H bond strength. The photoreaction of 2 was sensitized by xanthene, but efficiently quenched by anthracene, indicating that these photoreaction occurred from their

triplet excited state. Accordingly, a plausible mechanism for the formation of 3 is given in Scheme 1. Photocyclization may begin with intramolecular γ -H-abstraction⁸ by the quinone carbonyl of 2 to give a biradical (4), which then cyclizes to (5). Intramolecular disproportionation of the biradical (5) produces hydroquinone (6), which is oxidized by the unreacted quinone (2)⁹ or air, to give 3.



Scheme 1

tionation of the biradical (5) produces hydroquinone (6), which is oxidized by the unreacted quinone (2)⁹ or air, to give 3.

ACKNOWLEDGEMENT

Helps given by Dr. H. Uno at Advanced Instrumentation Center for Chemical Analysis, Ehime University is greatly appreciated. The present work was partially supported by a Grant-in-Aid for Scientific Research No. 03740282 from the Ministry of Education, Science and Culture, Japan.

REFERENCES

1. H. Inoue and E. Leistner, "Biochemistry of quinones," in "The Chemistry of Quinonoid Compounds Vol. II," ed. by S. Patai and Z. Rappaport, John Wiley, London, 1988, p. 1293.
M. G. Peter, *Angew. Chem., Int. Ed. Engl.*, 1989, 28, 555.
S. G-Renault, J. Renault, P. G-Servolles, M. Baron, L. Paoletti, S. Cros, M-S. Bissery, F. Larelle, and G. Atassi, *J. Med. Chem.*, 1991, 34, 38.
K. Maruyama and A. Osuka, "Recent Advances in Photochemistry of Quinones," in "The Chemistry of Quinonoid Compounds Vol. II," ed. by S. Patai and Z. Rappaport, John Wiley, London, 1988, p. 759.
2. M. N. Preobrazhenskaya, N. V. Holpne-Kozlova, and E. I. Lazhko,

- J. Antibiot.*, 1992, **45**, 227.
- A. Godard, J-C. Rovera, F. Marsais, N. Ple, and G. Queguiner, *Tetrahedron*, 1992, **48**, 4123.
- K. V. Rao and J. W. Beach, *J. Med. Chem.*, 1991, **34**, 1871.
- F. Marsais, J-C Rovera, A. Turck, A. Godard, and G. Queguiner, *J. Chem. Soc., Perkin Trans. 1*, 1990, 2611.
- P. Molina, P. M. Fresneda, and M. Canovas, *Tetrahedron Lett.*, 1992, **33**, 2891.
- D. L. Boger and S. Nakahara, *J. Org. Chem.*, 1991, **56**, 880.
3. K. Maruyama, A. Osuka, K. Nakagawa, T. Jinsenji, and K. Tabuchi, *Chem. Lett.*, 1988, 1505.
4. K. Nakagawa, *Chem. Express.*, 1991, **6**, 177.
5. 7-Acetyl-6-methoxyquinoline-5,8-quinone (**2a**)
mp 115 °C; ^1H Nmr (90 MHz, CDCl_3) δ : 2.30 (s, 3H), 3.80 (s, 2H), 4.17 (s, 3H), 7.63 (dd, J = 8.0, 4.0 Hz, 1H), 8.63 (dd, J = 8.0, 1.80 Hz, 1H), 9.00 (dd, J = 4.0, 1.80 Hz, 1H); ir (KBr) ν_{max} : 1713, 1678, 1622, 1578, 1158 and 1100 cm^{-1} ; uv(λ_{max} , nm (ϵ), CH_2Cl_2) 245 (15500), 362 (1200); GCms (m/z): 247(M^+ +2H), 203(100%). Anal. Calcd for $\text{C}_{13}\text{H}_{11}\text{NO}_4$: C, 63.66; H, 4.52; N, 5.71. Found: C, 63.39; H, 4.38; N, 5.64.
6. Y. T. Pratt and N. L. Drake, *J. Am. Chem. Soc.*, 1955, **77**, 37.
7. 4-Methylquino[6,7-d][1,3]dioxepine-6,11-dione (**3a**)
mp 215 °C; ^1H Nmr (90 MHz, CDCl_3) δ : 2.17 (s, 3H), 5.53 (s, 2H), 6.10 (s, 3H), 7.65 (dd, J = 8.3, 4.8 Hz, 1H), 8.45 (dd, J = 8.1, 1.0 Hz, 1H), 9.00 (dd, J = 4.8, 1.0 Hz, 1H); ir (KBr) ν_{max} : 1682, 1667, 1626, 1568, 1100 cm^{-1} ; uv(λ_{max} , nm (ϵ), CH_2Cl_2) 284 (13200), 449 (2680); ms (m/z): Exact mass: Calcd for $\text{C}_{13}\text{H}_9\text{NO}_4$: 243.0532. Found: 243.0528.
8. N. J. Turro and D. S. Weiss, *J. Am. Chem. Soc.*, 1968, **90**, 2185.

9. The evidence for existence of hydroquinone of 2 was confirmed by the 400 MHz ^1H Nmr spectrum of the photoreaction mixture in benzene- d_6 .

Received, 3rd February, 1993