


A NEW POTENTIAL ACYL IMINIUM ION FOR THE ASYMMETRIC SYNTHESIS OF PIPERIDINE DERIVATIVES

Jacques Royer and Henri-Philippe Husson*

Institut de Chimie des Substances Naturelles . CNRS, 1 avenue de la Terrasse,
91198 Gif-sur-Yvette Cedex, France


Abstract - Epimeric oxazolopiperidones (**1a**) and (**1b**) were prepared independently, both from *R*-(-)-phenylglycinol as starting material. They represent chiral potential iminium ions.

Acyl iminium ions are known as strong electrophiles widely used in C-C bond formation α to nitrogen.¹ The preparation, reactivity and application of these derivatives in the synthesis of nitrogen containing compounds have been thoroughly studied, particularly by Speckamp and co-workers.¹ In the course of our research of new starting materials for the asymmetric synthesis of piperidine derivatives,² we designed compound (**1**) as a chiral potential acyl iminium ion. Chiral bicyclic lactams (e.g. γ -lactams of type **2**), analogous to **1** have already been reported by Meyers' group for the asymmetric synthesis of quaternary carbon compounds.³ More recently, Naito and Ninomiya⁴ have described the synthesis and use of acyl lactam (**3**) for the synthesis of simple piperidine alkaloids while Meyers⁵ has reported the usefulness of lactam (**2**) for the synthesis of pyrrolidine alkaloids. These reports prompt us to disclose our preliminary results in the synthesis of **1**.




We first planned to synthesize compound (**1**) starting from the chiral 2-cyano-6-oxazolopiperidine (**4**) which was easily prepared in one step by condensation of *R*-(-)-phenylglycinol, glutaraldehyde and KCN² (Scheme 1). The

O₂ oxidation⁶ of the anion of **4** (generated by LDA at -78°C in THF) easily gave the expected lactam (**1a**) in a 46% yield (73% based on recovered starting material)). It was interesting to notice that this reaction gave lactam (**1a**)⁷ as a unique stereoisomer. The stereochemistry of oxazolopiperidone (**1a**) was difficult to be proved by nmr and it was expected according to its preparation that the configuration at the oxazolidine center of **1a** remained the same as that of the starting material. It then appeared that the relative stereochemistry of **1a** was reverse as that one found by Meyers for similar lactams in both pyrrolidine and piperidine series.³ To confirm this assumption, we decided to synthesize lactam (**1**) through the reduction of imide (**5**) following the procedure described by Speckamp.⁸ Some difficulties were encountered in the preparation of imide (**5**); attempts to condense glutaric anhydride with *R*-(-)-phenylglycinol, produced uncyclized products or polymeric materials under a variety of experimental conditions.⁹ Good results were finally obtained using acetyl chloride as a dehydrating reagent and acylated imide (**6**) was obtained in 65% yield. The NaBH₄ reduction of **6** in EtOH with added dilute HCl gave a mixture of epimeric ethoxy- and hydroxylactams (**7**) which upon treatment with 1M HCl in MeOH (15h, room temperature) gave rise to the clean formation of **1b** in 50% yield from **6**. (Scheme 1)



Scheme 1

Another approach was also tested for the preparation of **1**. Imide (**8**) was easily prepared (95% yield) from *R*-(-)-ethyl phenylglycinate and glutaric anhydride in the presence of AcCl and was reduced (NaBH₄, added HCl) to a mixture of **1b** (37%) and ethoxylactam (**9**) (19%) (Scheme 2)



Scheme 2

The stereochemical assignment of each isomer (**1a**) and (**1b**) was based on the stereochemical course of their formation: Lactam (**1a**) possesses the configuration of the starting material (**4**), while lactam (**1b**) has the configuration of the more stable oxazolidine, as observed by Meyers.³ The nmr data showed that the oxazolidine proton H-2 was shifted to lower field in **1b** (δ H-2: 5.02 ppm) than in **1a** (δ H-2: 4.82 ppm) due to the anisotropic effect of the phenyl group.

Oxazolopiperidinones (**1a**) and (**1b**) represent an extension of the chemistry of the 2-cyano-6-oxazolopiperidine system previously studied in our laboratory.²

REFERENCES AND NOTES

1. H. Hiemstra and W. N. Speckamp, *Tetrahedron*, 1985, **41**, 4367
2. L. Guerrier, J. Royer, D. S. Grierson, and H.-P. Husson, *J. Am. Chem. Soc.*, 1983, **105**, 7754; M. Bonin, D. S. Grierson, J. Royer, and H.-P. Husson, *Org. Syntheses*, 1992, **70**, 54 and references cited therein.
3. a) A. I. Meyers and K.T. Wanner, *Tetrahedron Lett.*, 1985, **26**, 2047; b) A. I. Meyers and B. A. Lefker, *Tetrahedron*, 1987, **43**, 5663; c) A. I. Meyers, R. Hanreich, and K.T. Wanner, *J. Am. Chem. Soc.*, 1985, **107**, 7776; d) A. I. Meyers, B. A. Lefker, T. J. Sowin, and L. J. Westrum, *J. Org. Chem.*, 1989, **54**, 4243.
4. T. Kiguchi, Y. Nakazono, S. Kotera, I. Ninomiya, and T. Naito, *Heterocycles*, 1990, **31**, 1525.
5. A. I. Meyers and L. E. Burgess, *J. Org. Chem.*, 1991, **56**, 2294.
6. Only one related reaction has been reported: D. S. Grierson, M. Urrea, and H.-P. Husson, *Heterocycles*, 1985, **23**, 2493.; for similar oxidation of nitriles, see: H. Yamanaka and S. Ohba, *Heterocycles*, 1990, **31**, 895; R. W. Freerksen, S. J. Selikson, and R. R. Wroble, *J. Org. Chem.*, 1983, **48**, 4087.
7. **1a**: mp 71°C (ether); $[\alpha]_D -51^\circ$ (CH_2Cl_2 , c 2.2); ^1H nmr (CDCl_3 , 400 MHz): 1.75 (m, 2H), 1.97 (m, 1H), 2.35 (m, 3H), 3.98 (d, $J=9$ Hz, 1H), 4.12 (dd, $J=9, 7$ Hz, 1H), 4.82 (dd, $J=9.5, 3.5$ Hz, 1H), 4.90 (d,

$J = 7$ Hz, 1H), 7.30 (m, 5H); ir (neat): 1640 cm^{-1} ; ms (EI) m/z : 217(M^+ , 70), 187(29), 159(20), 148(23), 120(39), 104(100); Hrms calcd for $C_{13}H_{15}NO_2$: 217.1103, found: 217.1109.

1b: oil; $[\alpha]_D -88^\circ$ (CH_2Cl_2 , c 0.6); 1H nmr ($CDCl_3$, 400 MHz): 1.50-2.05 ppm (m, 4H), 2.35 (dt, $J=18$, 6 Hz, 1H), 2.52 (dd, $J=18$, 6 Hz, 1H), 3.75 (dd, $J=8$, 7.5 Hz, 1H), 4.51 (dd, $J=8$, 7.5 Hz, 1H), 5.02 (dd, $J=8$, 4 Hz, 1H), 5.25 (t, $J=7.5$ Hz, 1H), 7.25 (m, 5H); ^{13}C nmr ($CDCl_3$, 50 MHz): 17.1, 28.5, 31.4, 58.2, 72.5, 88.9, 126.3, 127.8, 129.0, 139.9, 169.0; ir (neat): 1670 cm^{-1} ; ms (EI) m/z : 217(M^+ , 100), 187(26), 120(48), 104(79), 91(27); Hrms calcd for $C_{13}H_{15}NO_2$: 217.1103, found: 217.1108.

8. J. C. Hubert, J. B. P. A. Wijnberg, and W.N. Speckamp, *Tetrahedron*, 1975, **31**, 1437.

9. Meyers reported (ref. 3c) that condensation of *S*-valinol with glutaric anhydride gave only a moderate yield of maleimide (22%) and that the reaction was accompanied by extensive polymerization.

Received, 15th February, 1993