

ON THE DECOMPOSITION OF 1,2,4-TRIOXAN-5-ONES

Charles W. Jefford* and M. Graça H. Vicente

Department of Organic Chemistry, University of Geneva
1211 Geneva 4, Switzerland


Abstract - The thermal and chemical properties of six 3,6-substituted 1,2,4-trioxan-5-ones were studied by flash-vacuum thermolysis (FVT) and by treatment with electron-transfer agents. The 1,2,4-trioxan-5-ones exhibited unusual stability both chemically and thermally, but eventually underwent decomposition with loss of carbon dioxide yielding the carbonyl components.

Chemists have believed that the 1,2,4-trioxan-5-one ring is intrinsically unstable and that it would decompose spontaneously to carbon dioxide and its carbonyl constituents. Typically, the decarboxylation of α -hydroperoxy carboxylic acids (**1**) in the presence of the acetal of dimethylformamide¹ to give aldehydes and ketones (**3**) was thought to proceed through the intermediacy of a frangible 1,2,4-trioxan-5-one (**2**) (Scheme 1). Similarly, the photo-oxygenation² of α -pyrones (**4**) gave the bridged bicyclic trioxanones (**5**) which, however, lost carbon dioxide almost immediately at room temperature to give the 1,2-diacylethylenes (**6**) (Scheme 2). Besides these examples, few 1,2,4-trioxan-5-ones have been reported.³ However, as part of a program for discovering 1,2,4-trioxanes of anti-malarial potential,⁴ we recently devised an operationally simple method for preparing 3,3,6,6-tetra-substituted 1,2,4-trioxan-5-ones⁵ and were surprised to find that they were sufficiently stable to be characterized by X-ray.⁶ Consequently, we now present our findings on some of their chemical and physical properties, in particular on the behavior of representative members (**7-12**) towards heat and electron transfer agents (Scheme 3).


Submission of all of them to a first qualitative thermal test showed them to be more stable than expected. Injection onto a capillary column in a gas chromatograph at a temperature of less than 180 °C simply revealed the chromatogram characteristic of each particular trioxanone.⁷ However, raising the temperature to 210 °C resulted in decomposition to give the corresponding carbonyl products, *e.g.* cyclohexanone, 2-adamantanone, pivalaldehyde, or 1-adamantylcarboxaldehyde.

In a further experiment, heating **7-10** in boiling decalin (bp 189 °C) under a gentle stream of argon led to complete decomposition in 2-3 h, whereas **11** and **12** under the same conditions required 3-5 h. Passage of the ef-


Scheme 1




Scheme 2



Scheme 3



Scheme 4




fluent gas through an aqueous solution of barium hydroxide caused precipitation of barium carbonate in amounts indicative of yields of 80-100% of carbon dioxide.⁸ Flash-vacuum thermolysis (FVT)⁹ confirmed the thermal stability. The *t*-butyl and the spirocyclic adamantane derivatives (**7**) and (**11**) survived temperatures of 300 °C, but progressively fragmented into carbon dioxide, pivalaldehyde (**13**), cyclohexanone (**14**) and 2-adamantanone (**15**) at temperatures of 350-400 °C (Scheme 4).¹⁰ Trioxanones (**8-10**) showed similar FVT behavior, whereas **12** volatilized only difficultly owing to its high melting point (216 °C).

Next, the stability of the trioxanones (**7-12**) towards electron transfer agents was assayed.¹¹ It is well known that the peroxide bond is susceptible to scission, especially by electron donors.¹² Unexpectedly, trioxanones (**7-12**) were found to be quite resistant to the action of sodium azide, sodium iodide or triethylamine in diethyl ketone at 25 °C. In the boiling solvent (bp 103 °C), trioxanones (**7-10**) decomposed in 1 h, whereas when a higher boiling solvent, tetrachloroethylene (bp 122 °C), was used, decomposition was over in 10-20 min. In contrast, trioxanones (**11**) and (**12**) displayed greater stability to the aforementioned electron transfer agents and required 24-31 h in boiling diethyl ketone or tetrachloroethylene to undergo complete fragmentation. In the absence of the agents, exposure to the boiling solvents for more than 48 h was necessary for complete fragmentation.


Metal salts and complexes known to have an affinity for the peroxide bond, such as $\text{Fe}(\text{ClO}_4)_2$, $\text{Fe}(p\text{-TsO})_2$,¹³ $\text{Pd}(\text{PPh}_3)_4$,¹⁴ $\text{Co}(\text{II})m\text{-TPP}$,¹⁵ and $\text{Zn}(\text{II})m\text{-TPP}$ ¹⁶ at concentrations of 5-6% molar equivalent in either tetrachloroethylene, THF (bp 66 °C) or acetonitrile (bp 82 °C) had a similar effect on **7-12**. Reaction times for complete decomposition were usually 1-14 h.

The aforementioned agents (D) initiated fragmentation by a mechanism that is best explained by electron exchange (Scheme 5).¹⁷ Taking trioxanone (**11**) as an illustration, the electron donor, D, gives an electron to the oxygen-oxygen σ^* -bond to afford the radical anion (**16**). Subsequently, cleavage can occur giving either the adamantyl- or cyclohexanonyl radicals (**17**) or (**18**). Restitution of the electron by **17** or **18** to the donor radical cation (D^+) creates an excited donor (D^*) and liberates adamantanone (**15**) and cyclohexanone (**14**) respectively. No luminescence was visible to the naked eye. However, heating the trioxanone alone in 1-phenyldecane as solvent (bp 294 °C) above 200 °C resulted in light emission as detected by a photo-multiplier tube.¹⁸

Although triethylamine decomposes **11** and **12** by electron exchange, a different mechanism was followed by trioxanones (**7-10**). In boiling tetrachloroethylene after 50 min. they were effectively deprotonated at the C6



Scheme 6



position with excision of adamantanone or cyclohexanone¹⁹ to produce the corresponding α -keto carboxylic acids (**19**, **20**) in 100% yield (Scheme 6).

The conclusion is that 1,2,4-trioxanones (**7-12**) are unusually thermally stable. Their decomposition is facilitated by electron transfer agents but high temperatures (100-120 °C) are required. These trioxanones stand in contrast to the bridged bicyclic analogues (**5**) which probably owe their inherent fragility to the endocyclic double bond. Experiments on the chemistry and physical properties of **7-12** are under way and the results will be reported in due course.

Acknowledgments. We are grateful for support of this work by the Swiss National Science Foundation (grant No 20-32'166.91).

REFERENCES AND NOTES

1. H.H. Wasserman and B.H. Lipshutz, *Tetrahedron Lett.*, **1975**, 4611.
2. W. Adam and I. Erden, *Angew. Chem., Int. Ed. Engl.*, **1978**, 17, 211; *J. Am. Chem. Soc.*, **1979**, 101, 5692.
3. J.M. Holland and D.W. Jones, *J. Chem. Soc. C*, **1970**, 530; J.P. Smith and G.B. Schuster, *J. Am. Chem. Soc.*, **1978**, 100, 2564.
4. Eighth Programme Report, UNDP/World Bank/WHO, Special Programme for Research and Training in Tropical Medicine, **1987**, p. 24.

5. C.W. Jefford, J. Currie, G.D. Richardson, and J.C. Rossier, *Helv. Chim. Acta*, **1991**, *74*, 1239.
6. C.W. Jefford, G.D. Richardson, J.C. Rossier, and G. Bernardinelli, *Acta Cryst.*, **1991**, *C47*, 388.
7. Analytical glc was performed on a Hewlett Packard model 5880 A instrument equipped with a flame ionization detector and a Hewlett Packard Ultra 1 capillary column (stationary phase: cross-linked PhMe silicone, length x internal diameter x film thickness = 25 m x 0.22 mm x 0.11 μ m), carrier gas: N₂.
8. The yields of carbon dioxide obtained were calculated from the amount of barium carbonate recovered.
9. U.E. Wiersum, *Aldrichimica Acta*, **1984**, *17*, 31.
10. The products were identified by both gc and ¹H-nmr. The yields were determined on the basis of the ¹H-nmr spectra by using an internal standard (1,3,5-tribromobenzene). Low yields of pivalaldehyde were probably due to rearrangement, alternative reaction pathways, or loss at the high temperatures and low pressures used in these experiments.
11. See, for example, a) F.D. Greene and W.W. Rees, *J. Am. Chem. Soc.*, **1958**, *80*, 3432; b) J.K. Kochi and H.E. Mains, *J. Org. Chem.*, **1965**, *30*, 1862; c) R.M. Wilson, S.W. Wunderly, T.F. Walsh, A.K. Musser, R. Outcalt, F. Geiser, S.K. Gee, W. Brabender, L. Yerino, Jr., T.T. Conrad, and G.A. Tharp, *J. Am. Chem. Soc.*, **1982**, *104*, 4429.
12. a) "Organic Peroxides, Their Chemistry, Decomposition and Role in Polymerization", A.V. Tobolsky and R.B. Mesrobian, Interscience Publishers, Inc. New York, 1954; b) W. Herz, R.C. Ligon, J.A. Turner and J.F. Blount, *J. Org. Chem.*, **1977**, *42*, 1885.
13. M. Suzuki, Y. Oda, and R. Noyori, *Tetrahedron Lett.*, **1981**, *22*, 4413.
14. a) J.D. Boyd, C.S. Foote, and D.K. Imagawa, *J. Am. Chem. Soc.*, **1980**, *102*, 3641; b) M. Balci and Y. Sütbeyaz, *Tetrahedron Lett.*, **1983**, *24*, 4135; d) N. Akbulut, A. Menzek, and M. Balci, *Tetrahedron Lett.*, **1987**, *28*, 1689; e) Y. Sütbeyaz, H. Seçen, and M. Balci, *J. Org. Chem.*, **1988**, *53*, 2312; f) K.E. O'Shea and C.S. Foote, *J. Org. Chem.*, **1989**, *54*, 3475.
15. Y. Ito, M. Tone, H. Yokoya, T. Matsuura, and G.B. Schuster, *J. Org. Chem.*, **1986**, *51*, 2240.
16. J.-Y. Koo and G.B. Schuster, *J. Am. Chem. Soc.*, **1977**, *99*, 6107; b) **1978**, *100*, 4496.
17. N. Komblum and H.E. DeLaMare, *J. Am. Chem. Soc.*, **1951**, *73*, 880; b) W. Adam, M. Balci, and J. Rivera, *Synthesis*, **1979**, 807.
18. C.W. Jefford, M.G.H. Vicente, D. Lovy, H. Hagemann, and H. Bill, unpublished work.
19. C.W. Jefford, J.C. Rossier, and J. Boukouvalas, *J. Chem. Soc., Chem. Commun.*, **1986**, 1701.

Received, 29th January, 1993