



TOTAL SYNTHESIS OF (±)-*cis*-, *trans*-CEMBRANOLIDES AND (±)-SARCOPHYTONIN B FROM GERANYLGERANIOL¹

Kiyoshi Nishitani, Toshihiko Konomi, Kiyonobu Okada, and Koji Yamakawa*

Faculty of Pharmaceutical Sciences, Science University of Tokyo,
Ichigaya-funagawara, Shinjuku-ku, Tokyo 162, Japan


Abstract - (±)-*cis*-Cembranolide (**I**) was synthesized *via* Cr(II) mediated intramolecular macrocyclization of ω-formyl-β-methoxycarbonylallyl halides (**6a** and **b**), which were prepared starting from geranylgeraniol. The stereoselective synthesis of (±)-*trans*-cembranolide (**II**) was achieved by lactonization of 14-membered *cis*-intermediate (**7**) with inversion of the stereochemistry at C-2. (±)-Sarcophytonin B (**III**) was also synthesized from (±)-*cis*-cembranolide (**I**) *via* isomerization of the *exo*-cyclic double bond.

Many cembranolides, 14-membered diterpenoid lactones, have been found as marine natural products,² and some of them possess interesting physiological properties.³ Some efforts have been made to synthesize cembranolides in recent years.⁴ The unnamed *cis*- and *trans*-cembranolides (**I** and **II**) were isolated from *Sinularia mayi* by Uchio *et al.*,⁵ and also from *Lobophytum michaelae* by Coll *et al.*⁶ Sarcophytonin B (**III**) was isolated from *Sarcophyton* sp. by Kobayashi *et al.*⁷ Several syntheses of the cembranolides (**I** and **II**) had been reported.⁸




We recently described a facile synthesis of (±)-**I** *via* Cr(II) mediated macrocyclization of the ω-formyl-β-methoxycarbonylallyl halide (**6**), derived from geraniol, and subsequent lactonization reaction.⁹ We now describe another more facile synthesis of (±)-**I** starting from geranylgeraniol. We also report the synthesis of (±)-*trans*-cembranolide (**II**), and (±)-sarcophytonin B (**III**) using this intramolecular macrocyclization method.

Preparation of the ω -formylallyl halide (**6**) starting from geranylgeraniol was as follows. The regioselective hydroxy bromination of the terminal double bond of the THP ether of geranylgeraniol (**1**) was performed by the method of van Tamelen *et al.* (NBS in 40% H₂O-THF).¹⁰ The bromohydrin (**2**) was treated with K₂CO₃ followed by HClO₄ to give the diol (**3**), which was then converted into the aldehyde (**4**) with lead tetraacetate in good yield. The aldehyde was identified by the comparison with the authentic sample synthesized from geraniol.⁹ Conversion of the aldehyde into ω -formylallyl halides (**6a** and **6b**) had already been reported by us.^{9,11} Macrocyclization of the aldehydes (**6a** and **6b**) selectively affords the *cis*-cyclized product (**7**) in good yields, and subsequent lactonization with *p*-TsOH gave ($\pm$)-*cis*-cembranolid (**I**).⁹ Treatment of the methyl ester (**7**) with NaH in THF¹² gave the carboxylic acid (**8**), which was then treated with *N,N*-dimethylformamide dineopentylacetal *via* Walden inversion^{12,13} to give ($\pm$)-*trans*-cembranolid (**II**), mp 53.5-54.5 °C in 70% yield. The ¹H- and ¹³C-nmr spectra were coincident with those of the natural (-)-*trans*-cembranolid.¹⁴



Scheme 1

a) NBS/THF-H₂O (61%), b) 1) K₂CO₃/MeOH, 2) HClO₄/THF-H₂O (86%), c) Pb(OAc)₄, K₂CO₃/benzene (94%), d) LDA, THF, -78 °C (56%, conversion 70%), e) See reference 8 and note 10.



Scheme 2

a) 5.5 eq CrCl₂/DMF (0.02M), room temperature (79-81%), b) *p*-TsOH/benzene, room temperature 10 min (87%), c) 1) NaH/THF, room temperature, 2) Me₂NCH(O-neopentyl)₂/toluene, reflux (70%), d) RhCl₃/EtOH-H₂O, reflux (66%)

Isomerization of the *exo*-cyclic double bond of *cis*-cembranolid (I) into the *endo*-cyclic one was effected with rhodium chloride.¹⁵ An aqueous ethanol solution of I and RhCl₃ was heated for 7 h to afford (±)-sarcophytonin B (III) in 66 % yield. The ¹H-nmr spectrum and other spectral properties were identical with those of the natural (+)-sarcophytonin B.¹⁶

ACKNOWLEDGEMENTS

We thank Miss N. Sawabe and Mrs. F. Hasegawa of this laboratory for nmr and mass spectral measurements, and Mr. H. Igarashi for elemental analysis. Support for this work from the Ministry of Education, Science and Culture of Japan through Grants-in-Aid for Scientific Research is gratefully acknowledged. Geranylgeraniol was kindly provided by Kuraray Co., Ltd. and Takasago perfumery Co., Ltd.

REFERENCES AND NOTES

1. This paper forms "Studies on the Terpenoids and Related Alicyclic Compounds XLV". Part XLIV: reference 8.
2. A. J. Weinheimer, C. W. J. Chang, and J. A. Matson, *Fortschr. Chem. Org. Naturst.*, **1979**, 36, 286.
3. For examples; Y. Uchio, J. Toyota, H. Nozaki, M. Nakayama, Y. Nishizono, and T. Hase, *Tetrahedron Lett.*, **1981**, 22, 4089; A. J. Weinheimer, J. A. Matson, M. B. Hossain, and D. Helm, *Tetrahedron Lett.*, **1977**, 2923; I. Neeman, L. Fishelson, and Y. Kashman, *Toxicon*, **1974**, 12, 593.
4. Reviews see; M. A. Tius, *Chem. Rev.*, **1988**, 88, 719 and references therein. Recent report; M. A. Tius and N. K. Reddy, *Tetrahedron Lett.*, **1991**, 32, 3605.
5. Y. Uchio, S. Eguchi, M. Nakayama, and T. Hase, *Chem. Lett.*, **1982**, 277.
6. J. C. Coll, S. J. Mitchel and G. J. Stokie, *Aust. J. Chem.*, **1977**, 30, 1859.
7. M. Kobayashi and T. Hirase, *Chem. Pharm. Bull.*, **1990**, 38, 2442.
8. M. Aoki, T. Uyehara, and T. Kato, *Chem. Lett.*, **1984**, 69; J. A. Marshall and S. L. Crooks, *Tetrahedron Lett.*, **1987**, 28, 5081; J. A. Marshall, S. L. Crooks, and B. S. DeHoff, *J. Org. Chem.*, **1988**, 53, 1616; J. A. Marshall and W. Y. Gung, *Tetrahedron Lett.*, **1988**, 29, 1657; K. Kodama, T. Takahashi, and S. Itô, *Tetrahedron Lett.*, **1982**, 23, 5175.
9. K. Nishitani, T. Konomi, Y. Mimaki, T. Tsunoda, and K. Yamakawa, *Heterocycles*, **1993**, 36, 1957.
10. E. E. van Tamelen and T. J. Curphey, *Tetrahedron Lett.*, **1962**, 121; E. J. Corey and W. -C. Shieh, *Tetrahedron Lett.*, **1992**, 33, 6435.
11. The allylic bromide (6b) was synthesized by the same methods as described for 6a except bromination step (CBr₄ was used instead of CCl₄).
12. C. Kuroda, S. Shimizu, and J. Y. Satoh, *J. Chem. Soc., Perkin Trans. I*, **1990**, 519.
13. H. Vorbruggen and K. Krolikiewicz, *Angew. Chem., Int. Ed. Engl.*, **1977**, 16, 876.
14. Spectra of the natural *cis*- and *trans*-cembranolides were kindly provided by Prof. M. Nakayama and Prof. Y. Uchio to whom we are grateful.
15. P. A. Grieco, M. Nishizawa, N. Marinovic, and W. J. Ehmann, *J. Am. Chem. Soc.*, **1976**, 98, 7102.
16. We thank Dr. M. Kobayashi for kind providing of ¹H-, ¹³C-nmr, ir and mass spectra of (+)-sarcophytonin B.

Received, 16th September, 1993