

REACTION OF 1-HYDROXYPHTHALIMIDE DERIVATIVES WITH ALUMINUM CHLORIDE IN BENZENE

Kensaku Uto, Takeshi Sakamoto, Keita Matsumoto,[†] and Yasuo Kikugawa*

Faculty of Pharmaceutical Sciences, Josai University, 1-1 Keyakidai, Sakado, Saitama 350-02, Japan; Taisho Pharmaceutical Co., Ltd., 1 Yoshinocho, Ohmiya, Saitama 330, Japan[†]

Abstract --- The reaction of 1-hydroxyphthalimide derivatives with AlCl₃ in benzene has been investigated. From 1-hydroxyphthalimide (**1a**) 2-hydroxy-3,3-diphenyl-2,3-dihydroisoindol-1-one (**2**) and 1,1-diphenyl-1*H*-benzo[*d*][1,2]oxazin-4-one (**3**) are obtained by initial attack of benzene on the imide carbonyl, assisted by the neighboring oxygen atom. Heating **2** with Lewis acid (AlCl₃) in benzene results in ring expansion to afford **3**, while heating **3** with protic acid (H₂SO₄) leads to ring contraction to give **2**. From *O*-(2-phenethyl)- and *O*-benzoyl derivatives (**1b** and **c**), 1,2-diphenylethane (**4**) and benzophenone (**5**) are obtained, respectively, by the heterolytic cleavage of an C-O bond.

Previously we reported the AlCl₃-mediated heterolytic cleavage of the N-O bond of *N*-methoxy-*N*-phenylamides.¹ This cleavage reaction provides a new source of *N*-acyl-*N*-arylnitrenium ions² and leads to the nucleophilic intramolecular migration of the methoxy group from the nitrogen to the *ortho* position of the benzene ring.

In an extension of this work, we have investigated the reaction of 1-hydroxyphthalimide derivatives with AlCl₃ in benzene, in the expectation that AlCl₃-mediated cleavage of

the N-O bond might give a diacylnitrenium ion³ which will be trapped by solvent benzene. However, the results are different from expectation.

When the progress of the reaction of **1a** with AlCl_3 in benzene was monitored by tlc, production of two new compounds was observed: compound (**2**), having a lower R_f value, was gradually converted to **3** (which has a higher R_f value) and disappeared at the end of the reaction (7 h reflux). Isolation of **2** and **3** by silica gel column chromatography was performed and structures of them were determined unambiguously by spectral and X-ray analyses.

Scheme 1. Reaction of 1-hydroxyphthalimide derivatives with AlCl_3 in C_6H_6





Table. ^{13}C -Nmr chemical shifts (δ ppm) of **2** and **3**^{a)}

position	2		3	
a	164.20		156.57	
b	76.12		96.23	
c	139.48		141.67	
d	unassigned		128.07	
i	146.52		146.65	
e f g h j k l	123.63	123.97	122.05	123.88
	128.21	128.26	127.22	128.37
	128.37	132.15	129.04	131.0


^{a)}The data were taken in CDCl_3 at 67.8 MHz.

The ^{13}C -nmr spectrum of **2** shows 8 peaks for aromatic carbons, indicating that two peaks are overlapped. X-ray crystallographic analysis revealed the structure of **2** to be

2-hydroxy-3,3-diphenyl-2,3-dihydroisindol-1-one. The structure of **3** was determined to be 1,1-diphenyl-1*H*-benzo[*d*][1,2]oxazin-4-one by spectral analyses. ^{13}C -Nmr data are presented in the Table.


Heating **2** for 7 h with AlCl_3 in benzene resulted in ring expansion to afford **3** in 65% yield. On the other hand, heating **3** for 1 h in 30% H_2SO_4 -dioxane leads to ring contraction to give **2** in 93% yield. In the reverse reaction, protonation on the oxygen atom induces cleavage of the C-O bond and subsequent recombination forms a C-N bond, as shown in Scheme 2.

Scheme 2. Ring expansion and ring contraction



An acyclic *N*-alkoxyimide, *N*-methoxy-*N*-(4-methylbenzoyl)-4-nitrobenzamide (**6**), reacted with AlCl_3 in benzene (room temperature, 40 min) to give 4-nitrobenzophenone


Scheme 3. Reaction of 1-hydroxy- and 1-methoxyamide derivatives with AlCl_3 in C_6H_6



(**7**) (90%) and *N*-methoxy-4-methylbenzamide (**8**) (73%) (Scheme 3). However, phthalimide (**9a**) and 1-phenylphthalimide (**9b**) do not react with AlCl_3 under stronger reaction conditions and the starting materials were recovered quantitatively. *N*-Hydroxy-*N*-methylbenzamide (**10**) reacted with AlCl_3 in benzene (7 h reflux) to give

benzophenone (**5**) (4%) and starting material (**10**) (75%). These results indicate that the 1-hydroxy group of **1a** activates the imide carbonyl for Friedel-Crafts type reactions. The assumed mechanism is presented in Scheme 4.

Scheme 4. Assumed mechanism



When **1b** was submitted to similar reaction conditions, heterolytic cleavage of the C-O bond and subsequent Friedel-Crafts type phenylation occurred to give 1,2-diphenylethane (**4**) (86%) and **1a** (76%) (path b). Similarly, benzophenone (**5**) (81%) and **1a** (74%) were obtained from the *O*-benzoyl compound (**1c**) (path c).

EXPERIMENTAL

All the melting points were determined with a Yanagimoto hot-stage melting point apparatus and are uncorrected. ^1H Nmr spectra were measured at 60 MHz on a JEOL JNM-PNX60SI or at 270 MHz on a JEOL JNM-EX270 spectrometer with

tetramethylsilane (Me_4Si) as an internal reference and CDCl_3 as the solvent, unless otherwise noted. ^1H Nmr spectral data are reported in parts per million (δ) relative to Me_4Si . Infrared spectra were recorded on a JASCO IR 810 spectrophotometer. Mass spectra were obtained with a JEOL JMX-DX 300 spectrometer with a direct inlet system at 70 eV. Elemental analyses were performed in the Microanalytical Laboratory of this University. Compounds (**1a**, **4**, **5**, **7**, **8**, and **9a**) were purchased from Tokyo Kasei Kogyo Co. Compound (**1c**) and (**9b**) were prepared by the literature methods.^{4,5} Compound (**1b**) was prepared from the potassium salt of **1a** and 2-bromoethylbenzene in *N,N*-dimethylformamide in 71% yield, mp 93-94 °C (from benzene-hexane). ^1H Nmr δ 3.13 (2H, t, $J=6.9$, OCH_2CH_2), 4.40 (2H, t, $J=6.9$, OCH_2CH_2), 7.22 (5H, s, ArH), 7.73 (4H, s, ArH); $\nu_{\text{max}}(\text{KBr})/\text{cm}^{-1}$ 1735; m/z 163 (M^+-104 , 12.9%), 105 (100). Anal. Calcd for $\text{C}_{16}\text{H}_{13}\text{NO}_3$: C, 71.90; H, 4.90; N, 5.24. Found C, 72.12; H, 5.09; N, 5.20. Compound (**6**) was prepared by the Schotten-Baumann reaction, using 4-nitrobenzoyl chloride and **8**, mp 134-135 °C (benzene-hexane). ^1H Nmr δ 2.35 (3H, s, CH_3), 3.92 (3H, s, OCH_3), 7.12 (2H, d, $J=8.0$, ArH), 7.55 (2H, d, $J=8.0$, ArH), 8.28 (4H, s, ArH); $\nu_{\text{max}}(\text{KBr})/\text{cm}^{-1}$ 1740; m/z 314 (M^+ , 3.36), 119 (100). Anal. Calcd for $\text{C}_{16}\text{H}_{14}\text{N}_2\text{O}_5$: C, 61.14; H, 4.49; N, 8.91. Found C, 61.10; H, 4.63; N, 8.97.

2-Hydroxy-3,3-diphenyl-2,3-dihydroisoindol-1-one (2) and 1,1-diphenyl-1H-benzo[d][1,2]oxazin-4-one (3); path a

To **1a** (200 mg, 1.23 mmol) in benzene (8 ml) was added AlCl_3 (817 mg, 6.13 mmol) with cooling. After refluxing the reaction mixture for 1.5 h, 10% HCl (10 ml) was added with cooling. The aqueous layer was extracted with ethyl acetate (20 ml x 2), and the combined organic layer was washed with brine (30 ml), dried over Na_2SO_4 , and concentrated. The crude products were chromatographed on a column of silica gel. First elution with benzene-ethyl acetate (2:1) afforded **3** (232 mg, 63%), mp 193-195 °C (from benzene-hexane). ^1H Nmr δ 7.11-7.79 (14H, m, ArH), 8.12 (1H, br s, NH); $\nu_{\text{max}}(\text{KBr})/\text{cm}^{-1}$ 3325, 1675; m/z 301 (M^+ , 16.4), 208 (100). Anal. Calcd for $\text{C}_{20}\text{H}_{15}\text{NO}_2$: C, 79.72; H, 5.02; N, 4.65. Found C, 79.80; H, 5.19; N, 4.57. Further elution with the same solvent mixture afforded **2** (65 mg, 18%), mp 206-208 °C (AcOEt). ^1H Nmr δ 7.00-7.67 (14H, m, ArH); $\nu_{\text{max}}(\text{KBr})/\text{cm}^{-1}$ 3100, 1685; m/z 301

(M+, 14.6), 208 (100). Anal. Calcd for $C_{20}H_{15}NO_2$: C, 79.72; H, 5.02; N, 4.65. Found C, 79.68; H, 5.28; N, 4.92. Compound (**2**) was gradually converted to **3** under the reaction conditions and **3** was obtained in 91% yield when the reaction time was prolonged to 7 h.

Synthesis of **3** from **2**


A mixture of **3** (213 mg, 0.708 mmol), 30% H_2SO_4 (10 ml), and dioxane (10 ml) was refluxed for 1 h. After the reaction, H_2O (10 ml) was added and the solution was extracted with ethyl acetate (20 ml x 2), and the combined organic layer was washed with brine (30 ml), dried over Na_2SO_4 , and concentrated. The residue was chromatographed on a column of silica gel with benzene-ethyl acetate (2:1) as an eluent to give **2** (198 mg, 93%), mp 205-208 °C.

X-Ray Analysis of **2**

#A12001-001 $C_{20}H_{15}NO_2$ F.W.=301, mp 206-208 °C (acetone- CH_2Cl_2)

Crystal data: $a=9.584(3)$ Å, $b=9.704(4)$ Å, $c=9.254(2)$ Å, $\alpha=117.64(2)^\circ$, $\beta=92.47(2)^\circ$, $\gamma=94.90(3)^\circ$, $V=756.3(4)$ Å³, triclinic, $P\bar{1}$, $Z=2$, $D_x=1.32$ g/cm³, $F(000)=316$, $\mu(Cu K\alpha)=6.00$ cm⁻¹.

The diffraction experiment was carried out using a colorless transparent single crystal with dimension of 0.40x0.50x0.30 mm. The diffractometer MXC18(Mac Science) was used with graphite-monochromated Cu $K\alpha$ radiation ($\lambda=1.5418$ Å). The unit cell dimensions were determined from angular setting of 20 reflections (2θ values in the range of 50-60°). 2495 unique reflections ($2\theta\leq 130^\circ$) were measured, of which 2465 with $F_o>0$ $\sigma(F_o)$ were considered as observed. Absorption corrections were applied to the data.⁷ The structure was solved by the direct method using SHELX586⁶ and the difference Fourier method. The refinement of atomic parameters was carried out using the full matrix least squares methods.⁷ Anisotropic thermal parameters were used for all non-H atoms, while only coordinates of H atoms were refined. The atomic scattering factors were taken from "International Tables for X-ray Crystallography".⁸ The final R value is 0.042 ($R_w = 0.061$).

Figure 1. Parallel view of **2****1,2-Diphenylethane (4); path b**

To **1b** (200 mg, 0.80 mmol) in benzene (8 ml) was added AlCl_3 (531 mg, 3.98 mmol) with cooling. After stirring the reaction mixture for 0.5 h at room temperature, 10% HCl (10 ml) was added with cooling. The aqueous layer was extracted with CH_2Cl_2 (30 ml x 2), and the combined organic layer was washed with brine (30 ml), dried over Na_2SO_4 , and concentrated. The crude products were chromatographed on a column of silica gel. First elution with benzene-ethyl acetate (2:1) afforded **4** (124 mg, 86%), mp 46-48 °C (lit.,⁹ mp 52 °C), and further elution with the same solvent mixture afforded **1a** (98 mg, 76%), mp 234-236 °C (benzene-AcOEt) (lit.,¹⁰ mp 230 °C).

Benzophenone (5), path c

To **1c** (200 mg, 0.75 mmol) in benzene (8 ml) was added AlCl_3 (499 mg, 3.98 mmol) with cooling. After stirring the reaction mixture for 0.5 h at room temperature, 10% HCl (10 ml) was added with cooling. The aqueous layer was extracted with ethyl acetate (30 ml x 2), and the combined organic layer was washed with brine (30 ml), dried over Na_2SO_4 , and concentrated. The crude products were chromatographed on a column of silica gel. First elution with benzene-ethyl acetate (5:1) afforded **5** (111 mg, 81%), the structure of which was determined by comparison of spectral data with those of an authentic sample. Further elution with the same solvent mixture afforded **1a** (90 mg, 74%), mp 235-237 °C (benzene-AcOEt).

REFERENCES

1. Y. Kikugawa and M. Shimada, *J. Chem. Soc., Chem. Commun.*, **1989**, 1450.
2. R. A. Abramovitch and R. Jeyaraman, in 'Azides and Nitrenes,' ed. E. F. V. Scriven, Academic Press, New York, 1984, pp. 297-357.
3. R. A. Abramovitch, J. M. Beckert, P. Chinnasamy, H. Xiaohua, W. Pennington, and A. R. V. Sanjivamurthy, *Heterocycles*, 1989, **28**, 623; J. I. G. Cadogan and A. G. Rowley, *J. Chem. Soc., Perkin Trans. 1*, **1975**, 1069.
4. E. Grochowski and J. Jurczak, *Synthesis*, **1992**, 277.
5. M. Onda and Y. Kunugi, *Yakugaku Zasshi*, 1958, **78**, 690; R. Pummerer and G. Dorfmueller, *Ber.*, 1912, **45**, 292.
6. SHELXS86: Program for the Solution of Crystal Structure; G. M. Sheldrick, Univ. of Göttingen, Germany, 1986.
7. C. Katayama, *Acta Cryst.*, 1986, **A42**, 19.
8. International Tables for X-ray Crystallography, Vol. IV, Birmingham, Kynoch Press, 1974.
9. E. Sakellarios and T. Kyrimis, *Ber.*, 1924, **57**, 322.
10. G. H. L. Nefkens and G. I. Tesser, *J. Am. Chem. Soc.*, 1961, **83**, 1263.

Received, 1st November, 1995