

TWO NEW BISACRIDONE ALKALOIDS FROM *CITRUS* PLANTS¹


Motoharu Ju-ichi,*^a Yuko Takemura,^a Noriko Nagareya,^a Mitsuo Omura,^b Chihiro Ito,^c and Hiroshi Furukawa^c

Faculty of Pharmaceutical Sciences, Mukogawa Women's University,^a Nishinomiya, Hyogo 663, Japan, Fruit Tree Research Station,^b Ministry of Agriculture, Forestry and Fisheries, Okitsu, Shimizu, Shizuoka 424-02, Japan, and Faculty of Pharmacy, Meijo University,^c Tempaku, Nagoya 468, Japan

Abstract — Two new binary acridone alkaloids named citbismine-D (1) and -E (2) were isolated from the roots of Marsh grapefruit (*Citrus paradisi*). The structures were elucidated by spectroscopic studies as 1 and 2. Citbismine-E (2) was also obtained from the roots of Hirado-Buntan (*C. grandis*).

Previous chemical studies in our laboratory of the constituents of *Citrus* plants (Rutaceae) have resulted in the isolation of many acridone alkaloids,² coumarins³ and acridone-coumarin dimers.⁴ Recently, we reported the isolation and structure elucidation of new type bisacridone alkaloid citbismine-A⁵ from the roots of Marsh grapefruit (*C. paradisi* Macf.) and Hirado-Buntan (*C. grandis* Osbeck f. Hirado). Further study of these plants resulted in the isolation of two additional bisacridone alkaloids named citbismine-D (1) and -E (2). This paper deals with the structure elucidation of these alkaloids.

Citbismine-D (1) was obtained from the roots of Marsh grapefruit as a pale yellow oil, $[\alpha]_D^{25} \pm 0.0$ (CHCl₃). The molecular formula C₄₀H₃₈N₂O₁₁ was obtained by HRms (m/z 722.2473). The ir (1620, 1595 cm⁻¹) and uv [224 (sh), 267, 298 (sh), 338, 387 nm] absorptions indicated the presence of a 9-acridone skeleton.⁶ The ¹H nmr spectrum had signals due to two chelated hydroxyl groups (δ 15.84, 14.82), two *ortho*-coupled aromatic protons [δ 7.86, 6.81 (each 1H, d, J = 8.5 Hz), 8.04, 7.05 (each 1H, d, J = 9.2 Hz)], a lone aromatic proton [δ 6.18 (1H, s)], and a 2,2-dimethylpyran ring [δ 6.58, 5.51 (each 1H, d, J = 9.8 Hz), 1.30 (6H, s)]. The lowest signals of aromatic proton at δ 7.86 and 8.04 were deshielded by the 9- and 9'-carbonyl group, and thus were assigned as signals of H-8 and H-8'. The presence of 2-hydroxyisopropyl substituted dihydrofuran moiety was assumed based on the signals at δ 5.70, 4.56 (each 1H, d, J = 5.5 Hz), 1.38, 1.43 (each 3H, s) and 4.05 (1H, br s). The characteristic high field signal at δ 2.91 (3H, s) was assigned to the 5-methoxyl group, shielded by another acridone nucleus⁷ and resonating at an unusually high field. The remaining signals at δ 3.77, 3.64 and 3.88 indicated the presence of methoxyl and *N*-methyl groups. The assignment of these



signals was facilitated by the following NOE experiments. Irradiation of the signal at δ 3.77 showed 9.5% increments of the signal at δ 5.70 (H-11), suggesting the presence of a 10-*N*-methyl group and angular orientation of the dihydrofuran ring. When the signal at δ 3.64 was irradiated, 9.8% increment was observed of the signal at δ 6.58, indicating the presence of a 10'-*N*-methyl group and angular orientation of the 2,2-dimethylpyran ring. Irradiation of the signal at δ 3.88 showed no increments of any aromatic protons, leading to the conclusion that this group is located at C-5'. On the basis of these results, the linkage of two acridone moieties was assigned between C-11 and C-2'. From the above results, the structure of citbismine-D was concluded to be 1.

Citbismine-E (2) was obtained from the roots of Marsh grapefruit and Hirado-Buntan as a pale yellow oil, $[\alpha]_D^{25} \pm 0^\circ$ (CHCl₃). The HRms gave a molecular ion at m/z 670.2160, consistent with the molecular formula of C₃₆H₃₄N₂O₁₁. The presence of a 9-acridone skeleton⁶ was assumed by ir (1620, 1595 cm⁻¹) and uv (221, 264, 275 (sh), 298 (sh), 335, 379 nm) absorptions. The ¹H-nmr spectrum showed similar signal patterns with that of 1: the signals of two chelated hydroxyl groups (δ 15.92, 14.78), two *ortho*-coupled [δ 8.14, 7.03 (each 1H, d, *J* = 9.2 Hz), 7.86, 6.80 (each 1H, d, *J* = 8.5 Hz)] aromatic protons and a 2-hydroxyisopropyl substituted dihydrofuran ring [δ 5.72, 4.54 (each 1H, d, *J* = 5.5 Hz), 3.9 (1H, br s), 1.43, 1.38 (each 3H, s)], and methoxyl and *N*-methyl [δ 4.02, 3.88, 3.78, 3.57 and 2.80 (each 3H, s)] groups. The differences were the presence of signals of one additional methoxyl group (δ 3.57) and a lone aromatic proton (δ 6.46) instead of the disappearance of the signals of the 2,2-dimethylpyran moiety. The location of methoxyl and *N*-methyl groups was elucidated by the following NOE experiments. Irradiation of the signals at δ 4.02 and 3.57 showed 12 and 11% enhancements of the signal at δ 6.46, respectively. The result led us to conclude that the signals at δ 4.02 and 3.57 were due to the 10' *N*-methyl and 3'-methoxyl group, respectively. Irradiation of the signal at δ 3.78 showed 8.7% increments of the signal at δ 5.72, providing evidence for the presence of the 10-*N*-methyl group. Irradiation of signals at δ 2.80 and 3.88 showed no

increments of aromatic protons, suggesting the location of these methoxyl groups to be C-5 and C-5', respectively. Thus, the structure of citbismine-E was concluded to be 2.

Both citbismine-D (1) and -E (2), in analogy with other citbismine, were obtained in the optically inactive form and we assumed that they may be artifacts or were formed without participation of enzymes in the plant cells.⁸

EXPERIMENTAL

Isolation : The extraction procedures for the roots of Marsh grapefruit and Hirado-Buntan are described in the previous paper.⁹ The CH₂Cl₂ eluate (43.11 g), obtained from silica gel column chromatography of the acetone extract of Marsh grapefruit, was separated with a silica gel column, centrifugal chromatography and finally repeated PTLC [solvents: CHCl₃-MeOH (19:1), acetone-hexane (1:1), benzene-MeOH (9:1)] to give citbismine-D (1)(2.5 mg) and citbismine-E (2)(1.4 mg). The CH₂Cl₂ eluate (29.394 g), obtained from the silica gel column chromatography of the acetone extract of Hirado-Buntan, was treated in a similar manner to give citbismine-E (2)(4.6 mg).

Citbismine-D (1): Pale yellow oil, $[\alpha]_D^{25} \pm 0^\circ$ (CHCl₃, c=0.140), ir $\nu_{\max}$ (CHCl₃) 1620, 1595 cm⁻¹; uv $\lambda_{\max}$ (EtOH) 224 (sh), 267, 298 (sh), 338, 387 nm; HRms m/z 722.2473 (M⁺, calcd for C₄₀H₃₈N₂O₁₁ 722.2476); EIms m/z 722 (M⁺), 663, 649 (base peak), 369, 353, 352, 338, 311; ¹H nmr (acetone-d₆) δ 15.84 (1H, s, 1'-OH), 14.82 (1H, s, 1-OH), 9.18 (1H, s, 6' or 6-OH), 8.62 (1H, s, 6 or 6'-OH), 8.04 (1H, d, J= 9.2 Hz, H-8'), 7.86 (1H, d, J= 8.5 Hz, H-8), 7.05 (1H, d, J= 9.2 Hz, H-7'), 6.81 (1H, d, J= 8.5 Hz, H-7), 6.58 (1H, d, J= 9.8 Hz, H-11'), 6.18 (1H, s, H-2), 5.70 (1H, d, J= 5.5 Hz, H-11), 5.51 (1H, d, J= 9.8 Hz, H-12'), 4.56 (1H, d, J= 5.5 Hz, H-12), 4.05 (1H, br s, 13-OH), 3.88 (3H, s, 5'-MeO), 3.77 (3H, s, 10-NMe), 3.64 (3H, s, 10'-NMe), 2.91 (3H, s, 5-MeO), 1.43, 1.38 (each 3H, s, 13-Me), 1.30 (6H, s, 13'-Me); NOE: irradiation at δ 3.77 (10-NMe) - 9.5% enhancement at δ 5.70 (H-11); irradiation at δ 3.64 (10'-NMe) - 9.8% enhancement at δ 6.58 (H-11'); irradiation at δ 5.70 (H-11) - 7.9% enhancement at δ 3.77 (10-NMe).

Citbismine-E (2): Pale yellow oil, $[\alpha]_D^{25} \pm 0^\circ$ (CHCl₃, c=0.120), ir $\nu_{\max}$ (CHCl₃) 1620, 1595 cm⁻¹; uv $\lambda_{\max}$ (EtOH) 221, 264, 275 (sh), 298 (sh), 335, 379 nm; HRms m/z 670.2160 (M⁺, calcd for C₃₆H₃₄N₂O₁₁ 670.2163); EIms m/z 670 (M⁺), 612, 611, 599, 598, 597, 582, 369, 351, 336, 311, 301 (base peak), 296, 287, 286; ¹H-nmr (acetone-d₆) δ 15.92 (1H, s, 1'-OH), 14.78 (1H, s, 1-OH), 9.3 (1H, br s, 6' or 6-OH), 8.7 (1H, br s, 6 or 6'-OH), 8.14 (1H, d, J= 9.2 Hz, H-8'), 7.86 (1H, d, J= 8.5 Hz, H-8), 7.03 (1H, d, J= 9.2 Hz, H-7'), 6.80 (1H, d, J= 8.5 Hz, H-7), 6.46 (1H, s, H-4'), 6.14 (1H, s, H-2), 5.72 (1H, d, J= 5.5 Hz, H-11), 4.54 (1H, d, J= 5.5 Hz, H-12), 4.02 (3H, s, 10'-NMe), 3.9 (1H, br s, 13-OH), 3.88 (3H, s, 5'-MeO), 3.78 (3H, s, 10-NMe), 3.57 (3H, s, 3'-MeO), 2.80 (3H, s, 5-MeO), 1.43, 1.38 (each 3H, s, 13-Me); NOE: irradiation at δ 4.02 (10'-NMe) - 12.4% enhancement at δ 6.46 (H-4'); irradiation at δ 3.78 (10-NMe) - 8.7% enhancement at δ 5.72 (H-11); irradiation at δ 3.57 (3'-MeO) - 10.8% enhancement at δ 6.46 (H-4'); irradiation at δ 5.72 (H-11) - 7.2% enhancement at δ 3.78 (10-NMe).

ACKNOWLEDGEMENT

The authors express their deep gratitude to Misses K. Suwa and S. Takeyama, Mukogawa Women's University, for ms and nmr spectra measurements.

REFERENCES AND NOTES

- 1 Part XXIV of "Constituents of Domestic Citrus Plants". Part XXIII: Y. Takemura, Y. Matsushita, N. Nagareya, M. Abe, J. Takaya, M. Ju-ichi, T. Hashimoto, Y. Kan, S. Takaoka, Y. Asakawa, M. Omura, C. Ito, and H. Furukawa, submitted to Chem. Pharm. Bull..
- 2 Y. Takemura, H. Uchida, M. Ju-ichi, M. Omura, C. Ito, K. Nakagawa, T. Ono, and H. Furukawa, Heterocycles 1992, 34, 2123 and references cited therein.
- 3 Y. Takemura, T. Nakata, M. Ju-ichi, M. Okano, N. Fukamiya, C. Ito, and H. Furukawa, Chem. Pharm. Bull., 1994, 42, 1213 and references cited therein.
- 4 Y. Takemura, M. Ju-ichi, M. Omura, M. Haruna, C. Ito, and H. Furukawa, Heterocycles, 1994, 38, 1937 and references cited therein.
- 5 Y. Takemura, M. Ju-ichi, T. Hashimoto, Y. Kan, S. Takaoka, Y. Asakawa, M. Omura, C. Ito, and H. Furukawa, Chem. Pharm. Bull., 1994, 42, 1548.
- 6 J. Reisch, K. Szendri, E. Minker, and I. Novak, Pharmazie, 1972, 27, 208.
- 7 S. Funayama and G. A. Cordell, J. Nat. Prod., 1985, 48, 536.
- 8 P. G. Watermann, "Alkaloids: Chemical and Biological Perspectives," Vol. 4, ed. by S. W. Pelletier, John Wiley & Sons Inc., NY, 1986, p 331.
- 9 Y. Takemura, J. Kuwahara, N. Nagareya, M. Ju-ichi, M. Omura, I. Kajiura, C. Ito, and H. Furukawa, Heterocycles, 1994, 39, 315.

Received, 27th January, 1995