

THE SYNTHESIS OF 1 β -FLUOROCARBAPENEM DERIVATIVE†



Masataka Ihara,^a Ken Satoh,^a Yohhei Ishida,^b Nobuaki Taniguchi,^a Yuji Tokunaga,^a Makoto Takemura,^b and Keiichiro Fukumoto*^a

^a Pharmaceutical Institute, Tohoku University, Aobayama, Sendai 980-77, Japan

^b Exploratory Research Laboratories I, Daiichi Pharmaceutical Co., Ltd., 16-13, Kitakasai 1-chome, Edogawa-ku, Tokyo 134, Japan


Abstract—The synthesis of 1 β -fluorocarbapenem derivative (16) was carried out starting with the 4-acetoxiazetidin-2-one (3) through the carbapenem formation utilizing the Sankyo's procedure.

Since the presence of a 1 β -methyl substituent such as 1 has been found to enhance the chemical and metabolic stability of synthetic carbapenem antibiotics,¹ the introduction of substituent(s), other than methyl group, at the C (1) position has been attracted by several groups.² We interested in the synthesis of 1 β -fluorocarbapenems (2), in order to study their biological activities and stabilities, and would like to describe our findings.³




With the aim of the preparation of 7, the substitution reaction of the commercially available 3 with fluoromalonate was examined. The desired fluoro compound (4) was obtained in 96% yield when the reaction was carried out in the presence of lithium hexamethyldisilazide (LHMDS). One of the methoxycarbonyl groups of 4 was removed by hydrolysis of 4 using one equivalent mole of LiOH·H₂O followed by decarboxylation of the resulting carboxylic acid (5). Heating 5 in xylene at 135 °C produced a 2 : 1 diastereoisomeric mixture of 6 in 63% overall yield from 4. The corresponding carboxylic acids (7) were obtained in 96% yield by hydrolysis of 6 with LiOH·H₂O.

† Dedicated to the memory of the late Professor Yoshio Ban.



Scheme 1


The construction of the carbapenem skeleton was first studied by the application of the Merck's method.⁴ Although the side-chain extension to a β -keto ester was rather difficult, the transformation of **7** into **8** was performed in 44% yield, when MeCN was used as the reaction solvent. After the deprotection of the *tert*-butyldimethylsilyl (TBDMS) group with 6N HCl in MeOH, followed by diazoexchange with *p*-toluenesulfonyl azide in the presence of Et₃N, the formed **10** was subjected to the carbene insertion reaction using Rh₂(OAc)₄ in AcOEt at 80 °C. Reactions proceeded in reasonable yields, but the products (**11**) were too unstable to isolate as pure forms. Furthermore, activation of the C(2) position of **11** for the introduction of a thiol group failed. Therefore, we next focused our attention on the Sankyo's method⁵ for the assembly of the carbapenem nucleus.



Scheme 2

After conversion of **6** into thioester (**12**) (87% overall yield), reaction of **12** with *p*-nitrobenzyl chlorooxalate in the presence of *i*Pr₂NEt and CaCO₃ provided the oxalimides (**13**) in 86% yield. Cyclization of **13** was achieved

by heating with $(\text{EtO})_3\text{P}$ in refluxing xylene for 1 h. Two stereoisomers (**14**) and (**15**) were isolated in 45 and 33% yields, respectively, after chromatographical separation. Stereostructures of products were determined by the nOe experiment: 9.3% nOe was observed between hydrogens at C(1) and C(5) positions. The assigned stereochemistry was further supported by their chemical behaviors. Treatment of **14** with $n\text{Bu}_4\text{NF}$ in the presence of AcOH produced **16** in 31% yield; 64% yield based on the consumed starting material (**14**), while



Scheme 3

the deprotection of TBDMS group of **15** to **17** was not successful under various conditions. It was reported that 1 α -methylcarbapenems were unstable comparing with the corresponding 1 β -isomers.⁶ The rule could be applied to this case. Unfortunately, attempts to deblock the *p*-nitrobenzyl ester resulted in failure. Only the pyrrole derivative (**20**) was obtained by deprotection procedures carried out under a variety of conditions. It is assumed that the formed acid (**18**) is readily attacked with H₂O at the lactam carbonyl group to expel the fluorine atom to afford **20** via **19**. This result suggests the necessity of further elaboration for isolation of deprotected 1-fluorocarbapenems.

EXPERIMENTAL

(3S,4S)-3-[(1'R)-1'-(*tert*-Butyldimethylsiloxy)ethyl]-4-bis(methoxycarbonyl)fluoromethylazetidin-2-one (4). To a stirred solution of dimethyl fluoromalonate (5.0 g, 33.3 mmol) in dry THF (50 ml) at -78°C was added 1 M LHMDs-hexane (31.8 ml, 31.8 mmol). After being stirred for 30 min at -78°C , a solution of **3** (8.7 g, 30.3 mmol) in dry THF (90 ml) was added to it. The mixture was stirred for 3 h at -78°C and for 9 h at ambient temperature. After neutralization with 1N HCl under ice cooling, the mixture was concentrated under reduced pressure and the residue was taken up into AcOEt. The extract was washed with H₂O and brine, dried (Na₂SO₄), and evaporated. Chromatography of the residue on silica gel with CHCl₃-MeOH (99 : 1 v/v) as eluent gave **4** (11.0 g, 96%) as crystals, mp $158-160^{\circ}\text{C}$, $[\alpha]_{\text{D}}^{29} -47.7^{\circ}$ ($c = 1.1$, CHCl₃); ν_{max} (CHCl₃, cm⁻¹) 3440 (NH), 1755 (C=O); δ_{H} (CDCl₃, 500 MHz) 0.07 (6H, s, SiMe₂), 0.88 (9H, s, ^tBu), 1.06 (3H, d, $J = 6.1$ Hz, 1'-Me), 3.29 (1H, br s, 3-H), 3.87 and 3.88 (each 3H, each s, 2 $\times$ OMe), 4.25-4.30 (1H, m, 1'-H), 4.41 (1H, dd, $J = 20.8$ and 1.8 Hz, 4-H), 5.85 (1H, br s, NH); HRms m/z calcd for C₁₅H₂₅NO₆FSi (M⁺ - Me): 362.1436, found: 362.1411. Anal. Calcd for C₁₆H₂₈NO₆FSi: C, 50.91; H, 7.48; N, 3.71. Found: C, 50.99; H, 7.52; N, 3.69.

(3S,4S)-3-[(1'R)-1'-(*tert*-Butyldimethylsiloxy)ethyl]-4-methoxycarbonyl(fluoro)methylazetidin-2-one (6). To a solution of **4** (90 mg, 0.24 mmol) in MeOH (5 ml) was added a solution of LiOH·H₂O (10 mg, 0.24 mmol) in H₂O (2 ml), and the mixture was stirred for 2 h at ambient temperature. After concentration under reduced pressure, the residue was partitioned between Et₂O (15 mL) and H₂O (5 ml). The ethereal layer was washed with saturated NaHCO₃ (3 ml). The combined aqueous layer was acidified with 1N HCl under ice cooling and then thoroughly extracted with AcOEt. The extract was washed with brine, dried (MgSO₄), and evaporated to give a 2 : 1 diastereoisomeric mixture of the acids (**5**) (71 mg, 82%) as a powder: δ_{H} (CDCl₃, 500 MHz) 0.05 and 0.07 [each 3H, each s, SiMe₂], 0.87 and 0.88 [9H (2 : 1), each s, ^tBu], 1.07 and 1.13 [3H (1 : 2), each d, each $J = 6.1$ Hz, 1'-Me], 3.33 and 3.38 [1H (1 : 2), each br s, 3-H], 3.88 and 3.89 [3H (2 : 1), each s, OMe], 4.24-4.30 (1H, m, 1'-H), 4.44 and 4.46 [1H (2 : 1), each dd, each $J = 19.0$ and 1.8 Hz, 4-H], 6.45 and 6.93 [1H (2 : 1), each s, NH], which was used in the next reaction without purification. The starting material (**4**) (16 mg) was obtained from the above ethereal solution.

A mixture of the above acids (**5**) (100 mg, 0.28 mmol) in xylene (5 ml) was heated for 3 h at 135°C . After evaporation of the solvent, the residue was taken up into CH₂Cl₂. The extract was washed with brine, dried

(MgSO₄), and evaporated to afford a residue, which was subjected to flash chromatography on silica gel. Elution with hexane–AcOEt (7 : 3 v/v) yielded a 2 : 1 diastereoisomeric mixture of **6** (68 mg, 77%) as a solid, mp 136–138 °C; $\nu_{\max}$ (CHCl₃, cm⁻¹) 3405 (NH), 1764 (C=O); δ_{H} (CDCl₃, 400 MHz) 0.06 and 0.07 (each 3H, each s, SiMe₂), 0.86 (9H, s, ^tBu), 1.13 and 1.15 [3H (2 : 1), each d, each $J = 6.4$ Hz, 1'-Me], 3.23 and 3.26 [1H (1 : 2), each br s, 3-H], 3.83 and 3.84 [3H (1 : 2), each s, OMe], 4.00–4.15 (1H, m, 4-H), 4.20–4.30 (1H, m, 1'-H), 4.91 (1/3 $\times$ 1H, dd, $J = 48.3$ and 5.6 Hz, CHF), 4.97 (2/3 $\times$ 1H, dd, $J = 48.3$ and 4.9 Hz, CHF), 5.85 and 5.90 [1H (2 : 1), each br s, NH]; HR ms m/z calcd for C₁₃H₂₃NO₄FSi (M⁺ – Me): 304.1381, found: 304.1378. Anal. Calcd for C₁₄H₂₆NO₄FSi: C, 52.64; H, 8.20; N, 4.38. Found: C, 52.63; H, 8.18; N, 4.24.

(3S,4S)-3-[(1'R)-1'-(*tert*-Butyldimethylsiloxy)ethyl]-4-[1-carboxy-1-fluoromethyl]azetidin-2-one (7). To a solution of **6** (62 mg, 0.19 mmol) in MeOH (3 ml) was added a solution of LiOH·H₂O (8.1 mg, 0.19 mmol) in H₂O (2 ml), and the mixture was stirred for 1 h at ambient temperature. After concentration under reduced pressure, the residue was partitioned between Et₂O (10 ml) and H₂O (4 ml). The ethereal layer was extracted with saturated NaHCO₃. The combined aqueous layer was acidified with 1N HCl under ice cooling and then thoroughly extracted with AcOEt. The extract was washed with brine, dried (MgSO₄), and evaporated to provide a 2 : 1 mixture of **7** (57 mg, 96%) as an amorphous powder: $\nu_{\max}$ (CHCl₃, cm⁻¹) 3405 (NH), 1760 (C=O); δ_{H} (CDCl₃, 500 MHz) 0.10 and 0.11 (2/3 $\times$ 6H, each s, SiMe₂), 0.11 and 0.12 (1/3 $\times$ 6H, each s, SiMe₂), 0.90 and 0.91 [9H (2 : 1), each s, ^tBu], 1.18 (2/3 $\times$ 3H, $J = 6.7$ Hz, 1'-Me), 1.24 (1/3 $\times$ 3H, $J = 6.1$ Hz, 1'-Me), 3.22–3.24 and 3.24–3.26 [1H (2 : 1), each m, 3-H], 4.10–4.17 (1H, m, 4-H), 4.25–4.32 (1H, m, 1'-H), 5.15 (1/3 $\times$ 1H, dd, $J = 48.8$ and 3.1 Hz, CHF), 5.25 (2/3 $\times$ 1H, dd, $J = 48.8$ and 4.9 Hz, CHF), 7.30 and 7.49 [1H (2 : 1), each br s, NH]; HRms m/z calcd for C₁₂H₂₁NO₄FSi (M⁺ – Me): 290.1224, found: 290.1244.

(3S,4S)-3-[(1'R)-1'-(*tert*-Butyldimethylsiloxy)ethyl]-4-[(1-fluoro-3-(*p*-nitrobenzyloxy)-carbonyl)-2-oxopropyl]azetidin-2-one (8). To a solution of **7** (367 mg, 1.2 mmol) in MeCN (10 ml) was added a solution of *N,N*-carbonyldiimidazole (234 mg, 1.44 mmol) in MeCN (5 ml), and the mixture was stirred for 30 min at ambient temperature. After addition of a solution of magnesium *p*-nitrobenzyl malonate (661 mg, 1.32 mmol) in MeCN (5 ml), the resulting mixture was stirred for 12 h at ambient temperature. Concentration of the mixture under reduced pressure gave a residue, which was taken up into AcOEt. The organic extract was washed with 10% HCl, H₂O, saturated NaHCO₃, and brine. After being dried (MgSO₄), evaporation of the solvent afforded a residue, which was subjected to flash chromatography on silica gel. Elution with benzene–acetone (92 : 8 v/v) provided **8** (253 mg, 44%) as an oil: $\nu_{\max}$ (CHCl₃, cm⁻¹) 3400 (NH), 1765 (C=O), 1520 and 1350 (NO₂); δ_{H} (CDCl₃, 500 MHz) 0.06 (6H, s, SiMe₂), 0.86 (9H, s, ^tBu), 1.15 and 1.18 [3H (2 : 1), each d, each $J = 6.1$ Hz, 1'-Me], 3.18–3.23 (1H, m, 3-H), 3.69–3.87 (2H, m, COCH₂CO), 4.05–4.10 (1H, m, 4-H), 4.20–4.27 (1H, m, 1'-H), 4.90 and 4.98 [1H (2 : 1), each dd, each $J = 47.6$ and 6.1 Hz, CHF], 5.28 and 5.31 [2H (2 : 1), each s, CH₂Ar], 5.90 and 5.97 [1H (2 : 1), each br s, NH], 7.52 and 7.53 (2H, each d, $J = 8.6$ Hz, 2 $\times$ ArH), 8.23 and 8.24 (2H, each d, $J = 8.6$ Hz, 2 $\times$ ArH); HRms m/z calcd for C₁₈H₂₂N₂O₇FSi (M⁺ – ^tBu): 425.1180, found: 425.1153.

(3*S*,4*S*)-3-[(1*R'*)-1'-(*tert*-Butyldimethylsiloxy)ethyl]-4-[1-fluoro-1-(phenylthiocarbonyl)-methyl]azetidin-2-one (12). The esters (**6**) (735 mg, 2.3 mmol) was converted, by the same procedure as above, into the acids (**7**), which was used in the reaction without purification. To a stirred solution of the crude acids (**7**), thiophenol (0.236 ml, 2.30 mmol) and 1-hydroxybenzotriazole (HOBt) (31 mg, 0.23 mmol) in THF (30 ml) was added DCC (712 mg, 3.45 mmol) under ice cooling. The mixture was stirred for 20 min at 0 °C and for 18 h at ambient temperature. Evaporation of the solvent gave a residue, which was taken up into AcOEt. After filtration, the filtrate was washed with H₂O, dried (Na₂SO₄), and evaporated to afford a residue, which was chromatographed on silica gel. Elution with hexane–AcOEt (85 : 15 v/v) provided **12** (792 mg, 87% from **6**) as an oil: $\nu_{\max}$ (CHCl₃, cm⁻¹) 3405 (NH), 1770 and 1690 (C=O); δ_{H} (CDCl₃, 400 MHz) 0.07 and 0.08 [6H (2 : 1), each s, SiMe₂], 0.86 and 0.87 [9H (2 : 1), each s, ^{*t*}Bu], 1.16 and 1.17 [3H (1 : 2), each d, each *J* = 5.9 Hz, 1'-Me], 3.25–3.35 (1H, m, 3-H), 4.10–4.20 (1H, m, 4-H), 4.23–4.30 (1H, m, 1'-H), 5.04 and 5.07 [1H (1 : 2), each dd, each *J* = 48.3 and 4.9 Hz, CHF], 5.95 and 6.00 [1H (2 : 1), each br s, NH], 7.38–7.48 (5H, m, 5 × ArH); HR ms *m/z* calcd for C₁₈H₂₅NO₃FSSi (M⁺ – Me): 382.1308, found; 382.1309.

***p*-Nitrobenzyl (1*R* and 1*S*,5*S*,6*S*)-6-[(1'*R*)-1'-(*tert*-Butyldimethylsiloxy)ethyl]-1-fluoro-2-(phenylthio)carbapen-2-em-3-carboxylate (14 and 15).** To a solution of **12** (792 mg, 1.99 mmol) in CH₂Cl₂ (40 ml) were added ClCOCO₂PNB (970 mg, 3.98 mmol), ^{*i*}Pr₂NEt (0.69 ml, 3.98 mmol) and CaCO₃ (399 mg, 3.98 mmol) under ice cooling, and the mixture was stirred for 1 h at the same temperature. After evaporation, the residue was taken up into AcOEt. The extract was washed with H₂O, dried (MgSO₄), and evaporated to give a residue, which was purified by flash chromatography on silica gel. Elution with CHCl₃–MeOH (99 : 1 v/v) provided **13** (1.04 g, 86%) as an oil.

A mixture of **13** (1.04 g, 1.72 mmol) and (EtO)₃P (1.47 ml, 8.57 mmol) in xylene (50 ml) was heated for 1 h under reflux. Evaporation of the solvent afforded a residue which was subjected to flash chromatography on silica gel. Elution with hexane–AcOEt (95 : 5 v/v) gave **14** (438 mg, 45%) as an oil: $\nu_{\max}$ (CHCl₃, cm⁻¹) 1790 and 1720 (C=O), 1520 and 1340 (NO₂); δ_{H} (CDCl₃, 400 MHz) 0.05 and 0.06 (each 3H, each s, SiMe₂), 0.84 (9H, s, ^{*t*}Bu), 1.13 (3H, d, *J* = 6.4 Hz, 1'-Me), 3.46 (1H, t, *J* = 2.9 Hz, 6-H), 4.19 (1H, ddd, *J* = 29.3, 6.4 and 2.9 Hz, 5-H), 4.30–4.40 (1H, m, 1'-H), 5.03 (1H, dd, *J* = 54.2 and 6.4 Hz, 1-H), 5.37 and 5.50 (each 1H, each d, *J* = 14.2 Hz, OCH₂Ar), 7.35–7.60 (3H, m, 3 × ArH), 7.62 (2H, d, *J* = 6.8 Hz, 2 × ArH), 7.68 (2H, d, *J* = 8.8 Hz, 2 × ArH), 8.23 (2H, d, *J* = 8.8 Hz, 2 × ArH); HRms *m/z* calcd for C₂₈H₃₃N₂O₆FSSi (M⁺): 572.1813, found: 572.1799.

Further elution with hexane–AcOEt (9 : 1 v/v) furnished **15** (322 mg, 33%) as an oil: δ_{H} (CDCl₃, 400 MHz) 0.05 and 0.06 (each 3H, each s, SiMe₂), 0.84 (9H, s, ^{*t*}Bu), 1.18 (3H, d, *J* = 6.3 Hz, 1'-Me), 3.18 (1H, t, *J* = 3.9 Hz, 6-H), 4.17 (1H, dt, *J* = 33.7 and 3.9 Hz, 5-H), 4.20–4.30 (1H, m, 1'-H), 5.34 and 5.49 (each 1H, each d, *J* = 14.2 Hz, OCH₂Ar), 5.58 (1H, dd, *J* = 54.2 and 3.9 Hz, 1-H), 7.30–7.45 (3H, m, 3 × ArH), 7.61 (2H, d, *J* = 7.8 Hz, 2 × ArH), 7.67 (2H, d, *J* = 8.6 Hz, 2 × ArH), 8.22 (2H, d, *J* = 8.6 Hz, 2 × ArH).

***p*-Nitrobenzyl (1*R*,5*S*,6*S*)-6-[(1'*R*)-1'-Hydroxyethyl]-1-fluoro-2-(phenylthio)carbapen-2-em-3-carboxylate (16).** A mixture of **14** (236 mg, 0.42 mmol), AcOH (0.24 ml, 4.10 mmol) and 1 M Bu₄NF-THF (1.24 ml, 1.24 mmol) in THF (20 ml) was stirred for 24 h at ambient temperature. After concentration

under reduced pressure, the residue was subjected to flash chromatography on silica gel. Elution with CHCl_3 -MeOH (99 : 1 v/v) gave **14** (122 mg) and further elution with CHCl_3 -MeOH (99 : 1 v/v) provided **16** (59 mg, 31%; 64% based on the consumed **14**) as an oil: $[\alpha]_{\text{D}}^{21} +145.6^\circ$ ($c = 0.49$, CHCl_3); ν_{max} (CHCl_3 , cm^{-1}) 1790 and 1710 (C=O), 1520 and 1350 (NO_2); δ_{H} (CHCl_3 , 400 MHz) 1.29 (3H, d, $J = 6.3$ Hz, 1'-Me), 3.49 (1H, dd, $J = 5.4$ and 2.9 Hz, 6-H), 4.12 (1H, ddd, $J = 28.3$, 6.4 and 2.9 Hz, 5-H), 4.25–4.40 (1H, m, 1'-H), 5.06 (1H, dd, $J = 54.2$ and 6.4 Hz, 1-H), 5.34 and 5.56 (each 1H, each d, $J = 13.7$ Hz, OCH_2Ar), 7.40–7.50 (3H, m, $3 \times \text{ArH}$), 7.61 (2H, d, $J = 7.3$ Hz, $2 \times \text{ArH}$), 7.69 (2H, d, $J = 8.6$ Hz, $2 \times \text{ArH}$), 8.24 (2H, d, $J = 8.6$ Hz, $2 \times \text{ArH}$); HRms m/z calcd for $\text{C}_{22}\text{H}_{19}\text{NO}_4\text{FS}$ ($\text{M}^+ - \text{NO}_2$): 412.1019, found 412.0996.

(1'S,2'R)-5-(1'-Carboxy-2'-hydroxypropyl)-3-(phenylthio)pyrrole-2-carboxylic Acid (20). A mixture of **16** (53 mg, 0.11 mmol) and PtO_2 (10 mg) in THF-1/15 M phosphate buffer (pH 7) (1 : 1 v/v, 10 ml) was shaken for 30 min at ambient temperature under H_2 (4 kg/cm²). After filtration, followed by concentration of the filtrate under reduced pressure, the residue was purified by chromatography using DIAION HP-20 with H_2O and THF- H_2O (5 : 95 v/v) as eluent. Elution with H_2O provided **20** (15 mg, 41%) as a powder: δ_{H} (D_2O , 400 MHz) 1.20 (3H, d, $J = 6.4$ Hz, Me), 3.36 (1H, d, $J = 8.3$ Hz, 1'-H), 4.18 (1H, dq, $J = 8.3$ and 6.4 Hz, 2'-H), 5.90 (1H, s, 4-H), 7.25 - 7.55 (5H, m, Ph).

ACKNOWLEDGMENT

This work was partly supported by Daiichi Pharmaceutical Co., Ltd.

REFERENCES

1. D. H. Shih, F. Baker, L. Cama, and B. G. Christensen, *Heterocycles*, 1984, **21**, 29.
2. N. V. Shah and L. D. Cama, *Heterocycles*, 1987, **25**, 221; Y. Nagao, T. Abe, H. Shimizu, T. Kumagai, and Y. Inoue, *J. Chem. Soc., Chem. Commun.*, 1989, 821; R. L. Rosati, L. V. Kapili, P. Morrissey, and J. A. Retsema, *J. Med. Chem.*, 1990, **33**, 291; J. H. Bateson, A. M. Robins, and R. Southgate, *J. Chem. Soc., Perkin Trans. 1*, 1991, 2399; Y. Sendo, M. Kii, M. Sakanoue, K. Motokawa, and Y. Kimura, *Chem. Pharm. Bull.*, 1992, **40**, 2410; T. Shibata, K. Iino, T. Tanaka, T. Hashimoto, Y. Kameyama, and Y. Sugimura, *Tetrahedron Letters*, 1985, **26**, 4739.
3. A part of this work (synthesis of **7**) has been published as a preliminary form: M. Ihara, K. Satoh, N. Taniguchi, K. Fukumoto, Y. Ishida, and M. Takemura, *Heterocycles*, 1993, **35**, 139.
4. T. N. Salzmann, R. W. Ratcliffe, B. G. Christensen, and F. A. Bouffard, *J. Am. Chem. Soc.*, 1980, **102**, 6161; R. W. Ratcliffe, T. N. Salzmann, and B. G. Christensen, *Tetrahedron Lett.*, 1980, **21**, 31.
5. A. Yoshida, Y. Tajima, N. Takeda, and S. Oida, *Tetrahedron Lett.*, 1984, **25**, 2793.
6. D. H. Shih, L. Cama, and B. G. Christensen, *Tetrahedron Lett.*, 1985, **26**, 587.

Received, 28th April, 1995