


2-METHYL-4-NITROISOXAZOLIN-5-ONE:
RING TRANSFORMATION TO 3-NITROPYRROLES

Masahiro Ariga,* Nagatoshi Nishiwaki, Yuko Miwa, Keita Tani, and Yasuo Tohda

Department of Chemistry, Osaka Kyoiku University,
Asahigaoka, Kashiwara, Osaka 582, Japan


Abstract - Ring transformation of 2-methyl-4-nitroisoxazolin-5-one with some enolate anions afforded 3-nitropyrroles. A ring-opened intermediate of the ring transformation was isolated.

In our course of the study on electron deficient pyridones, 4-nitroisoxazolin-5-one was obtained.¹ Functional varieties of the isoxazolone such as heterodiene, α -nitrolactone, and β -nitroenamine suggest us that it may undergo various types of reactions. We already reported 2-methyl-4-nitroisoxazolin-5-one (**1**) acted as a precursor of a nitrile oxide in the reaction with various dipolarophiles to give isoxazoles.² Availability of 2-acetyl-4-nitroisoxazolin-5-one (**2**) as an acetylating reagent also was revealed.³



Here, we wish to deal a new ring transformation of **1** with enolate anions of β -keto esters and β -diketones. Treatment of **1** with 1.2 equiv. of ethyl 2-sodio-3-oxobutanoate in pyridine at 70 °C gave ethyl 1,3-dimethyl-4-nitro-2-pyrrolicarboxylate (**3a**)⁴ as a major product (77.0%) and a trace of diethyl 1,3,5-trimethyl-2,4-pyrroledicarboxylate (**4a**).⁵ The structure of **3a** was assigned by its ¹H nmr and ir spectra, and by comparison with those of an authentic sample, which was independently synthesized from ethyl 3-methylpyrrole-2-carboxylate⁶ by nitration, separation from 5-nitro isomer on silica gel column, and

N-methylation. The similar reaction with diethyl 2-sodio-3-oxopentanedioate yielded ethyl 3-ethoxycarbonylmethyl-1-methyl-4-nitro-2-pyrrolicarboxylate (**3b**).⁷ The results are summarized in Table 1.



It is clear that the C(2)-C(3) part of the obtained pyrrole (**3**) is derived from the reagent and the rest arises from the N(2)-C(3)-C(4) moiety of the isoxazolone (**1**). Formation of **4a** must involve the condensation of two molar ethyl 3-oxobutanoate during the ring transformation, but the detail is in progress.


Table 1 Ring transformation of 2-methyl-4-nitroisoxazoloin-5-one (**1**)

Substrate	Reagent ^{a)}	Reaction Conditions				Products (Yield/%)	
		equiv.	Solvent	Temp./°C	Time/h		
1	a	1.2	Pyridine	70	5	3a (77.0)	4a (trace)
1	a	3.0	"	"	"	3a (48.0)	4a (36.0)
1	a	1.2	DMF	"	"	3a (56.8)	
1	b	1.2	"	"	"	3b (56.6)	
1	c	1.2	Pyridine	20	"	3c (5.8)	5c (10.4)
1	c	3.0	DMF	70	"		5c (64.2)
5c	(NH ₄ Cl)	3.0	Ethanol	80	10	3c (83.7)	


a) **a**: ethyl 2-sodio-3-oxobutanoate; **b**: diethyl 2-sodio-3-oxo-pentanedioate;
c: 3-sodiopentane-2,4-dione

2,4-Pentanedione, a kind of β -diketone, yielded a similar ring transformed product (**3c**)⁸ in a low yield, and gave **5c**⁹ as a major product. The ring-opened product (**5c**) presented in an equilibrium mixture of *E*- and *Z*-isomers in CDCl₃ solution in a ratio of about 4:3.^{10,11}

The product (**5c**) is a β -nitroenamine and has an electrophilic carbonyl carbon in the molecule. Cyclization of **5c** to pyrrole (**3c**) could be achieved in good yield by heating **5c** at 80 °C for 10 h in ethanol containing a small amount of ammonium chloride as an acid catalyst.




In the present reaction, a new C-N bond is formed between the active methylene carbon of the reagent and the ring nitrogen of isoxazolone (**1**). Taking account of isolation of **5c**, one of a plausible course of the reaction is proposed as that direct nucleophilic attack of the enolate anion to the nitrogen atom of **1** accompanied by decarboxylation to form **6**, then intermediate anion (**6**) recycled to nitropyrrole (**3**).



We, however, can not neglect that the intramolecular attack of the enolate anion introduced at the 3-position of **1** to the ring nitrogen atom caused the N-O bond cleavage and decarboxylation to give aziridine (**7**), and successive ring opening of the aziridine ring forms intermediate (**6**).

Since, it is not likely known that the C-anion of the reagent attacks directly at the ring nitrogen of an isoxazolone before ring cleavage. It has been well known that the nucleophilic attack at the 3-position of 5-isoxazolone resulted in the N-O bond cleavage and decarboxylation.¹²⁻¹⁵ There are also some precedents for the ring opening of isoxazole or isoxazolone followed by succeeding recyclization to an aziridine.¹⁶



Though more examples with a variety of carbonyl compounds are necessary to show the general utility of the reaction, the ring transformation furnishes a convenient route for synthesis of functionalized β -nitropyrrole derivatives which were tediously obtainable by direct nitration.

In conclusion, this new ring transformation suggests that 2-methyl-4-nitroisoxazol-5(2H)-one (**1**) behaves as a masked β -nitroenamine besides a precursor of nitrile oxide.²

REFERENCES

1. M. Ariga, Y. Tohda, H. Nakashima, K. Tani, Y. Mori, and E. Matsumura, *Bull. Chem. Soc. Jpn.*, 1991, **64**, 3544.
2. S. Higashida, H. Nakashima, Y. Tohda, K. Tani, N. Nishiwaki, and M. Ariga, *Heterocycles*, 1992, **34**, 1511.
3. N. Nishiwaki, Y. Takada, Y. Tohda, and M. Ariga, *Chem. Express*, 1992, **7**, 845.
4. Mp 143.0-145.0 °C. Ir: ν 3140, 1690, 1550, 1340, 1100. $^1\text{H Nmr}$ (CDCl_3): δ 1.42 (t, $J=7$ Hz, 3H), 2.64 (s, 3H), 3.94 (s, 3H), 4.36 (q, $J=7$ Hz, 2H), 7.60 (s, 1H).
5. Ir: ν 1690, 1540, 1350. $^1\text{H Nmr}$ (CDCl_3): δ 1.37 (t, $J=7$ Hz, 3H), 1.38 (t, $J=7$ Hz, 3H), 2.51 (s, 3H), 2.55 (s, 3H), 3.78 (s, 3H), 4.28 (q, $J=7$ Hz, 2H), 4.30 (q, $J=7$ Hz, 2H).
6. W. G. Terry, A. H. Jackson, G. W. Kenner, and G. Kornis, *J. Chem. Soc.*, **1965**, 4389.
7. Mp 88.0-89.0 °C. Ir: ν 3130, 1720, 1700, 1550, 1346. $^1\text{H Nmr}$ (CDCl_3): δ 1.27 (t, $J=7$ Hz, 3H), 1.37 (t, $J=7$ Hz, 3H), 3.96 (s, 3H), 4.18 (q, $J=7$ Hz, 2H), 4.28 (s, 2H), 4.34 (q, $J=7$ Hz, 2H), 7.64 (s, 1H).
8. Mp 133.5-134.5 °C. Ir: ν 1655, 1550, 1310. $^1\text{H Nmr}$ (CDCl_3): δ 2.52 (s, 3H), 2.69 (s, 3H), 3.90 (s, 3H), 7.63 (s, 1H).
9. Mp 125.0-128.0 °C. Ir: ν 1610, 1560, 1310. $^1\text{H Nmr}$ (CDCl_3): *E-5c*: δ 2.10 (s, 6H), 3.29 (s, 3H), 6.46 (d, $J=11$ Hz, 1H), 8.16 (d, $J=11$ Hz, 1H), 15.71 (s, 1H); *Z-5c*: δ 2.10 (s, 6H), 3.03 (s, 3H), 6.72 (d, $J=11$ Hz, 1H), 7.92 (d, $J=11$ Hz, 1H), 15.53 (s, 1H).
10. N. Nishiwaki, E. Wakamura, Y. Nishida, Y. Tohda, and M. Ariga, *Heterocyclic Commun.*, 1996, **2**, in press.
11. S. Rajappa and K. Nagarajan, *J. Chem. Soc., Perkin Trans. II*, **1978**, 912.
12. F. de Sarlo and G. Renzi, *Tetrahedron*, 1966, **22**, 2995.
13. D. J. Woodman, C. H. Borman, N. Tontapanish, and P. M. Stonebraker, *J. Org. Chem.*, 1969, **34**, 2981.
14. H. J. Wollweber and C. Wentrup, *J. Org. Chem.*, 1985, **50**, 2041.
15. N. Nishiwaki, Y. Takada, Y. Inoue, Y. Tohda, and M. Ariga, *J. Heterocycl. Chem.*, 1995, **32**, 473.
16. H. C. van der Plas, "Ring Transformations of Heterocycles", Vol. 1, Chapt. 3, Academic Press, New York, 1973.